

USN

**10MAT11**

First Semester B.E. Degree Examination, June / July 2014
Engineering Mathematics - I

4

Time: 3 hrs.

Max. Marks: 100

a

Note: 1. Answer any FIVE full questions, choosing at least two from each part.

2. Answer all objective type questions only on OMR sheet page 5 of the answer booklet.

3. Answer to objective type questions on sheets other than OMR will not be valued.

b-c

PART — A

1 a. Choose the correct answers for the following : (04 Marks)

i) If $y_n = (4177)^n e^{4n} \cos x + n \tan^{-1} \frac{1}{4}$ then $y =$ _____A) $e^4 x \cos x$ B) $e^{2n} \sin 3x$ C) $ex \cos x$ D) None of theseii) $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots$ is,

A) Taylor's series B) Exponential series C) Maclaurin's series D) None of these

iii) In the Rolle's theorem if $F'(c) = 0$ then the tangent at the point $x = c$ is,

iv) A) parallel to y-axis B) parallel to x-axis C) parallel to both axes D) None of these

iv) If $y = 3^x$ then $y' =$ _____A) $(\log x)3^x$ B) $3(\log x)x^2$ C) $3^x \log 3$ D) $3^x (\log_e 3)^x$ b. If $x = \sin t$, $y = \sin pt$ prove that, $(1 - x^2)y_{n+2} - (2n+1)xy_{n+1} + (p^2 - n^2)y_n = 0$. (04 Marks)c. State and prove Cauchy's mean value theorem in $[0, 16]$. (06 Marks)d. Expand $41 + \sin 2x$ by using Maclaurin's expansion. (06 Marks)

III. MAT111P

2 a. Choose the correct answers for the following : (04 Marks)

i) The value of $\lim_{x \rightarrow 0} \frac{\sin x}{x}$ is,A) 0 B) 1 C) $\frac{1}{e}$ D) 0.9ii) The angle between two curves $r = ae^{\theta}$ and $r = b e^{p\theta}$ is,A) $\frac{\pi}{2}$ B) $\frac{\pi}{4}$ C) 0 D) π iii) $\int \frac{dx}{dt} \left(\frac{dy}{dt} \right)^2$

A) Polar form B) Parametric form C) Cartesian form D) None of these

$$\int \frac{\log x}{x^2} \cot x \, dx$$

A) 1 B) 0 C) 2 D) -2

b. Find a & b, if $\lim_{x \rightarrow 0} \frac{x(1 + \cos x) - bx \sin x}{x} = 1$. (04 Marks)c. Find the pedal equation of the curve $r^2 = a^2 \cos 2\theta$ (06 Marks)d. Find the radius of curvature at any point t of the curve $x = a(t + \sin t)$ and $y = a(1 - \cos t)$. (06 Marks)

3 a. Choose the correct answer

(04 Marks)

i) If $u = (x - y)^2 + (y - z)^2 + (z - x)^2$ then $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z}$ is,

- A) 1 B) 24 C) $2(x + y + z)$ D) 0

ii) $ex \cos y = \pi [1 + (x-1)(y - \frac{\pi}{4}) + \frac{(x-1)^2}{2}(x-1)(y - \frac{\pi}{4}) + \dots]$

- A) (1, i) B) (0, 0) C) (1, 1) D) (4, 1)

iii) At (a, b) $\frac{\partial^2 u}{\partial x^2} = A$, $\frac{\partial^2 u}{\partial y^2} = B$ and $\frac{\partial^2 u}{\partial x \partial y} = H$ and if $AB - H^2 < 0$ the point is called,

- A) Maximum B) Minimum C) Saddle 17) Extremum

iv) If $J = \frac{a(u, v)}{a(x, y)}$, $J = \frac{a(x, y)}{a(u, v)}$, then DI is,

- A) 0 B) 2 C) D) 1

b. If $u = f(\frac{x}{y}, \frac{y}{z}, \frac{z}{x})$ then prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 0$

(04 Marks)

c. If $u = \frac{xy}{z}$, $v = \frac{yz}{x}$, $w = \frac{zx}{y}$ then show that $J \begin{vmatrix} u & v & w \\ \frac{\partial u}{\partial x} & \frac{\partial v}{\partial x} & \frac{\partial w}{\partial x} \\ \frac{\partial u}{\partial y} & \frac{\partial v}{\partial y} & \frac{\partial w}{\partial y} \end{vmatrix} = 4$ verify $JJ = 1$.

(06 Marks)

d. For the kinetic energy $E = \frac{1}{2}mv^2$ find approximately the change in E as the mass m changes from 49 to 49.5 and the velocity v changes from 1600 to 1590.

(06 Marks)

4 a. Choose the correct answers for the following:

(04 Marks)

i) The value of $V \times V^2$ is,

- A) 0 B) R C) 9 D) 3

ii) Any motion in which the curl of the velocity vector is zero, then the vector v is said to be,

- A) Constant B) Solenoidal C) Vector D) Irrotational

iii) In orthogonal curvilinear co-ordinates the Jacobian $J = \frac{a(x, y, z)}{a(u, v, w)}$ is,

- A) $\frac{h_1}{h_2 h_3}$ B) $\frac{1}{h_1 h_2 h_3}$ C) $h_1 h_2 h_3$ D) —

iv) A gradient of the scalar point function ϕ , $V\phi$ is,

- A) Scalar function B) Vector function C) ϕ D) zero

b. Find the value of the constant a such that the vector field,

$F = (axy - z^3)i + (a - 2)x^2j + (1 - a)xz^2k$ is irrotational and hence find a scalar function ϕ such that $F = V\phi$.

(04 Marks)

C. Prove that $\text{curl}(\text{curl } A) = V(V.A) - V^2 A$.

(06 Marks)

d. Express $V^2 \phi$ in orthogonal curvilinear co-ordinates.

(06 Marks)

5 a. Choose the correct answers for the following :

(04 Marks)

- i) The value of $\int_{0}^{\pi} \cos^3(4x) dx$ is,

A) 3

B) 6

C)

D) 2

- ii) If the equation of the curve remains unchanged after changing 0 to -0 the curve $r = f(\theta)$ is symmetrical about,

- A) A line perpendicular to initial line through pole
- B) Radially symmetric about the point pole.
- C) Symmetry does not exist
- D) Initial line

volume of the curve $r = a(1 + \cos\theta)$ about the initial line is,A) $4na^3$ B) $2na^3$

3

C) $8na^3$ D) $\frac{4a^3}{3}$

- iv) The asymptote for the curve $x^3 + y^3 = 3axy$ is equal to,

- A) $x+y=0$
- B) $x-y-a=0$
- C) No Asymptote
- D) $x+y-a=0$

- b. Evaluate $\int_0^{2\pi} \frac{\log(1 + \cos x)}{\cos x} dx$

(04 Marks)

- c. Evaluate $\int_0^{\pi} \sqrt{2ax - x^2} dx$.

(06 Marks)

- d. Find the area of surface of revolution about x-axis of the astroid $x + y^3 = a^3$. (06 Marks)

6 a. Choose the correct answers for the following : (04 Marks)

- i) In the homogeneous differential equation, $\frac{dy}{dx} = \frac{f_1(xy)}{f_2(xy)}$ the degree of the function, $f_1(xy)$ and $f_2(xy)$ are,

- A) Different
- B) Relatively prime
- C) Same
- D) None of these

- ii) The integrating factor of the differential equation, $\frac{dy}{dx} + \cot xy = \cos x$ is,

- A) $x \sin x$
- B) $\sin x$
- C) $-\sin x$
- D) $\cot x$

- iii) Replacing $\frac{dy}{dx} = -\frac{dy}{dx}$ in the differential equation $f(x, y, \frac{dy}{dx}) = 0$ we get the differential equation of,

- A) Polar trajectory
- B) Orthogonal trajectory
- C) Parametric trajectory
- D) Parallel trajectory.

- iv) Two families of curves are said to be orthogonal if every member of either family cuts each member of the other family at,

- A) Zero angle
- B) Right angle
- C) $\frac{\pi}{6}$
- D) $\frac{27c}{3}$

- b. Solve $(1 + e^{xy})dx + e^{xy}(1 - \frac{dy}{y}) = 0$. (04 Marks)

- c. Solve $\frac{dy}{dx} + x \sin 2y = x^3 \cos^2 y$. (06 Marks)

- d. Find the orthogonal trajectories of $r^2 = a^2 \cos^2 \theta$. (06 Marks)

7 a. Choose the correct answer

$$\text{I) } A = \begin{vmatrix} 7 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 7 \end{vmatrix}$$

- is called,
- A) Scalar matrix B) Diagonal matrix C) Identity matrix D) None of these
 - ii) If $r = n$ and $x = y = z = 0$. The equations have only _____ solution.
 A) Non trivial B) Trivial C) Unique D) Infinite
 - iii) In Gauss Jordan method, the coefficient matrix can be reduced to,
 A) Echelon form B) Unit matrix C) Triangular form D) Diagonal matrix
 - iv) The inverse square matrix A is given by,

A) A^{-1}

B) $\frac{\text{adj}A}{|A|}$

C) $\text{adj}A$

: $\text{adj}A$

b. Find the Rank of the matrix, $\begin{vmatrix} 1 & 2 & 3 & 2 \\ 2 & 3 & 5 & 1 \\ 1 & 3 & 4 & 5 \end{vmatrix}$.

- c. Investigate the values of 2 and $.i$ such that the system of equations, $x + y + z = 6$, $x + 2y + 3z = 10$, $x + 2y + Xz = g$ may be i) Unique solution ii) Infinite solution iii) No solution.

- d. Using Gauss elimination method solve,

$$2x_1 - x_2 + 3x_3 = 1, \quad -3x_1 - 4x_2 - 5x_3 = 0, \quad x_2 - 5x_3 = 0$$

8 a. Choose the correct answers for the following :

- i) A square matrix A of order 3 has 3 linearly independent eigen vectors then a matrix P can be found such that $P^{-1}AP$ is a,
 - A) Diagonal matrix
 - B) Unit matrix
 - C) Singular matrix
 - D) Symmetric matrix
- ii) The eigen values of matrix $\begin{bmatrix} 2 & 1 & 0 \\ 4 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ are,
 - A) $2 \pm i$
 - B) $2 \pm \sqrt{3}$
 - C) $2 \pm \sqrt{5}$
 - D) None of these
- iii) Solving the equations $x + 2y + 3z = 0$, $3x + 4y + 4z = 0$, $7x + 10y + 12z = 0$. x, y and z values are,
 - A) $x=y=z=0$
 - B) $x=y=z=1$
 - C) $x = y = z$
 - D) None of these
- iv) The index and significance of the quadratic form, $x^2 + 2x_2^2 - 3x_3^2$; are respectively _____ and _____
 - A) Index = 1, Signature = 1
 - B) Index = 1, Signature = 2
 - C) Index = 2, Signature = 1
 - D) None of these.

- b. Find all the eigen values and the corresponding eigen vectors of the matrix,

$$A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & 4 \\ 2 & 4 & 3 \end{bmatrix}$$

- c. Reduce the matrix $A = \begin{bmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & \frac{6}{5}y^2 - 3xz^2 \end{bmatrix}$ into a diagonal matrix.

- d. Reduce the quadratic form $3x^2 + 2yz + 2zx - 2xy$ to the canonical form.
