

# www.FirstRanker.com

www.FirstRanker.com

ł.

USN

1

# Third Semester B.E. Degree Examination, Dee.24<sup>1</sup>04ftn.2020 **Basic Thermodynamics**

3 BIGS SIGHEME

Time: 3 hrs.

Max. Marks: 100

**18M E33** 

Note: I. Answer any FIVE full questions, choosing ONE full question from each module.

2. Use of thermodynamic data hand book and steam tables is permitted.

3. Assume missing data suitably-

# Module-1

a. Differentiate between micro and macroscopic approach.

- b. Define the following terms with neat sketch:
  - (i) Open system
  - (ii) Closed system
  - (iii) Isolated system
  - (iv) Quasi-static process

c. The temperature 'T' on a thermometric scale is defined as T = al n(K) + b, where a and b are constants. The values of K are found to be 1.83 and 6.78 at 0°C and 100°C, respectively. Calculate the temperature for value of K = 2.42. (08 Marks)

OR

- 2 a. Define:
  - (i) Thermodynamic equilibrium
  - (ii) Zeroth law of thermodynamics
  - b. With neat sketch explain the working principle of: (i) Electrical resistance thermometer (ii) Thermocouple .6

c. Two Celsius thermometer 'A' and 'B' agree at ice point and steam point, and related by the , where L, M and N are constants. When both thermometers are equation  $t = L + Mt_B + ...$ immersed in a fluid, 'A' registers 26°C, while 'B' registers 25°C. Determine the reading of `A' when 'B' reads 37.4°C. (08 Marks)

# Module-2

- a. Define thermodynamic work and heat. 3
  - b. Write an expression for displacement of work for the following process with P-V diagrams.
    - Constant pressure (i)
    - (ii) Constant volume
    - (iii) Constant temperature
    - (iv) Polytropic process
  - c. A quantity of gas is compressed in a piston-cylinder from a volume of  $0.8611 \text{ m}^{-3}$  to a final volume of 0.1721 m<sup>3</sup>. The pressure in (bar) and as a function of volume (m<sup>3</sup>) is given by:

$$P^{-0.8611\ 8.6067\ x\ 10^{-5}}$$

(i) Find the amount of work done in KJ.

(ii) If the atmospheric pressure is I bar, acting on the other side of piston is considered. Find the net work done in KJ. (08 \larks)

# www.FirstRanker.com

(08 Marks)

(08 Marks)

(04 Marks)

(04 Marks)

### (08 Marks)

(04 Marks)



www.FirstRanker.com

18M

# OR

- 4 a. State I <sup>s</sup> law of thermodynamics. Derive an expression for 1 <sup>st</sup> law of thermodynamics for open system (SFEE). (10 Marks)
  - b. The working fluid, in a steady flow process at a rate of 220 kg/min. The fluid rejects 100 KJ/s of heat passing through the system. The condition of the fluid at inlet and outlet are given as  $V_{c}$ . 220 m/s,  $p_i = 6.0$  bar,  $u_i = 2000$  KJ/kg, ui = 0.36 m<sup>3</sup>/kg and  $p_i = 1.2$  bar,  $V_i = 140$  m/s,  $u_i = 1400$  kJ/kg,  $0_i = 1.3$  m<sup>3</sup>/kg. The suffix 1 and 2 indicates at inlet and outlet conditions respectively. Determine the power capacity of the system in MW.

(10 Marks)

(10 Marks)<sup>-</sup>

(04 Marks)

(08 Marks)

(06 Marks)

### Module-3

- 5 a. Define the following terms:
  - (i) Thermal reservoir
  - (ii) Heat engine
  - (iii) Kelvin-Plank statement of 2<sup>-1</sup> law
  - (iv) Clausius statement of r<sup>i</sup> law
  - (v) Heat pump
  - b. A heat engine working on a Carnot cycle absorbs heat from three thermal reservoirs at 1000 K, 800 K and 600 K, respectively. The engine does 10 KW of net work and rejects 400 kJ/min of heat to a heat sink at 300 K. If the heat supplied by the reservoir at 1000 K is 60% of heat supplied by the reservoir at 600K. Find the quantity of heat supplied by each reservoirs.

6 a. Define entropy and prove that it is a point function.

- b. Discuss the Clausius Inequality.
  - c. A steel ball mass of 10 kg at 627°C is dropped in 100 kg of oil at 30°C. The specific heat of steel and oil are 0.5 kJ/kgK and 3.5 kJ/kgK, respectively. Calculate the entropy change of steel, oil and the universe.

### Module-4

7 a. With neat sketch, explain available and Unavailable energy on T-S diagram (06 Marks,

- b. Explain the concept of second law of efficiency.
- C. A Carnot engine works between the temperature limits 225°C and 25°C in which water is used as the working fluid. If heat is supplied to the saturated liquid at 225°C, until it is converted into saturated vapour, determine per kg of water.
  - (i) Amount of heat absorbed by the fluid
  - (ii) Available energy
  - (iii) Unavailable energy

(Take latent heat of water = 1858.5 klikg)

(08 Marks)

### OR

- 8 a. With neat sketch explain the working of separating and throttling calorimeter. (10 Marks)
  - A vessel of volume 0.04 m<sup>3</sup> contains a mixture of saturated water and saturated state at a temperature of 250°C. The mass of the liquid present is 9 kg. Find the mass, specific volume, enthalpy, entropy and internal energy of the steam. (10 Marks)

# www.FirstRanker.com

FirstRanker.com

www.FirstRanker.com

www.FirstRanker.com



# Module-5

## 9 a. Define:

- (i) Mole fraction
- (ii) Mass fraction
- (iii) Dalton's law
- (iv) Amgat's law of volume additives
- b. A mixture of gases contain 1 kg of CO2 and 1.5 kg of N2. The pressure and temperature of the mixture are 3.5 bar and 27°C. Determine:
  - (i) Mole fraction of each constituent
  - (ii) Partial pressure
  - (iii) Partial volume
  - (iv) Volume of mixture
  - (v) Density of mixture

### OR

- 10 a. State and explain the following terms:
  - (i) Compressibility factor
  - (ii) Reduced properties
  - (iii) Real gases
  - (iv) Relative humidity
  - b. With usual notations, write the Vandeer Waal equation and explain the terms involved in it.
  - C. Determine the pressure exerted by CO<sup>•</sup> in a container of 1.5 m<sup>3</sup> capacity when it contains 5 kg at 27°C:
    - (i) Using ideal gas relation
    - (ii) Using Vandeer Waal's equation
    - [Take a = 364.3 kPa (m<sup>3</sup>lkg.mol)<sup>2</sup>: b = 0.0427 (m<sup>3</sup>/kg.mol) for Vandeer Waal's constants] (08 Marks)

18ME33

(10 Marks)

(10 Marks)

(08 Marks)