

Set No. 1
Code No: R10102 / R10
I B.Tech I Semester Regular/Supplementary Examinations January 2012
MATHEMATICS - I
(Common to all branches)
Time: 3 hours
Max Marks: 75
**Answer any FIVE Questions
All Questions carry equal marks**

- 1.(a) Find the differential equations of all parabolas with x-axis as its axis and $(\alpha, 0)$ as its focus.
(b) Find the orthogonal trajectories of coaxial circles $x^2 + y^2 + 2\lambda y + c = 2$, where λ is the parameter. [7M + 8M]
- 2.(a) Solve $(D^2 - 2)y = e^{\sqrt{x}} + \cos x$
(b) Solve $\frac{d^2 y}{dx^2} + 4\frac{dy}{dx} + 5y = 2 \sin hx$ subject to $y=0$ and $\frac{dy}{dx} = 1$ at $x=0$. [7M + 8M]
- 3.(a) If $u = xy + yz + zx$, $v = x^2 + y^2 + z^2$ and $w = x + y + z$, verify whether there exists a possible relationship in between u, v and w. If so find the relation.
(b) Find the minimum value of $x^2 + y^2 + z^2$ on the plane $x + y + z = 3a$ [7M + 8M]
- 4.(a) Trace the curve $x(x_2 + y_2) = 4(x^2 - y^2)$
(b) Trace the polar curve $r = 2 + 3 \cos \theta$. [7M + 8M]
- 5.(a) Find the perimeter of one loop of the curve $3ay^2 = x(x - a)^2$.
(b) Find the volume generated by revolving the area bounded by one loop of the curve $r = a(1 + \cos \theta)$ about the initial line. [7M + 8M]
- 6.(a) Evaluate $\int_0^\infty \int_x^\infty \frac{e^{-y}}{y} dx dy$ by changing the order of integration.
(b) Evaluate $\int_0^2 \int_0^{\sqrt{2x-x^2}} \frac{x}{\sqrt{x^2 + y^2}} dy dx$ by changing into polar coordinates. [7M + 8M]
- 7.(a) Find the directional derivative of $\phi(x, y, z) = xy^2 + yz^3$ at the point $(2, -1, 1)$ in the direction of the vector $i + 2j + 2k$.
(b) Find $\text{curl}[rf(r)]$ where $r = xi + yj + zk$, $r = |r|$ [7M + 8M]

firstranker.com

**Code No: R10102 / R10****Set No. 1**

- 8.(a) Compute the line integral $\int (y^2 dx - x^2 dy)$ round the triangle whose vertices are (1,0), (0,1) and (-1,0) in the xy-plane.
- (b) Evaluate the integral $I = \int \int_S x^3 dydz + x^2 y dzdx + x^2 z dxdy$ using divergence theorem, where S is the surface consisting of the cylinder $x^2 + y^2 = a^2$ ($0 \leq z \leq b$) and the circular disks $z=0$ and $z=b$ ($x^2 + y^2 \leq a^2$).

[7M + 8M]

Set No. 2

Code No: R10102 / R10
I B.Tech I Semester Regular/Supplementary Examinations January 2012
MATHEMATICS - I
(Common to all branches)
Time: 3 hours
Max Marks: 75

Answer any FIVE Questions
All Questions carry equal marks

- 1.(a) Find the solution of the differential equation $\frac{dy}{dx} = x e^{y-x^2}$ and $y(0) = 0$.
- (b) A body initially at 80°C cools down to 50°C in 10 minutes, the temperature of the air being 40°C . What will be the temperature of the body after 20 minutes?
[7M + 8M]
- $\frac{d^2y}{dx^2} + 9y = e^{2x} x^2$
- 2.(a) Solve dx^2
- (b) Find the general solution of $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = e^x \sin 2x$
[7M + 8M]
- 3.(a) Verify whether the functions $u = \sin^{-1} x + \sin^{-1} y$ and $v = x\sqrt{1-y^2} + y\sqrt{1-x^2}$ are functionally dependent. If so, find the relation between them. [7 M+8 M]
- (b) Prove that the rectangular solid of maximum volume that can be inscribed into a sphere of radius 'a' is a cube.
[7M + 8M]
- 4.(a) Trace the parametric curve $x = a \left(\cos \theta + \frac{1}{2} \log \tan^2 \left(\frac{t}{2} \right) \right)$ and $y = a \sin t$.
- (b) Trace the lemniscate $r^2 = a^2 \cos 2\theta$.
[7M + 8M]
- 5.(a) Find the surface area generated by revolving the arc of the curve $y = a \cosh (x/c)$ from $x=0$ to $x=c$ about the x-axis.
- (b) Find the total length of the lamniscate $r^2 = a^2 \cos 2\theta$.
[7M + 8M]
- 6.(a) Find the area of the region which is outside the circle $r=1$ and inside the cardioid $r = (1 + \cos \theta)$
- (b) Evaluate the following integral by changing into polar coordinates

$$\int \int \sqrt{\frac{1-(x^2+y^2)}{1+x^2+y^2}} dx dy$$
 over the positive coordinate of the circle $x^2 + y^2 = 1$
[7M + 8M]

**Code No: R10102 / R10****Set No. 2**

- 7.(a) Find the directional derivative of the divergence of $F = x y i + y z j + z^2 k$ at the point $(2,1,2)$ in the direction of the outer normal to the sphere $x^2 + y^2 + z^2 = 9$.
- (b) Find the value of a, b and c such that $(x + y + a z) i + (b x + 2 y - z) j + (-x + c y + 2 z) k$ is irrotational.
- [7M + 8M]
- 8.(a) If $f = (x^2 + y - 4) i + 3 x y j + (2 x z + z^2) k$ and S is the upper half of the sphere $x^2 + y^2 + z^2 = 16$. Show by using Stokes theorem that $\int_S \text{Curl } f \cdot n \, ds = 2\pi a^3$.
- (b) If S is the surface of the tetrahedron bounded by the planes $x = 0$, $y = 0$, $z = 0$ and $ax + by + cz = 1$. Show that $\int_S r \cdot n \, ds = \frac{1}{2abc}$.
- [7M + 8M]

Set No. 3

Code No: R10102 / R10
I B.Tech I Semester Regular/Supplementary Examinations January 2012
MATHEMATICS - I
(Common to all branches)
Time: 3 hours
Max Marks: 75
**Answer any FIVE Questions
All Questions carry equal marks**

$$(x^2 + y^2) \frac{dy}{dx} = xy$$

1.(a) Solve

(b) A colony of bacteria is grown under ideal condition in laboratory so that the population increases exponentially with time. At the end of 3 hours there are 10000 bacteria. At the end of 5 hours there are 40000. How many bacteria were present initially?

[7M + 8M]

 2.(a) Solve $(D^3 - 6D^2 + 11D - 6)y = e^{-2x} + x^3$

 (b) Solve $(D^2 + 1)y = x^2 e^{2x} + x \cos x$

[7M + 8M]

 3.(a) If $u = x + y + z$, $u^2 v = y + z$ and $u^3 w = z$, then find $\frac{\partial(u, v, w)}{\partial(x, y, z)}$

 (b) Find the minimum and maximum distances of a point on the curve $2x^2 + 4xy + 4y^2 - 8 = 0$.

[7M + 8M]

 4.(a) Trace the parametric curve $x = a(t - \sin t)$ and $y = a(1 + \cos t)$

 (b) Trace the curve $y^2(x - a) = x^2(x + a)$ and $a > 0$

[7M +

 8M] 5.(a) Find the volume of the solid formed by revolving the area bounded by the curve $27ay^2 = 4(x - 2a)^3$ about x-axis

 (b) Find the length of the loop of the curve $r = a(1 - \cos \theta)$.

[7M + 8M]

 6.(a) Find the area of the loop of the curve $x^3 + y^3 = 3axy$, by transforming it into polar coordinates.

 (b) Change the order of integration and evaluate $I = \int_0^1 \int_x^{\sqrt{x}} xy \, dy \, dx$.

[7M + 8M]

firstranker.com

**Code No: R10102 / R10****Set No. 3**

- 7.(a) In what direction from the point $(1, 3, 2)$ is the directional derivative of $\phi = 2xz - y^2$ is maximum and what is its magnitude.
- (b) Show that $\vec{F} = (y^2 \cos x + z^3)i + (2y \sin x - 4)j + (3xz^2 + 2)k$ is a conservative force field and find its scalar potential. [7M + 8M]
- 8.(a) Show that $\vec{F} = (2xy + z^3)i + x^2j + 3xz^2k$ is a conservative force field. Find the scalar potential and the work done in moving an object in this field from $(1, -2, 1)$ to $(3, 1, 4)$.
- (b) Verify Green's theorem, if $\oint_C Mdx + Ndy$ is $(xy + y^2)dx + x^2dy$ with C : closed curve of the region bounded by $y = x$ and $y = x_2$. [7M + 8M]

Set No. 4

Code No: R10102 / R10
I B.Tech I Semester Regular/Supplementary Examinations January 2012
MATHEMATICS - I
(Common to all branches)
Time: 3 hours
Max Marks: 75
**Answer any FIVE Questions
All Questions carry equal marks**

$$x \frac{dy}{dx} - y = \sqrt{x^2 + y^2}$$

1.(a) Solve

 (b) A body is heated to 110°C is placed in air at 10°C . After 1 hour its temperature is 80°C . When will the temperature be 30°C ?

[7M + 8M]

 2.(a) Solve $(D^2 + 3D + 2)y = \sin x \sin 2x$

 (b) Solve $(D^2 + 2D - 3)y = x^3 e^{-2x}$.

[7M + 8M]

 3.(a) Verify whether the functions $u = \frac{x-y}{x+z}$ and $v = \frac{x+z}{y+z}$ are functionally dependent. If so, find the relation in between them.

 (b) The temperature T at any point (x, y, z) in the space is given as $T = 400 x^2 y z$. Find the highest temperature on the surface of the sphere $x^2 + y^2 + z^2 = 1$

[7M + 8M].

 4.(a) Trace the curve $x^3 + y^3 = 3a x y$

 (b) Trace the polar curve $r = a(1 - \sin \theta)$.

[7M + 8M]

 5 (a) Find the surface area generated by revolving the arc $x^{2/3} + y^{2/3} = a^{2/3}$ about x-axis.

 (b) Find the volume of the solid generated by revolving the cardioid $r = a(1 + \cos \theta)$ about the initial line.

[7M + 8M]

 6.(a) Find the area of a plate in the form of a quadrant of an ellipse $x^2/a^2 + y^2/b^2 = 1$ by changing into polar coordinates.

 (b) By changing the order of integration, evaluate the integral $\int_0^{4a} \int_{\frac{y^2}{4a}}^{\sqrt{4a-y}} dx dy$.

[7M + 8M]

**Code No: R10102 / R10****Set No. 4**

- 7.(a) Find the constants a and b so that the surface $a x^2 - b y z = (a + 2) x$ will be orthogonal to the surface $4 x^2 y + z^3 = 4$ at the point $(1, -1, 2)$.
- (b) Determine the constant b such that $\vec{A} = (b x^2 y + y z) \vec{i} + (x y^2 - x z^2) \vec{j} + (2 x y z - 2 x^2 y^2) \vec{k}$ has zero divergence.
- [7M + 8M]
- 8.(a) Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F} = x^2 \vec{i} + y^2 \vec{j}$ and curve c is the arc of the parabola $y = x^2$ in the xy -plane from $(0,0)$ to $(1,1)$.
- (b) Evaluate by Stokes theorem $\int_C (x + y) dx + (2x - z) dy + (y + z) dz$, where C is the boundary of the triangle vertices $(0,0,0)$, $(1,0,0)$ and $(1,1,0)$.
- [7M + 8M]