

Total No. of Pages : 02

B.Tech.(Petroleum Refinery Engineering) (2013 Batch) (Sem.-4)

Subject Code : BTPC-404/BTCH-305

Time : 3 Hrs.

Max. Marks : 60

1. **SECTION-A is COMPULSORY consisting of TEN questions carrying TWO marks each.**
2. **SECTION-B contains FIVE questions carrying FIVE marks each and students have to attempt any FOUR questions.**
3. **SECTION-C contains THREE questions carrying TEN marks each and students have to attempt any TWO questions.**

1. Answer briefly :

- Find the density of nitrogen gas at NTP.
- State 2nd law of thermodynamics.
- What is throttling process? Give an example.
- What is the significance of H-x diagram?
- What do you understand by dew point and bubble point?
- What is the physical significance of partial molar properties?
- Define equilibrium constant. What is its significance?
- What is the effect of pressure on equilibrium constant in a gas phase reaction?
- Write two major applications of Gibb's Duhem equation.
- What do you understand by theoretical flame temperature?

2. Calculate ΔU and ΔH in kJ for 1 kmol water, as it is vaporized at a constant temperature of 373 K and constant pressure of 101.3 kPa. The specific volumes of liquid and vapour at these conditions are 1.04×10^{-3} and $1.675 \text{ m}^3/\text{kmol}$ respectively ; 1030 kJ of heat is added to water for this change.

- Write the Clapeyron equation and hence find the melting point of mercury at 10 bar, where mercury has a density of $13.69 \times 10^3 \text{ kg/m}^3$ in the liquid state and $14.193 \times 10^3 \text{ kg/m}^3$ in the solid state, both measured at the melting point of 234.33 K and 1 bar. Heat of fusion of mercury is 9.7876 kJ/kg.
- Find the volume of *n*-pentane at 500 K and 20 bar following Van der waals equation of state. $T_c = 469.6 \text{ K}$, $P_c = 33.7 \text{ bar}$
- State Hess's law of constant heat summation. Calculate heat of formation of the gaseous ethyl alcohol at 298 K using following data :
 Standard heat of formation of $\text{CO}_2 (\text{g}) = -393.51 \text{ kJ/mol}$
 Standard heat of formation of $\text{H}_2\text{O} (\text{l}) = -285.83 \text{ kJ/mol}$
 Heat of combustion of gaseous ethyl alcohol at 298 K = -1410.09 kJ/mol
- Derive the expression for effect of temperature on fugacity coefficient.

SECTION-C

- Show that the fugacity of a gas obeying the van der waals equation of state is given by $\ln f = \frac{b}{V-b} - \frac{2a}{RTV} + \ln \frac{RT}{V-b}$, where *a* and *b* are van der Waals constants.
- Describe the criteria for chemical reaction equilibria. Calculate the equilibrium constant for the reaction $\text{N}_2 + 3 \text{H}_2 \rightarrow 2 \text{NH}_3$ at 500 K assuming that the standard heat of reaction is constant in temp. range 298 K to 500 K. Standard heat of formation and standard free energy of formation of NH_3 at 298 K are -46100 J/mol and -16500 J/mol respectively.
- For a binary system of components (1) and (2), the vapour pressures are given by the Antoine's equations

$$\ln P_1^{\text{sat}} = 13.818 - \frac{2477.07}{T - 40.00} \quad ; \quad \ln P_2^{\text{sat}} = 13.859 - \frac{2911.32}{T - 56.56}$$

P is in kPa and *T* in K. Assume the solution as ideal. Calculate :

- The composition of liquid and vapour in equilibrium at 95 kPa and 335 K.
- The composition of the vapour in equilibrium with a liquid containing 40% (mol) pentane and the equilibrium temperature at *P* = 95 kPa.
- The total pressure and the vapour composition in equilibrium with a liquid of composition $x_1 = 0.40$ at *T* = 333.2 K.

NOTE : Disclosure of Identity by writing Mobile No. or Making of passing request on any page of Answer Sheet will lead to UMC against the Student.