


Learning Objectives

- Introduction
- Gross features
- Bronchial Tree
- Microscopic features of different part
- Functional correlations
- Clinical correlates


Introduction

- Provide molecular oxygen for cellular oxidation
- Remove carbon dioxide generated
- Vascular or circulatory transport AND Air/Gas transport


Gross Anatomy




Bronchopulmonary Segments



Continued.....



Continued.....



Functional Requirements

- The Respiratory System has tubular components for air/gas conduction
- Plus respiratory components for gas exchange.
- Inspiratory pressure tends to collapse the conducting tubes. They must be held open.
- Other essential functions include: warming; moistening; cleansing by removal of particulate matter; detoxification by absorption of harmful gases; and entrapment of harmful bacteria and viruses.


Conduction

- The nasal cavity is held open by bone and cartilage.
- Most of the additional functions are carried out by **pseudostratified columnar** ciliated epithelial cells and **goblet cells (mucous secretors)**.
- The cilia beat rhythmically in one direction only
- Move debris and pathogen-laden mucous to the oropharynx and mouth where it is expectorated or swallowed.


Continued.....

- Air temperature control is by the **profuse capillary beds** that are beneath the epithelium
- Which warm or cool inspired air.
- Extensive venous plexus, largely replace capillaries in the nasal cavities to modify air temperature

Trachea



Bronchus and Bronchial Tree




- The trachea typifies the conducting system.
- The cilia are paralyzed by cigarette smoke.
- At rest, smooth muscle contraction decreases tracheal (and bronchial) diameter to decrease respiratory dead space.

Continued.....


- Bronchioles - lumen of **1mm or less in diameter**.
- The epithelium **ciliated columnar**.
- Cartilage and glands **disappear**
- Bronchiole is held open by the surrounding lung tissue.
- The smooth muscle in the wall may excessively narrow the lumen in **asthma**.

Bronchiole




- Bronchioles with a lumen of **1 mm or less** in diameter.
- Epithelium becomes ciliated columnar.
- Cartilage and glands disappear
- The bronchiole is held open by the surrounding lung tissue.
- The smooth muscle may excessively narrow the lumen in asthma.

Bronchiole




- Bronchioles continue to divide and decrease in size
- Terminal Bronchioles
- Respiratory Bronchioles , which give the alveolar ducts, alveolar sacs and alveoli.

Terminal and Respiratory Bronchioles



- Epithelium - **non-ciliated cuboidal** and **goblet cells disappear**.
- Gas exchange - in the respiratory alveoli
- That bud from the respiratory bronchioles.


Alveoli




- The alveoli - functional unit
- Out-pocketings of respiratory bronchioles, alveolar ducts and alveolar sacs.
- The most conspicuous feature of the alveolar wall - Presence of many small capillaries in the septae between them.
- Opposite the septae the alveoli are open to allow air entry.

Epithelium

surface.



- Type I are the **flattened epithelial cells** (squamous pneumocytes) that have cytoplasmic extensions about 0.1 - 0.3 μm thick and have no microvilli.
- In **some areas** the basal lamina of the type I cell is **fused** to the capillary
- Endothelium - possible areas of gas exchange.
- Type II, the **large alveolar cells (granular pneumocytes)** are **cuboidal** without broad cytoplasmic extensions and occupy about 5% of the alveolar surface.



Epithelium

- Type II Cells produce Surfactant, a stabilizing, surface-active material consisting mainly of the phospholipid lecithin.
- Surfactant is crucial to lung function
- Its **high viscosity** and **low surface tension**, stabilizes the diameter of alveoli preventing collapse after expiration, while retaining residual air in the alveoli after each breath;
- Because alveoli are already partially open, energy expenditure in subsequent inspirations is much less.


Continued....

- Acting as a **detergent** surfactant facilitates the removal of amniotic fluid from the lungs at birth, before the first breath.
- **Dust cells** (macrophages) usually lie free in the alveoli and air passageways.
- Their role is to clear the lungs of invading bacteria and inspired particulate material (e.g. dust and carbon particles).

Summary


- Massive surface area for gas exchange (**75 square metres**)
- As lumen decreases so do cartilage, cilia, epithelial height and mucous
- Smooth muscle controls diameter, but excessively narrowing in asthma
- Surfactant is thick in polycystic fibrosis

Emphysema



Bronchiectasis

Bronchiectasis



MCQs

Question :- Which of the following is correct regarding Clara cell?

- a. Ciliated secretory cuboidal cells
- b. Present in larger terminal bronchiole
- c. Present in smaller terminal bronchiole
- d. Number decreases in small bronchiole