

Refraction - II

Department of Ophthalmology

2

Acknowledgement

- Photographs in this presentation are courtesy of Kanski's Clinical Ophthalmology.

Learning Objectives

At the end of the class, students shall be able to

- Understand what is refraction.
- Have basic knowledge of hypermetropia and astigmatism and their management.

What is Refraction

- When rays of light traveling through air enter a denser transparent medium, the speed of the light is reduced and the light rays proceed at a different angle, i.e., they are refracted.
- Except when the rays are normal

Refraction in Ophthalmology

- **Methods for evaluating the optical and refractive state of the eye**

Emmetropia


- Parallel light rays, from an object more than 6 m away, are focused at the plane of the retina when accommodation is at rest.
- Clear image of a distant object formed without any internal adjustment of the optics of the eye.
- Absence of emmetropia = Ametropia

Refractive errors

Anomalies of the optical state of the eye

- **Myopia**
- **Hypermetropia**
- **Astigmatism**

Hypermetropia



Hypermetropia

- Refractive or Diopteric state of eye wherein **incident parallel rays of light coming from infinity are focused behind the retina** with accommodation being at rest.
- Near images can be blurred unless there is sufficient accommodation, as in a child.
- They have blurred images for distant objects also
- Most children are born about +3 D hyperopic, but this usually resolves by age 12 years.


Types

|
|
|
|
|

Axial Curvature Index Positional Absence of lens



- Axial is the commonest form.
- In this condition the total refractive power of eye is normal but there is axial shortening of eye wall.



- Each millimeter of shortening represents approximately 3D of refractive change and thus a hypermetropia of over 6D is uncommon.
- Physiological: Infant, child.
- Pathological: Orbital tumour, or inflammatory mass may indent the posterior pole of the eye and flatten it

- **Curvature Hypermetropia** : When the radius of curvature of any of the refracting surfaces is increased,
- congenitally (cornea plana) or as a result of trauma
- Increase of 1 mm produces a hypermetropia of 6 D.
- **Index Hypermetropia** : Usually manifests itself as a decrease in the effective refractivity of the lens and is responsible for the hypermetropia which occurs physiologically in old age and pathologically in diabetes.

- **Positional Hypermetropia** : Posterior placed lens also produced hypermetropia whether it occurs as a congenital anomaly or as a result of trauma and disease.
- **Aphakia** : Surgical, posterior dislocation of lens

Clinical Types:

- **Simple Hypermetropia** : Commonest form.
- It results from normal biological variations in the development of eye e.g., axial and curvatural.
- **Pathological Hypermetropia** : Either congenital or acquired conditions of eyeball which are outside the normal biological variation of development e.g.
 - index , positional (Aphakia).
- **Functional Hypermetropia** : Results from paralysis of accommodation as seen in patients with third nerve palsy.

Components of hypermetropia

Total hypermetropia = Latent+manifest (facultative + absolute)

Accommodation in Hypermetropia

- **Contraction of the ciliary muscle** in the act of accommodation increases the refractive power of the lens so that it corrects a certain amount of hypermetropia.
- Normally there is an appreciable amount corrected by the contraction involved in the **physiological tone** of this muscle.
- Consequently the full degree of hypermetropia is revealed only when this muscle is paralysed by the use of a drug such as atropine.
- This is called **latent** hypermetropia, normally 1D.


Manifest Hypermetropia consists of:

- **Facultative Hypermetropia:** Corrected by the effort of accommodation
- **Absolute Hypermetropia:** Cannot be overcome by effort of accommodation
- As tone of ciliary muscle decreases with age, some latent hypermetropia becomes manifest
- As range of accommodation reduces with age, more facultative hypermetropia becomes absolute, all of it after age 60.

Symptoms

- Vary with degree of hypermetropia and accommodative effort
- Blurred vision: near > distant
- Accommodative asthenopia
- Convergent squint due to continuous effort of accommodation, excess of convergence leads to dissociation of muscle balance
- Early onset of presbyopia

Signs



- Small eyeball
- Smaller cornea
- Shallow anterior chamber predisposes to angle closure glaucoma since size of lens is normal
- Apparent divergent squint

- **Retina** : Have peculiar sheen : a reflex effect so called “shot silk retina” on ophthalmoscopic finding.
- **Optic disc** : Characteristic appearance which may resemble an optic neuritis (Pseudopapillitis).

Treatment


- In young children below the age of 6-7 years, some degree of hypermetropia is physiological and a correction need be given only if the error is high or if strabismus is present.
- In those between 6 and 16 years especially when they are working strenuously at school smaller error may require correction.
- Required in middle aged patient, in high hypermetropia and if patient is having symptoms
- Optical:
 - Glasses
 - Contact lens

- Convex lenses prescribed after full cycloplegic refraction, particularly in children
- Child with convergent squint may need “full atropine correction”
- Contact lens power is a little more than spectacle power

Surgical


- **Conductive keratoplasty.**
- Non contact Holmium YAG laser **thermokeratoplasty** for lower hypermetropia (+1D – 2.5 D).
- **Phakic Intraocular lens** (+6D – +10 D)

Astigmatism



Astigmatism

- Astigmatism is a type of refractive error where in the refraction varies in the different meridia.
- Consequently the ray of light entering in the eye cannot converge to a point focus but form focal lines.



Astigmatism

- Light rays passing through a steep meridian are deflected more than those passing through a flatter meridian.

1. Corneal Astigmatism e.g. keratoconus
2. Lenticular Astigmatism
 - (i) Curvatural – e.g. lenticonus
 - (ii) Positional – subluxation
 - (iii) Index – cataract
3. Retinal astigmatism – due to oblique placement of macula.

Types of Regular Astigmatism

1. **With the rule astigmatism** : The two principal meridians are placed at right angles to one another but the **vertical meridian is more curved** than horizontal- more common.
2. **Against the rule astigmatism** : Horizontal meridian is more curved than the vertical meridian.
3. **Oblique astigmatism** : Is a type of regular astigmatism where the two principal meridians are not horizontal and vertical though they are at right angles to one another (45 and 135 deg)

Oblique astigmatism :

1. **Simple (or) meridional**


in both eyes.

2. **Compound (or) elementary**

one eye and at 150° in the other eye.


4. **Bi-oblique astigmatism :** In this type of regular astigmatism the two principal meridians are not at right angles to each other, one eye at 30° and other at 100° .

- **Optics of regular astigmatism :** In regular astigmatism the parallel rays of light are not focused on a point but form two focal lines – **Sturm's conoid**




Refractive types of Regular astigmatism

- Depending upon the position of two focal lines in relation to retina, regular astigmatism is further classified
- Simple :** Where one of the foci falls upon retina, the other focus may fall in front of or behind so that one meridian is emmetropic the other is either hypermetropic or myopic.



- Compound :** Where neither of two foci lie upon the retina but both are placed in front or behind it.


The state of the refraction is then entirely hypermetropic or entirely myopic. The former is known as compound hypermetropic, the latter as



compound myopic astigmatism.

- 3. Mixed :** Where one focus is in front of and other behind the retina so that the refraction is hypermetropic in one direction and myopic in the other.

AN UNCORRECTED ASTIGMATIC EYE WITH THE CIRCLE OF LEAST CONFUSION ON THE RETINA



- **Irregular Astigmatism :** Refraction in different meridians are irregular.

Etiological types:

1. Curvatural irregular astigmatism: irregular healing of cornea after trauma and inflammation (particularly ulceration & keratoconus)
2. Index irregular astigmatism : incipient cataract

Symptoms

1. Defective vision
2. Blurring of objects
3. Asthenopic symptoms - eyeyeache and headache
4. Running of lines

Treatment

- Optical – Spectacles with cylindrical lenses, Contact lens
(Toric contact lenses with prism ballast)
- Surgical
 1. Astigmatic keratotomy: Limbal Relaxing Incision, arcuate keratectomy, removal of sutures
 2. Photo-astigmatic refractive keratotomy (PARK)
 3. Laser: Excimer laser: LASIK or Femtosecond laser

Guidelines for Optical treatment

1. If the patient does not complain of asthenopic symptoms small astigmatic errors (0.5 D or less) generally do not require correction
2. If asthenopic symptoms are there, error should be corrected by cylindrical lenses.
3. Undercorrect the error initially
4. At a later date, full correction may be worn comfortably.