

ACCOMMODATION AND CONVERGENCE

1

Revision Question

- You have gone fishing and see a fish in the water. You do not have a fishing rod. The only equipment that you have is a spear to catch the fish.
- Where do you throw the spear?
 - a. in front of the fish.
 - b. behind the fish.
 - c. directly at the fish.
 - d. It is not possible to hit the fish as it is a virtual image.

Learning Objectives

At the end of the class, students shall be able to

- Understand the basic mechanism of accommodation and clinical importance of anomalies of accommodation
- Understand the pathway for near reflex and importance of convergence insufficiency.

3

Looking at the pupils




Looking at the pupils



5

ACCOMMODATION

- Definition: Accommodation is the **mechanism by which the eye changes its refractive power** by altering shape of lens in order to focus objects at variable distances.




6

Mechanism of accommodation

- Increase in the curvature of lens affects mainly anterior surface.
- Radius of curvature of anterior surface : **10 mm**

During accommodation




6 mm

This alteration in shape **increases** the converging power of lens.

7

RELAXATION THEORY OF HELMHOLTZ

- He considered that lens was elastic and in normal state is stretched and flattened by the tension of suspensory ligament.
- During accommodation: **Ciliary muscle contracts** causing lens zonules to slacken, lens assumes more spherical form increasing thickness and decreasing diameter, forward protrusion at centre and relative flattening at periphery.



9

NEAR REFLEX

- It has 2 components :
- **Convergence reflex** comprising convergence of visual axes of eyes and associated constriction of pupil.
- **Accommodation reflex** includes increased accommodation and associated constriction of pupil.
- The near reflex comprises :
Accommodation , convergence and miosis of pupils.

This diagram illustrates the visual pathway and the pathways for ocular motor control. At the top, the brain is shown with labels for the **Primary visual cortex** and **Visual association cortex**. The visual pathway is depicted with green lines, showing the flow from the eyes through the **LGN** (Lateral Geniculate Nucleus) to the visual cortex. The ocular motor pathways are shown with blue and orange lines, originating from the **Oc** (Oculomotor nucleus) and **CN III** (Cranial Nerve III). These pathways branch out to innervate the **Medial rectus** and **To ciliary muscle** of the eyes, as well as the **Pupillary sphincter**. The **CG** (Ciliary Ganglion) is also labeled. The diagram is numbered **11** in the bottom right corner.

12

❑ The **far point** of distinct vision is the position of an object such that its image falls on the retina in the relaxed eye, i.e. in the absence of accommodation. The **far point of emmetropic eye is at infinity.**

❑ The **near point** of distinct vision is the nearest point at which an object can be clearly seen when maximum accommodation is used.

13

❑ The **range of accommodation** is the distance between far point and near point.

❑ The **amplitude of accommodation** is the difference in dioptric power between the eye at rest and fully accommodated eye.

- The **amplitude of accommodation** is given by the formula
- $A = P - R$
- where A is amplitude of accommodation in dioptries
- P is dioptric value of near point distance
- R is dioptric value of far point distance.

15

- Applying this formula to the case of an emmetropic eye with a near point of 10 cm,
- $P = 10 \text{ D}$ (the reciprocal of 0.10 m)
- $R = 0$ (the reciprocal of infinity is zero)
- $A = 10 \text{ D}$

- **Far point and near point** of the eye **vary with the static refraction** of the eye
- In a **hypermetrope eye** far point is **virtual** and lies behind the eye
- In a **myopic eye** far point is **real** and lies in front of the eye.

17

In an emmetropic eye

- Far point is at **infinity**
- Near point varies with age
- About 7 cm at age of 10 years
- About 25 cm at age of 40 years
- 33 cm at age of 45 years

ANOMALIES OF ACCOMMODATION

DIMINISHED ACCOMMODATION

1. PHYSIOLOGICAL
(**PRESBYOPIA**)
2. PHARMACOLOGICAL
(Cycloplegia)
3. PATHOLOGICAL
 - Insufficiency of accommodation
 - Ill sustained accommodation
 - Paralysis of accommodation

INCREASED ACCOMMODATION

1. EXCESSIVE
ACCOMMODATION
2. SPASM OF
ACCOMMODATION

19

PRESBYOPIA

- Presbyopia is not an error of refraction but a condition of **physiological insufficiency** of accommodation due to reduced amplitude of accommodation, leading to a progressive fall in near vision.
- This usually begins between 40 years and 45 years of age.

- After the age of 40 years ,the NPA recedes beyond the normal reading distance.
- This condition of falling near vision due to age related decrease in the amplitude of accommodation or increase in punctum proximum is presbyopia.

21

Causes of presbyopia

- ☐ Age related changes in lens which include
 - Decrease in elasticity of lens capsule
 - Progressive increase in size and hardness (sclerosis) of lens substance.
- ☐ Age related decline in ciliary muscle power.

Causes of premature presbyopia include

- Uncorrected hypermetropia
- Premature sclerosis of crystalline lens
- General debility causing presenile weakness of ciliary muscle
- Chronic simple glaucoma

23

Symptoms

- **Difficulty in near vision** : patients start complaining of inadequacy of vision for small print and finer objects at usual reading distance. Such problems start occurring in evening, and in dim light.
- **Asthenopic** symptoms due to fatigue of ciliary muscle
- Intermittent diplopia at near may develop.

Treatment

Optical correction of presbyopia

Done by supplementing accommodation with **convex lens** of appropriate power.

The difference between distance correction and strength needed for near vision is called **the add**.

25

PRESBYOPIC ADD

- If patient is presbyopic, calculate the likely reading addition and add this to the distance lenses in the trial frame. In practice the reading addition is estimated from patient's age.

AGE RANGE	READING ADDITION
45-50 YEARS	+1.00 D
50-55 YEARS	+1.50 D
55-60 YEARS	+2.00D
OVER 60 YEARS	+2.50 D

MODES OF PRESCRIBING PRESBYOPIC ADD

- SPECTACLES
- CONTACT LENSES FOR PRESBYOPIA

2.SURGICAL TREATMENT OF PRESBYOPIA

- Refractive surgeries

27

Insufficiency of accommodation

Accommodative power is significantly and persistently below normal physiological limits for patient's age

- **Causes**

- Premature sclerosis of lens
- Weakness of ciliary muscle due to systemic causes such as diabetes mellitus.

Insufficiency of accommodation

Clinical features :

Headache

Fatigue

Blurring of vision for near work

Intermittent diplopia

Treatment :


1.Treatment of systemic cause

2.Near vision spectacles

3.Accommodation exercises

Paralysis of accommodation

- Paralysis of accommodation ,also known as **cycloplegia**, refers to complete absence of accommodation.



Causes

- Drug induced
- Internal ophthalmoplegia
- Paralysis of accommodation as a component of 3rd nerve palsy.

31

DRUG	MAXIMUM MYDRIASIS	MAXIMUM CYCLOPLEGIA	DURATION OF MYDRIASIS	DURATION OF CYCLOPLEGIA
ATROPINE 1% TID	30-40 MIN	1 DAY	7-10 DAYS	2 WEEKS
CYCLOPENTOLAT E 0.5%-1%	15 MIN	15-30 MIN	1 DAY	24 HRS
HOMATROPINE 2%	30-60 MIN	30-60 MIN	1-2 DAYS	1-2 DAYS
TROPICAMIDE 0.5%-1%	<u>15-30 MIN</u>	20-25 MIN	<u>4-6 HRS</u>	5-6 HRS
		www.FirstRanker.com		32

- **Clinical features**

- Blurring of near vision
- Photophobia (due to mydriasis)
- Micropsia
- Abnormal receding of near point
- Signs of 3rd nerve palsy

- **Treatment**

- Self recovery-drug induced paralysis and in cases when systemic cause is treated.
- Dark glasses – reduce glare
- Convex lenses –for near vision if paralysis is permanent.

Excessive accommodation

- A situation in which an individual exerts more than normal required accommodation for performing a certain near work.
- Excessive near work is an important precipitating factor especially when done in inadequate illumination.

Clinical features

- Varying degrees of blurred vision
- Symptoms of accommodative asthenopia
- Difficulty in performing near vision tasks

35

Treatment

1. Optical treatment : refractive error to be corrected
2. General treatment : Near work should be minimised and when done should be in proper illumination.
3. The general health of the patient should be improved.

CONVERGENCE


- Definition: Convergence is a **disconjugate movement in which both eyes rotate inward** so that lines of sight intersect in front of eyes.
- Allows bifoveal single vision to be maintained at any fixation distance.
- Convergence does not deteriorate with increasing age.
- The power of convergence can be increased by exercises.

Convergence reflex

- **Afferent pathway** –the afferents from medial recti travel centrally via the **3rd nerve** to the mesencephalic nucleus of the 5th nerve, to a **presumptive convergence centre in tectal or pretectal region**.
- **Internuncial fibres** : from the convergence centre go to the Edinger Westphal nucleus .


39

- **Efferent pathway**- along the 3rd nerve. From the **3rd nerve** efferent fibres of convergence reflex relay in the **accessory ganglion**, before reaching **sphincter pupillae**.



• Angle of convergence


- It refers to the angle that is formed between primary lines of sight during convergence
- Its size depends on
- the fixation distance
- and
- interpupillary distance (IPD)



41

Metre angle

- One metre angle convergence is exerted by each eye when the eyes are directed to object at a distance of 1 m of the meridian line between the two eyes.



- In an emmetropic eye, the number of dioptries of accommodation required to see an object clearly is equal to the number of metre angles through which each eye must converge to see the object singly.
- Thus 1D of accommodation is associated with 1 ma of convergence of each eye

43


- ❑ The **near point** of convergence is the closest point at which an object can be seen singly during bifoveal vision when maximum convergence is exerted.
- ❑ The **far point** of convergence refers to relative position of eyes when they are completely at rest, usually at infinity.

- ❑ The **range of convergence** is the distance between far point and near point of convergence
- ❑ The **amplitude of convergence** is the difference in convergence power exerted to maintain the eye in a position at rest and in a position of maximum convergence.

45

Measurement of amplitude of convergence

1. Prism bar
2. Synoptophore



Anomalies of convergence

1) Convergence insufficiency

Inability to maintain adequate binocular convergence for any length of time without undue effort.

47

■ ***Aetiology***

A. Primary or idiopathic –

Wide IPD, general debility, overwork.

B. Refractive errors- uncorrected high hypermetropia and myopia

C. Presbyopia

D. Muscular imbalances-

Exophoria, intermittent exotropia and vertical muscle imbalances.

Clinical features

1. Symptoms of muscular fatigue

- Eyestrain
- Headache and eye ache
- Difficulty in changing the focus from distant to near
- Itching, burning and soreness of eyes

2. Symptoms due to failure to maintain binocular vision

- Blurred near vision
- Intermittent crossed diplopia

Diagnosis

1. Remote NPC – if NPC > 10 cm, Convergence insufficiency is said to exist.
2. Decreased fusional convergence for near-when measured on synoptophore, if there is difficulty in attaining 30° of convergence.
3. Exophoria

Treatment

1. **Optical** treatment- Myopes are given full correction and hypermetropes undercorrection to stimulate their accommodation and simultaneously convergence.
2. **Orthoptic** treatment- exercises to increase the near point of convergence (NPC) and also to increase amplitude of fusional convergence.

51

Treatment

3. Prism Therapy

Base in (BI) prisms reading glasses or bifocals with prism in lower segment are useful.

4. Surgical treatment

- Last resort
- Medial rectus muscle resection may be performed.

Convergence paralysis

CAUSES

- Head injury
- Encephalitis
- Tabes dorsalis
- Narcolepsy
- Tumours

53

Convergence paralysis

- **Clinical features**
- Convergence is completely absent
- Exotropia and crossed diplopia occurs on attempted near vision
- Adduction is normal

Convergence paralysis

- TREATMENT

- ☐ Base In (BI) prisms

- ☐ Plus lenses with BI prisms

55

Summary

- **Accommodation** is the mechanism by which eye changes its refractive power by altering the shape of the lens in order to **focus objects at variable distances**.
- **Convergence** is a disconjugate movement in which both eyes rotate inward so that the lines of sight intersect in front of eyes.
- Near reflex comprises : Accommodation , convergence and miosis.

Question

- Which of the following is true regarding accommodation and contact lenses?
- **a.** In a myopic patient, contact lenses decrease the accommodative demand compared to spectacles.
- **b.** In a hyperopic patient, contact lenses increase the accommodative demand compared to spectacles.
- **c.** In a myopic patient, contact lenses increase accommodative demand compared to spectacles.
- **d.** There is no difference noted in accommodation when wearing contact lenses.