


Heart Sounds & Murmurs

Department of Physiology

Heart Sounds






Heart sounds

- \gt S₁ "lubb" caused by the closing of the AV valves
- > S₂ "dupp" caused by the closing of semilunar valves
- > S₃ –associated with blood flowing into the ventricles
- S₄ –associated with atrial contraction

Four heart sounds can be recorded by phonocardiography, but normally only the first and the second heart sounds, are audible through a stethoscope

First heart sound


- occurs when the atrioventricular (AV) valves close at the beginning of ventricular contraction
- generated by the vibration of the blood and the ventricular wall
- is louder, longer, more in duration than the second heart sound





Second heart sound

- occurs when aortic and pulmonary semilunar valves close at the beginning of ventricular dilation
- generated by the vibration of the blood and the aorta
- Aortic valve closes slightly before pulmonary valve
- Splitting of S₂



Heart Murmur

- Cardiac murmurs are abnormal heart sounds in the form of noises resulting from turbulent blood flow,
- Murmurs may be:
 - Systolic murmurs, if they occur during systole.
 - Diastolic murmurs, if they occur during diastole
- Causes:
 - Stenosis (narrowing) of heart valves
 - Incompetence of heart valves
 - Increase of blood flow or decrease of blood viscosity (as in anemia).



Two Basic Types of Valvular Diseases


valvular stenosis:

narrowing of the valve


valvular insufficiency (incompetence):

 valve is unable to close fully; so there is regurgitation





Phonocardiograms examples





Thank You

www.FireiRanker.com