

**B.Sc. Part-III (Semester-VI) Examination**
**MATHEMATICS**
**(Special Theory of Relativity)**
**Paper—XII**

Time : Three Hours]

[Maximum Marks : 60

**Note :—** (1) Question No. 1 is compulsory, attempt once.

 (2) Attempt **ONE** question from each unit.

1. Choose the correct alternative :

 (i) The interval  $ds^2 = -(dx^1)^2 - (dx^2)^2 - (dx^3)^2 + (dx^4)^2$  is said to be space like if : 1

- (a)  $ds^2 > 0$  (b)  $ds^2 < 0$   
 (c)  $ds^2 = 0$  (d) None of these

 (ii) The electric and magnetic field strengths  $E$  and  $H$  are invariant under : 1

- (a) Galilean Transformations (b) Laplace Transformations  
 (c) Fourier Transformations (d) Gauge Transformations

 (iii)  $A^i = (\bar{A}, \phi) = (Ax, Ay, Az, \phi)$  is a four potential then : 1

- (a)  $A_i = (\bar{A}, \phi)$  (b)  $A_i = (\bar{A}, -\phi)$   
 (c)  $A_i = (-\bar{A}, \phi)$  (d)  $A_i = (-\bar{A}, -\phi)$

 (iv)  $A^r = (A^1, A^2, A^3, A^4)$  is a four vector or four dimensional vector where  $A^2 < 0$  then  $A^r$  is : 1

- (a) Time like (b) Null or light like  
 (c) Space like (d) None of these

 (v) Covariant tensor of rank one  $T'_r$  is defined as : 1

- (a)  $T'_r = \frac{\partial x'^r}{\partial x^s} T_s$  (b)  $T'_r = \frac{\partial x'^r}{\partial x^s} T_r$   
 (c)  $T'_r = \frac{\partial x^s}{\partial x'^r} T_s$  (d)  $T'_r = \frac{\partial x^s}{\partial x'^r} T_r$

(vi) The special Lorentz transformations will reduce to simple Galilean transformations when : 1

- (a)  $V = C$  (b)  $C \ll V$   
 (c)  $V \ll C$  (d) None of these

(a)  $F_{ij} = \frac{\partial A_i}{\partial x^j} - \frac{\partial A_j}{\partial x^i}$

(b)  $F_{ij} = \frac{\partial A_j}{\partial x^i} - \frac{\partial A_i}{\partial x^j}$

(c)  $F_{ij} = \frac{\partial A_i}{\partial x^j} + \frac{\partial A_j}{\partial x^i}$

(d) None of these

 (viii) The transformations  $\bar{r}' = \bar{r} - \bar{v}t$  and  $t' = t$  are : 1

(a) Laplace transformations

(b) Lorentz transformations

(c) Galilean transformations

(d) None of these

 (ix) If  $\bar{A}$  is a vector potential then the magnetic field is : 1

(a)  $\bar{H} = \text{div} . \bar{A}$

(b)  $\bar{H} = \text{Curl} \bar{A}$

(c)  $\bar{H} = \text{div} . (\text{Curl} \bar{A})$

(d) None of these

 (x) Four velocity of a particle is : 1

(a) a unit space-like vector

(b) a unit time-like vector

(c) a unit light-like vector

(d) None of these

### UNIT—I

 2. (a) Obtain Galilean transformation equations for two inertial frames in relative motion. 3

 (b) Show that simultaneity is relative in special relativity. 3

(c) Show that the electromagnetic wave equation :

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2 f}{\partial t^2} = 0$$

 is not invariant under the Galilean transformations. 4

 3. (p) Discuss the geometrical interpretation of Lorentz transformations. 4

 (q) Prove that  $\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}$  is invariant under special Lorentz transformations. 4

 (r) Show that  $x^2 + y^2 + z^2 - c^2 t^2$  is Lorentz invariant. 2

### UNIT—II

 4. (a) Obtain the transformations for the velocity of a particle under special Lorentz transformations. 5

- (b) If  $\bar{u}$  and  $\bar{u}'$  be the velocities of a particle in two inertial systems  $s$  and  $s'$  respectively where  $s'$  is moving with velocity  $v$  relative to  $s$  along the  $XX'$  axis then show that :

$$\tan \theta' = \frac{\sin \theta \left(1 - \frac{v^2}{c^2}\right)^{1/2}}{\cos \theta - \frac{v}{u}}$$

and

$$u'^2 = \frac{u^2 \left[1 - 2 \frac{v}{u} \cos \theta + \frac{v^2}{u^2} - \frac{v^2}{c^2} \sin^2 \theta\right]}{\left(1 - \frac{uv}{c^2} \cos \theta\right)^2}$$

where  $\theta$  and  $\theta'$  are the angles made by  $u$  and  $u'$  with the  $X$ -axis respectively. 5

5. (p) If  $\bar{u}$  and  $\bar{u}'$  be the velocities of a particle in two inertial systems  $s$  and  $s'$  respectively then prove that :

$$\sqrt{1 - \frac{u^2}{c^2}} = \frac{\sqrt{1 - \frac{u'^2}{c^2}} \sqrt{1 - \frac{v^2}{c^2}}}{\left(\frac{1 + \frac{u'_x v}{c}}{c}\right)},$$

where  $s'$  is moving with velocity  $v$  relative to  $s$  along  $XX'$  axis. 5

- (q) Show that in nature no signal can move with a velocity greater than the velocity of light relative to any inertial system. 5

### UNIT—III

6. (a) Define time-like, space-like and light-like intervals for the space time geometry of special relativity. 3
- (b) Define a four tensor of the second order. Prove that :

$$(i) \quad T'^{11} = \alpha^2 \left\{ T^{11} - \frac{v}{c} T^{14} - \frac{v}{c} T^{41} + \frac{v^2}{c^2} T^{44} \right\} \text{ and}$$

$$(ii) \quad T'^{14} = \alpha^2 \left\{ -\frac{v}{c} T^{11} + T^{14} + \frac{v^2}{c^2} T^{41} - \frac{v}{c} T^{44} \right\} \quad 1+3+3$$

7. (p) Define a four vector  $A^i$ . Show that :

$$A^1 = -A_1, A^2 = -A_2, A^3 = -A_3, A^4 = A_4. \quad 1+3$$

- (q) Prove that there exists an inertial system  $s'$  in which the two events occur at one and the same time if the interval between two events is space-like. 4

- (r) Write the Lorentz transformations in index form. 2

#### UNIT—IV

8. (a) Deduce Einstein's mass energy equivalence relation. 5  
 (b) Define : Four velocity. Prove that the four velocity in component form can be expressed as :

$$u^i = \left( \frac{\bar{u}}{c\sqrt{1-u^2/c^2}}, \frac{1}{\sqrt{1-u^2/c^2}} \right)$$

where  $\bar{u} = (u_x, u_y, u_z) =$  velocity of the particle. 1+4

9. (p) Define : Four momentum vector  $p^i$ . Prove that the square of the magnitude of the four momentum vector  $p^i$  is  $m^2 \circ c^2$ . 1+4

- (q) A particle is given a kinetic energy equal to  $n$  times its rest energy  $m \circ c^2$ . Find speed and momentum of the particle.  $\left( \text{Kinetic energy} = T = m \circ c^2 \left\{ \frac{1}{\sqrt{1-v^2/c^2}} - 1 \right\} \right)$  5

#### UNIT—V

10. (a) Show that the Hamiltonian for a charged particle moving in an electromagnetic field is :

$$H = \left\{ m^2 c^4 + c^2 \left( p - \frac{e}{c} A \right)^2 \right\}^{1/2} + e\phi. \quad 5$$

- (b) Define : Current four vector. Show that  $c^2 p^2 - J^2$  is invariant and its value is  $\rho^2 \circ c^2$ . 1+4

11. (p) Prove that the set of Maxwell's equations  $\text{div. } \vec{H} = 0$  and  $\vec{E} = -\frac{1}{c} \frac{\partial \vec{H}}{\partial t}$  can be written

$$\text{as } \frac{\partial F_{ij}}{\partial x^k} + \frac{\partial F_{jk}}{\partial x^i} + \frac{\partial F_{ki}}{\partial x^j} = 0, \text{ where } F_{ij} \text{ is the electro-magnetic field tensor.} \quad 5$$

- (q) Define electromagnetic field tensor  $F_{ij}$ . Express the components of  $F_{ij}$  in terms of the electric and magnetic field strengths. 1+4