FirstRanker.com
Firstranker's choice
www.FirstRanker.com www.FirstRanker.com

B.Sc. Part-III (Semester-VI) Examination

MATHEMATICS

(Special Theory of Relativity)

Paper—XII

Time : Three Hours] [Maximum Marks : 60

Note :— (1) Question No. **1** is compulsory, attempt once.

(2) Attempt **ONE** question from each unit.

1. Choose the correct alternative :

(i) The interval $ds^2 = -(dx^1)^2 - (dx^2)^2 - (dx^3)^2 + (dx^4)^2$ is said to be space like if : 1

(a) $ds^2 > 0$ (b) $ds^2 < 0$

(c) $ds^2 = 0$ (d) None of these

(ii) The electric and magnetic field strengths E and H are invariant under : 1

(a) Galilean Transformations (b) Laplace Transformations

(c) Fourier Transformations (d) Guage Transformations

(iii) $A^i = (\overline{A}, \phi) = (Ax, Ay, Az, \phi)$ is a four potential then : 1

(a) $A_i = (\overline{A}, \phi)$ (b) $A_i = (\overline{A}, -\phi)$

(c) $A_i = (-\overline{A}, \phi)$ (d) $A_i = (-\overline{A}, -\phi)$

(iv) $A^r = (A^1, A^2, A^3, A^4)$ is a four vector or four dimensional vector where $A^2 < 0$ then A^r is : 1

(a) Time like (b) Null or light like

(c) Space like (d) None of these

(v) Covariant tensor of rank one T'_r is defined as : 1

(a) $T'_r = \frac{\partial x'^r}{\partial x^5} T_s$ (b) $T'_r = \frac{\partial x'^r}{\partial x^5} T_r$

(c) $T'_r = \frac{\partial x^5}{\partial x'^r} T_s$ (d) $T'_r = \frac{\partial x^5}{\partial x'^r} T_r$

(vi) The special Lorentz transformations will reduce to simple Galilean transformations when : 1

(a) $V = C$ (b) $C << V$

(c) $V << C$ (d) None of these

www.FirstRanker.com

FirstRanker.com
Firstranker's choice
www.FirstRanker.com www.FirstRanker.com

(vii) The electromagnetic field tensor (or Maxwell tensor) F_{ij} is defined as : 1

(a) $F_{ij} = \frac{\partial A_i}{\partial x^j} - \frac{\partial A_j}{\partial x^i}$ (b) $F_{ij} = \frac{\partial A_j}{\partial x^i} - \frac{\partial A_i}{\partial x^j}$

(c) $F_{ij} = \frac{\partial A_i}{\partial x^j} + \frac{\partial A_j}{\partial x^i}$ (d) None of these

(viii) The transformations $\bar{r}' = \bar{r} - \bar{v}t$ and $t' = t$ are : 1

(a) Laplace transformations (b) Lorentz transformations

(c) Galilean transformations (d) None of these

(ix) If $\bar{A}$ is a vector potential then the magnetic field is : 1

(a) $\bar{H} = \text{div}.\bar{A}$ (b) $\bar{H} = \text{Curl}\,\bar{A}$

(c) $\bar{H} = \text{div}.(\text{Curl}\,\bar{A})$ (d) None of these

(x) Four velocity of a particle is : 1

(a) a unit space-like vector (b) a unit time-like vector

(c) a unit light-like vector (d) None of these

UNIT—I

2. (a) Obtain Galilean transformation equations for two inertial frames in relative motion. 3

(b) Show that simultaneity is relative in special relativity. 3

(c) Show that the electromagnetic wave equation :

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} - \frac{1}{c^2}\frac{\partial^2 f}{\partial t^2} = 0$$

is not invariant under the Galilean transformations. 4

3. (p) Discuss the geometrical interpretation of Lorentz transformations. 4

(q) Prove that $\nabla^2 - \frac{1}{c^2}\frac{\partial^2}{\partial t^2}$ is invariant under special Lorentz transformations. 4

(r) Show that $x^2 + y^2 + z^2 - c^2t^2$ is Lorentz invariant. 2

UNIT—II

4. (a) Obtain the transformations for the velocity of a particle under special Lorentz transformations. 5

www.FirstRanker.com

FirstRanker.com
Firstranker's choice

(b) If $\bar{u}$ and $\bar{u}'$ be the velocities of a particle in two inertial systems s and s' respectively www.FirstRanker.com www.FirstRanker.com where s' is moving with velocity v relative to s along the XX' axis then show that :

$$\tan\theta' = \frac{\sin\theta\left(1-\frac{v^2}{c^2}\right)^{1/2}}{\cos\theta - \frac{v}{u}}$$

and

$$u'^2 = \frac{u^2\left[1-2\frac{v}{u}\cos\theta+\frac{v^2}{u^2}-\frac{v^2}{c^2}\sin^2\theta\right]}{\left(1-\frac{uv}{c^2}\cos\theta\right)^2}$$

where θ and θ' are the angles made by u and u' with the X-axis respectively. 5

5. (p) If $\bar{u}$ and $\bar{u}'$ be the velocities of a particle in two inertial systems s and s' respectively then prove that :

$$\sqrt{1-\frac{u^2}{c^2}} = \frac{\sqrt{1-\frac{u'^2}{c^2}}\sqrt{1-\frac{v^2}{c^2}}}{\left(\frac{1+u'_x v}{c}\right)},$$

where s' is moving with velocity v relative to s along XX' axis. 5

(q) Show that in nature no signal can move with a velocity greater than the velocity of light relative to any inertial system. 5

UNIT—III

6. (a) Define time-like, space-like and light-like intervals for the space time geometry of special relativity. 3

(b) Define a four tensor of the second order. Prove that :

(i) $T'^{11} = \alpha^2\left\{T^{11} - \frac{v}{c}T^{14} - \frac{v}{c}T^{41} + \frac{v^2}{c^2}T^{44}\right\}$ and

(ii) $T'^{14} = \alpha^2\left\{-\frac{v}{c}T^{11} + T^{14} + \frac{v^2}{c^2}T^{41} - \frac{v}{c}T^{44}\right\}$ 1+3+3

7. (p) Define a four vector A^r. Show that :

$A^1 = -A_1$, $A^2 = -A_2$, $A^3 = -A_3$, $A^4 = A_4$. 1+3

www.FirstRanker.com

FirstRanker.com
Firstranker's choice

(q) Prove that there exists an inertial system s' in which the two events occur at one and the same time if the interval between two events is space-like. 4

www.FirstRanker.com www.FirstRanker.com

(r) Write the Lorentz transformations in index form. 2

UNIT—IV

8. (a) Deduce Einstein's mass energy equivalence relation. 5

(b) Define : Four velocity. Prove that the four velocity in component form can be expressed as :

$$u^i = \left(\frac{\bar{u}}{c\sqrt{1-u^2/c^2}}, \frac{1}{\sqrt{1-u^2/c^2}} \right)$$

where $\bar{u} = (u_x, u_y, u_z)$ = velocity of the particle. 1+4

9. (p) Define : Four momentum vector p^i. Prove that the square of the magnitude of the four momentum vector p^i is $m^2 \circ c^2$. 1+4

(q) A particle is given a kinetic energy equal to n times its rest energy $m \circ c^2$. Find speed and momentum of the particle. $\left(\text{Kinetic energy} = T = m \circ c^2 \left\{ \frac{1}{\sqrt{1-v^2/c^2}} - 1 \right\} \right)$ 5

UNIT—V

10. (a) Show that the Hamiltonian for a charged particle moving in an electromagnetic field is :

$$H = \left\{ m^2c^4 + c^2 \left(p - \frac{e}{c} A \right)^2 \right\}^{1/2} + e\phi .$$ 5

(b) Define : Current four vector. Show that $c^2\rho^2 - J^2$ is invariant and its value is $\rho^2 \circ c^2$. 1+4

11. (p) Prove that the set of Maxwell's equations $\text{div}.\bar{H} = 0$ and $\bar{E} = -\frac{1}{c}\frac{\partial \bar{H}}{\partial t}$ can be written as $\frac{\partial F_{ij}}{\partial x^k} + \frac{\partial F_{jk}}{\partial x^i} + \frac{\partial F_{ki}}{\partial x^j} = 0$, where F_{ij} is the electro-magnetic field tensor. 5

(q) Define electromagnetic field tensor F_{ij}. Express the components of F_{ij} in terms of the electric and magnetic field strengths. 1+4

www.FirstRanker.com