



B.Sc. (Part—I) (Semester—I) Examination

MATHEMATICS

(Algebra & Trigonometry)

Paper—I

Time : Three Hours]

[Maximum Marks : 60

Note :— (1) Question **ONE** is compulsory. Attempt once.(2) Attempt **ONE** question from each Unit.

1. Choose the correct alternative :—

(1) Which **one** of the following statements is true :—

10

(a) $\cosh(x + iy) = \cosh x \cdot \cos y + i \sinh x \cdot \sin y$

(b) $\cosh(x + iy) = \cos x \cos y + i \sin x \sin y$

(c) $\cosh(x + iy) = \cosh x + \cos y - i \sinh x \cdot \sin y$

(d) $\cosh(x + iy) = \cosh x \sin y + i \sinh x \cos y$

(2) What is the value of $\sinh^{-1}x$:

(a) $\log[x + \sqrt{x^2 + 1}]$

(b) $\log[x + \sqrt{x^2 - 1}]$

(c) $\log[x + \sqrt{1 - x^2}]$

(d) None of these

(3) The value of $4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99}$ is _____.

(a) $\pi/2$

(b) $\pi/4$

(c) $\pi/3$

(d) π

(4) Sum of the series $x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^{n+1} \frac{x^n}{n} + \dots$; $-1 < x < 1$ is denoted by ____.

(a) $\log(1 + x)$

(b) $\sinh x$

(c) $\cosh x$

(d) e^x

(5) If $q = 2 + 2i - j + 4k$ then the norm of q is _____.

(a) -5

(b) 5

(c) $1/5$

(d) None of these

(6) The inverse of unit quaternion is its ____.

(a) Purely imaginary

(b) Purely real

(c) Complex conjugate

(d) None of these

(7) If $\alpha + i\beta$ be the root of quadratic polynomial $f(x) = 0$ then its another root is ____.

(a) $\alpha - i\beta$

(b) α

(c) β

(d) None of these

(8) If α, β, γ are the roots of the equation $px^3 + qx^2 + rx + s = 0$ then $\Sigma \alpha$ is _____.

(a) $\frac{q}{p}$

(b) $-\frac{q}{p}$

(c) $\frac{r}{p}$

(d) $\frac{s}{p}$



- (9) If A and B are the non-singular matrices of order n then _____.
 (a) $(AB)^{-1} = AB$ (b) $(AB)^{-1} = A^{-1} \cdot B^{-1}$
 (c) $(AB)^{-1} = B^{-1} \cdot A^{-1}$ (b) None of these
- (10) 'Every square matrix satisfies its own characteristics equation' is the statement of _____.
 (a) Lagrange's MVT (b) De-Moivre's theorem
 (c) Cayley-Hamilton theorem (d) Cauchy's MVT

UNIT—I

2. (a) Prove that $\frac{1 + \sin \theta + i \cos \theta}{1 + \sin \theta - i \cos \theta} = \sin \theta + i \cos \theta$.
 Hence prove that $\left(1 + \sin \frac{\pi}{5} + i \cos \frac{\pi}{5}\right) + i \left(1 + \sin \frac{\pi}{5} - i \cos \frac{\pi}{5}\right) = 0$. 5
- (b) If $\sin(\alpha + i\beta) = x + iy$ then prove that $\frac{x^2}{\cosh^2 \beta} + \frac{y^2}{\sinh^2 \beta} = 1$ and $\frac{x^2}{\sin^2 \alpha} - \frac{y^2}{\cos^2 \alpha} = 1$ 5
3. (p) Prove that one of the value of :
 $(\cos \theta + i \sin \theta)^n$ is $(\cos n\theta + i \sin n\theta)$; when n is negative integer. 5
 (q) Separate real and imaginary parts of $\tan(x + iy)$. 5

UNIT—II

4. (a) Find the Sum of the series :
 $C = 1 + e^{\sin x} \cdot \cos(\cos x) + \frac{1}{2!} e^{2 \sin x} \cdot \cos(2 \cos x) + e^{3 \sin x} \cdot \frac{1}{3!} \cos(3 \cos x) + \dots$ 5
- (b) Prove that $4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99} = \frac{\pi}{4}$. 5
5. (p) Find the sum of the series $\sinh x + \frac{1}{2!} \sinh 2x + \frac{1}{3!} \sinh 3x + \dots$ 5
- (q) If $-\frac{\pi}{4} \leq x \leq \frac{\pi}{4}$ then prove that
 $x = \tan x - \frac{1}{3} \tan^3 x + \frac{1}{5} \tan^5 x + \dots + (-1)^{n-1} \frac{1}{2n-1} \tan^{2n-1} x + \dots$ 5

UNIT—III

6. (a) Prove that for $p, q \in H$, $N(pq) = N(p) N(q)$ and $N(q^*) = N(q)$. 5
- (b) For the quaternion $q = \cos \frac{\pi}{3} + i \sin \frac{\pi}{3}$ and the input vector $v = i$, compute the output vector w under the action of the operators L_q and L_{q^*} . 5
7. (p) Show that the quaternion product need not be commutative. 5
- (q) For any $p, q \in H$, show that $pq = qp$ if and only if p and q are parallel. 5

**UNIT—IV**

8. (a) Find the roots of the equation, $8x^3 + 18x^2 - 27x - 27 = 0$, if these roots are in geometric progression. 5
- (b) State Descartes rule of sign. Find the nature of the roots of the equation $2x^7 - x^4 + 4x^3 - 5 = 0$. 5
9. (p) Prove that in an equation with real coefficients complex roots occur in pairs. 5
- (q) Solve the equation $x^4 - 2x^3 - 22x^2 + 62x - 15 = 0$; given that $2\sqrt{3}$ is one of the root. 5

UNIT—V

10. (a) Show that if λ is the eigen value of a nonsingular matrix A then λ^{-1} is the eigen value of A^{-1} . 5
- (b) Find the eigen values and the corresponding eigen vector for smallest eigen value of the matrix $A = \begin{bmatrix} 8 & -8 & -2 \\ 4 & -3 & -2 \\ 3 & -4 & 1 \end{bmatrix}$. 5
11. (p) Show that the eigen values of any square matrix A and A' are same. 5
- (q) Reduce to canonical form and find the rank of the matrix $A = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix}$. 5



FirstRanker.com

FirstRanker's choice

www.FirstRanker.com

www.FirstRanker.com

firstRanker.com