

**B.Sc. (Part—I) (Semester—I) Examination
MATHEMATICS
(Algebra & Trigonometry)
Paper—I**

Time : Three Hours]

[Maximum Marks : 60]

Note :—(1) Question **ONE** is compulsory. Attempt once.
(2) Attempt **ONE** question from each Unit.

1. Choose the correct alternative :—

- (1) Which one of the following statements is true :— 10

- (a) $\cosh(x + iy) = \cosh x \cdot \cos y + i \sinh x \cdot \sin y$
 - (b) $\cosh(x + iy) = \cos x \cos y + i \sin x \sin y$
 - (c) $\cosh(x + iy) = \cosh x + \cos y - i \sinh x \cdot \sin y$
 - (d) $\cosh(x + iy) = \cosh x \sin y + i \sinh x \cos y$

- (2) What is the value of $\sinh^{-1}x$:

- (a) $\log[x + \sqrt{x^2 + 1}]$ (b) $\log[x + \sqrt{x^2 - 1}]$
 (c) $\log[x + \sqrt{1 - x^2}]$ (d) None of these

- (3) The value of $4\tan^{-1}\frac{1}{5} - \tan^{-1}\frac{1}{70} + \tan^{-1}\frac{1}{99}$ is _____.

- (a) $\pi/2$ (b) $\pi/4$
 (c) $\pi/3$ (d) π

- (4) Sum of the series $x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^{n+1} \frac{x^n}{n} + \dots$; $-1 < x < 1$ is denoted by ____.

- (a) $\log(1 + x)$ (b) $\sinh x$
 (c) $\cosh x$ (d) e^x

- (5) If $q = 2 + 2i - j + 4k$ then the norm of q is _____.

- (6) The inverse of unit quaternion is its ____.

- (a) Purely imaginary (b) Purely real
 (c) Complex conjugate (d) None of these

- (7) If $\alpha + i\beta$ be the root of quadratic polynomial $f(x) = 0$ then its another root is ____.

- (8) If α, β, γ are the roots of the equation $px^3 + qx^2 + rx + s = 0$ then $\Sigma\alpha$ is ____.

- $$(a) \frac{q}{p} \quad (b) -\frac{q}{p}$$

- (c) $\frac{r}{p}$ (d) $\frac{s}{p}$

(9) If A and B are the non-singular matrices of order n then ____.

- (a) $(AB)^{-1} = AB$ (b) $(AB)^{-1} = \bar{A}^T \cdot \bar{B}^T$
 (c) $(AB)^{-1} = B^T \cdot A^T$ (d) None of these

(10) 'Every square matrix satisfies its own characteristics equation' is the statement of ____.

- (a) Lagrange's MVT (b) De-Moivre's theorem
 (c) Cayley-Hamilton theorem (d) Cauchy's MVT

UNIT—I

2. (a) Prove that $\frac{1+\sin\theta+i\cos\theta}{1+\sin\theta-i\cos\theta} = \sin\theta+i\cos\theta$.

$$\text{Hence prove that } \left(1 + \sin \frac{\pi}{5} + i \cos \frac{\pi}{5}\right) + i \left(\left(1 + \sin \frac{\pi}{5} - i \cos \frac{\pi}{5}\right)\right) = 0. \quad 5$$

(b) If $\sin(\alpha + i\beta) = x + iy$ then prove that $\frac{x^2}{\cosh^2 \beta} + \frac{y^2}{\sinh^2 \beta} = 1$ and $\frac{x^2}{\sin^2 \alpha} - \frac{y^2}{\cos^2 \alpha} = 1 \quad 5$

3. (p) Prove that one of the value of :

$(\cos\theta + i\sin\theta)^n$ is $(\cos n\theta + i\sin n\theta)$; when n is negative integer. 5

(q) Separate real and imaginary parts of $\tan(x+iy)$. 5

UNIT—II

4. (a) Find the Sum of the series :

$$C = 1 + e^{\sin x} \cdot \cos(\cos x) + \frac{1}{2!} e^{2\sin x} \cdot \cos(2\cos x) + e^{3\sin x} \cdot \frac{1}{3!} \cos(3\cos x) + \dots \quad 5$$

(b) Prove that $4\tan^{-1}\frac{1}{5} - \tan^{-1}\frac{1}{70} + \tan^{-1}\frac{1}{99} = \frac{\pi}{4}$. 5

5. (p) Find the sum of the series $\sinh x + \frac{1}{2!} \sinh 2x + \frac{1}{3!} \sinh 3x + \dots \quad 5$

(q) If $-\frac{\pi}{4} \leq x \leq \frac{\pi}{4}$ then prove that

$$x = \tan x - \frac{1}{3} \tan^3 x + \frac{1}{5} \tan^5 x + \dots + (-1)^{n-1} \frac{1}{2n-1} \tan^{2n-1} x + \dots \quad 5$$

UNIT—III

6. (a) Prove that for $p, q \in H$, $N(pq) = N(p) N(q)$ and $N(q^*) = N(q)$. 5

(b) For the quaternion $q = \cos \frac{\pi}{3} + i \sin \frac{\pi}{3}$ and the input vector $v = i$, compute the output vector w under the action of the operators L_q and L_{q^*} . 5

7. (p) Show that the quaternion product need not be commutative. 5

(q) For any $p, q \in H$, show that $pq = qp$ if and only if p and q are parallel. 5

8. (a) Find the roots of the equation, $8x^3 + 18x^2 - 27x - 27 = 0$, if these roots are in geometric progression. 5
 (b) State Descartes rule of sign. Find the nature of the roots of the equation $2x^7 - x^4 + 4x^3 - 5 = 0$. 5
9. (p) Prove that in an equation with real coefficients complex roots occur in pairs. 5
 (q) Solve the equation $x^4 - 2x^3 - 22x^2 + 62x - 15 = 0$; given that $2\sqrt{3}$ is one of the root. 5

UNIT—V

10. (a) Show that if λ is the eigen value of a nonsingular matrix A then λ^{-1} is the eigen value of A^{-1} . 5
 (b) Find the eigen values and the corresponding eigen vector for smallest eigen value of the matrix $A = \begin{bmatrix} 8 & -8 & -2 \\ 4 & -3 & -2 \\ 3 & -4 & 1 \end{bmatrix}$. 5
11. (p) Show that the eigen values of any square matrix A and A' are same. 5
 (q) Reduce to canonical form and find the rank of the matrix $A = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix}$. 5



www.FirstRanker.com

www.FirstRanker.com