

ATAL BIHARI VAJPAYEE MEDICAL UNIVERSITY, LUCKNOW**MBBS DEGREE - IInd PROFESSIONAL - REGULAR EXAMINATION - MARCH-2024****PATHOLOGY - PAPER - II****SECTION A - Multiple Choice Questions (MCQs)**

PAPER CODE : 2412130004

SET : C

Note: Choose one correct answer in the OMR sheet provided. No overwriting should be done.

Time: 20 Minutes

(20 x 1 = 20 MARKS)

1. Neurofibrillary tangles are characteristically seen in:
 - a) Parkinson's disease
 - b) Multiple sclerosis
 - c) Alzheimer's disease
 - d) Prion disease
2. MEN 1 syndrome includes all except:
 - a) Parathyroid hyperplasia
 - b) Pituitary adenoma
 - c) Pancreatic tumor
 - d) Pheochromocytoma
3. Hirschsprung disease requires histological confirmation of loss of:
 - a) Villi
 - b) Muscularis
 - c) Stromal layer
 - d) Ganglion cells
4. Steroid resistant nephritic syndrome caused by mutation in gene encoding:
 - a) Podocin
 - b) Nephrin 1
 - c) Alpha actinin 4
 - d) Nephrin 2
5. Gluten intolerance is seen in:
 - a) Celiac disease
 - b) Whipple's disease
 - c) Short bowel syndrome
 - d) Tropical sprue

6. Which is the commonest tumor of uterus?
 - a) Lipoma
 - b) Leiomyoma
 - c) Adenocarcinoma
 - d) Fibroadenoma
7. Gleason's score is related to carcinoma of:
 - a) Pancreas
 - b) Pituitary gland
 - c) Prostate
 - d) Parathyroid gland
8. The commonest morphological variant of RCC is:
 - a) Granular cell type
 - b) Papillary type
 - c) Sarcomatoid type
 - d) Clear cell type
9. Lesion of thyroid associated with viral etiology is:
 - a) Grave's thyroiditis
 - b) Hashimoto thyroiditis
 - c) Riedel's thyroiditis
 - d) DeQuervain's thyroiditis
10. Reactivated TB is most commonly located near:
 - a) Bronchus
 - b) Lung apex
 - c) Base
 - d) Subpleural
11. Flexner Wintersteiner rosettes are seen in:
 - a) Neuroblastoma
 - b) Retinoblastoma
 - c) Medulloblastoma
 - d) Ependymoma
12. In obstructive jaundice, LFTs usually show:
 - a) Elevated direct bilirubin and alkaline phosphatase
 - b) Elevated indirect bilirubin and ALT
 - c) Elevated direct bilirubin and ALT
 - d) Elevated indirect bilirubin and GGT

13. Toxic megacolon is seen in:
- a) Celiac disease
 - b) Ulcerative colitis
 - c) Appendicitis
 - d) Crohn's disease
14. Elevated cANCA is seen in:
- a) Polyarteritis nodosa
 - b) Churg Strauss vasculitis
 - c) Wegener's granulomatosis
 - d) Microscopic polyangiitis
15. Increased serum calcitonin is observed in:
- a) Medullary carcinoma (Thyroid)
 - b) Follicular carcinoma (Thyroid)
 - c) Anaplastic carcinoma (Thyroid)
 - d) Papillary carcinoma (Thyroid)
16. Peutz-Jeghers polyp commonly located at:
- a) Appendix
 - b) Colon
 - c) Jejunum
 - d) Stomach
17. Ferruginous bodies are seen in:
- a) Asbestosis
 - b) Silicosis
 - c) Bagassosis
 - d) Byssinosis
18. Tumor of bone that usually affects children and adolescents is:
- a) Ewing's sarcoma
 - b) Osteoclastoma
 - c) Chondrosarcoma
 - d) Osteosarcoma
19. Sclerosing retroperitoneal fibrosis is related to:
- a) IgA
 - b) IgM
 - c) IgG2
 - d) IgG4

20. Libman-Sacks endocarditis is found in:

- a) Syphilis
- b) Lymphoma
- c) Rheumatoid arthritis
- d) Typhoid

www.FirstRanker.com