

Madhya Pradesh Medical Science University, Jabalpur
MBBS Final Part-2 Examination 2019
Subject – General Surgery II

Total MCQs : 20 $20 \times \frac{1}{2} = 10$

1. True about breast cancer in pregnancy:
 - a. Occurs in 1 every 3000 pregnant women
 - b. MC non-gynaecological cancer associated with pregnancy
 - c. Ductal carcinoma is MC type, accounting for 75–90% of breast cancer in pregnancy
 - d. All of the above
2. Mondor's disease is:
 - a. Thrombophlebitis of the superficial veins of breast
 - b. Carcinoma of the breast
 - c. Premalignant condition of the breast
 - d. Filariasis of the breast
3. Acute mastitis commonly occurs during
 - a. Pregnancy
 - b. Puberty
 - c. Lactation
 - d. Infancy
4. Best investigation for acute intestinal obstruction is:
 - a. Barium studies
 - b. X-ray
 - c. USG
 - d. ERCP
5. In case of newborn, the commonest cause of intestinal obstruction is:
 - a. Annular pancreas
 - b. Duodenal atresia
 - c. Jejunal atresia
 - d. Esophageal atresia
6. "Snow-storm" ascites is seen in:
 - a. Meconium ileus
 - b. Hirschsprung's disease
 - c. Ileocecal tuberculosis
 - d. Pseudomyxoma peritonei
7. Fluid levels are not visible in:
 - a. Meconium ileus
 - b. Intussusception
 - c. Colon pouch
 - d. Duodenal obstruction
8. Best treatment of acute suppurative cholangitis is:
 - a. Laparoscopic cholecystectomy
 - b. Open cholecystectomy
 - c. Endoscopic papillotomy
 - d. Choledochotomy
9. Most common cause of cholangitis:

- a. Viral infection
 - b. CBD stone
 - c. Surgery
 - d. Amebic infection
10. Most common surgical cause of obstructive jaundice:
- a. Periapillary carcinoma
 - b. Carcinoma gallbladder
 - c. Carcinoma head of pancreas
 - d. CBD stones
11. Common bile duct stones will manifest all except:
- a. Distended gallbladder
 - b. Jaundice
 - c. Itching
 - d. Clay-colored stools
12. Leucine aminopeptidase is elevated in obstruction of:
- a. Ureter
 - b. Urethra
 - c. Common bile duct
 - d. Spermatic cord
13. An absolute contraindication for IVP is:
- a. Allergy to the drug
 - b. Multiple myeloma
 - c. Blood urea > 200 mg
 - d. Renal tumor
14. Radiation exposure is the least in the following procedure:
- a. Micturating cystourethrogram
 - b. IVP
 - c. Bilateral nephrostogram
 - d. Spiral CT for stones
15. Central stellate scar on CT scans are seen in:
- a. Renal hemangioma
 - b. Renal oncocytoma
 - c. Wilm's tumor
 - d. Papillomas
16. All are features of hypernephroma except:
- a. Persistent pyrexia
 - b. Hematuria
 - c. Polycythemia
 - d. Lower pole involvement
17. Most important indicator for renal cell carcinoma:
- a. Nuclear grade
 - b. Histological type
 - c. Size
 - d. Pathological staging
18. All of the following tumors may be malignant except:
- a. Glioma

- b. Astrocytoma
- c. Hemangioblastoma
- d. Ependymoma

19. In children, most common posterior fossa tumour is:

- a. Meningiomas
- b. Astrocytoma
- c. Medulloblastoma
- d. Glioblastoma multiforme

20. Chang staging is used for:

- a. Retinoblastoma
- b. Medulloblastoma
- c. Ewing's sarcoma
- d. Rhabdomyosarcoma

firstranker.com
www.FirstRanker.com