

Dr. NTR UNIVERSITY OF HEALTH SCIENCES :: AP :: VIJAYAWADA – 520 008

FINAL M.B.B.S. PART – II – DEGREE EXAMINATION – DECEMBER, 2024

OBSTETRICS & GYNAECOLOGY – PAPER I

SET-B (Multiple Choice Questions)

Time: 20 Minutes

Max. Marks: 20

Note: Answer all questions

SECTION – I (MCQs – 20 MARKS)**1x20=20**

- 1) Bandi's ring is also called
 - a. Constriction ring
 - b. Schroeder's ring
 - c. Retraction ring
 - d. Cervical dystonia
- 2) What is the most favorable clinical scenario for a successful Trial of Labor (TOL) after cesarean birth
 - a. Only on previous cesarean delivery
 - b. Anesthesia immediately available
 - c. A previous vaginal delivery
 - d. Clear amniotic fluid
- 3) In which of the following cases might internal podalic version be indicated
 - a. Double foetaling breech
 - b. Impacted shoulder presentation
 - c. Vertex delivery of the first twin and transverse presentation of second twin
 - d. Term transverse lie with cervix completely dilated and membranes
- 4) A patient becomes unresponsive during active labor. The absolute diagnosis of amniotic fluid embolism is made by which of the following
 - a. Chest pain
 - b. Chest X-ray
 - c. Amniotic debris in pulmonary circulation
 - d. ECG changes
- 5) Most common cause of first trimester abortion
 - a. Intrauterine infection
 - b. Hemorrhage
 - c. Chromosomal anomalies
 - d. APLA
- 6) A patient with a history of recurrent abortions and hypercoagulable state is likely to be suffering from
 - a. APLA syndrome
 - b. Heparin induced thrombocytopenia
 - c. Leiden mucosa
 - d. Anti thrombin 3 deficiency

- 7) Which of the following factors tends to increase the average duration of labor
- Increased parity
 - Increasing age of mother
 - Decreasing size of the baby
 - Occipito posterior position of the baby
- 8) A patient has entered spontaneous premature labor at 28 weeks gestation. During the vertex delivery one should do which of the following
- Perform a episiotomy
 - Use prophylactic forceps
 - Use vacuum extraction
 - Allow spontaneous vaginal birth
- 9) Best diagnosed with USG for 1st trimester
- Anecephaly
 - NT defects
 - Meningocele
 - None of the above
- 10) A mother comes with history of antenatal fetal death in her first child. What is the amount of folic acid you will prescribe for prenatal counselling
- 4 mcg per day
 - 40 mcg per day
 - 400 mcg per day
 - 4000 mcg per day
- 11) Procedures done for cervical incompetency include
- Mac Donald suture
 - Porander operation
 - Khanna's sling operation
 - Abdominal sling operation
- 12) For MTP consent should be obtained from
- The male partner
 - The male as well as female partner
 - The female partner
 - No consent required
- 13) Deep transverse arrest is seen in
- Android pelvis
 - Gynecoid pelvis
 - Anthropoid pelvis
 - Platypelloid pelvis
- 14) A woman presents with 6 weeks amenorrhea and bleeding PV. Which screening test is useful for above
- Urine for pregnancy test
 - BETA HCG
 - USG
 - Laparotomy

15) In which part of the fallopian tube will an ectopic pregnancy have the longest survival

- a. Isthmus
- b. Ampulla
- c. Cornua
- d. Interstitium

16) Serum progesterone levels which excludes ectopic pregnancy are

- a. 2-4 ng/ml
- b. 5-10 ng/ml
- c. 10-20 ng/ml
- d. More than 25 ng/ml

17) In the pelvic inlet which is the shortest AP diameter

- a. True conjugate
- b. Obstetric conjugate
- c. Anatomical conjugate
- d. Bispinous diameter

18) Benign condyloma is associated with which condition. Infection by

- a. HPV type II
- b. PHPV 16
- c. HPV 16
- d. HPV 18

19) Engaging diameter in brow presentation is

- a. Submento bregmatic
- b. Mento vertical
- c. Occipito frontal
- d. Sub occipito bregmatic

20) Symphysiotomy is indicated in

- a. Contraction of brim
- b. Contraction of cavity
- c. Contraction of outlet
- d. All of the above
