

Dr. NTR UNIVERSITY OF HEALTH SCIENCES :: AP :: VIJAYAWADA – 520 008

FINAL M.B.B.S. PART – II – DEGREE EXAMINATION – DECEMBER, 2024

OBSTETRICS & GYNAECOLOGY – PAPER II

SET-D (Multiple Choice Questions)

Time: 20 Minutes

Max. Marks: 20

Note: Answer all questions

SECTION – I (MCQs – 20 MARKS)**1x20=20**

- 1) The most common site of ectopic pregnancy
 - a. Ampullary portion of Fallopian tube
 - b. Interstitium
 - c. Isthmus
 - d. Abdomen
- 2) Uterine leiomyoma is least likely to undergo
 - a. Malignant changes
 - b. Calcification
 - c. Hyaline changes
 - d. Red degeneration
- 3) Which of the following structures is most susceptible to unintentional damage during hysterectomy
 - a. Uterine artery
 - b. Urinary bladder
 - c. Ureter
 - d. Urethra
- 4) Ovary develops from which of the following
 - a. Genital ridge
 - b. Mullerian duct
 - c. Genital tubercle
 - d. Wolffian duct
- 5) A pregnant female with 16 weeks POG presented with bleeding. Upon USG fibroid detected, management is
 - a. Conservative
 - b. Curative
 - c. Lap myomectomy
 - d. Hysterectomy
- 6) What does satisfactory colposcopy means
 - a. Visualization of entire transformation zone
 - b. Visualization of endocervix
 - c. Visualization of ectocervix
 - d. Visualization of fornices
- 7) All of the following are the indicators for myomectomy in a case of fibroid uterus except
 - a. Associated infertility
 - b. Pressure symptoms
 - c. Recurrent pregnancy loss
 - d. Red degeneration

8) Condition that can be managed on laparoscopy

- a. Endometriosis
- b. Ca cervix
- c. Dysfunctional uterine bleeding
- d. Pyometra

9) A 16 year old female presents with primary amenorrhea. Examination shows a short blind vagina, with an absent uterus. The next investigation of choice is

- a. Karyotyping
- b. IVP
- c. Gonadotropin levels
- d. Serum prolactin

10) Gland homologous to prostate in female is

- a. Gartner's gland
- b. Skene gland
- c. Bartholine gland
- d. Cowper gland

11) In Stein – Leventhal syndrome which hormone is increased

- a. LH
- b. FSH
- c. GnRH
- d. Progesterone

12) Precocious puberty associated with bony dysplasia and café au lait spots in the skin is seen in

- a. Frohlich's syndrome
- b. Alport syndrome
- c. McCune – Albright syndrome
- d. Laurence – Moon – Biedl syndrome

13) Which of the following is true about wandering fibroid

- a. Attached to uterus
- b. Attached to uterus and surroundings
- c. Attached to surrounding viscera
- d. Neither attached to uterus nor to surrounding viscera

14) The first sign of sexual maturation in girl is

- a. Menarche
- b. Pubarche
- c. Enlargement of breast
- d. Maturation of breast

15) Fibromyoma is seen in age group

- a. Reproductive
- b. Early adolescence
- c. Late post menopausal
- d. Prepubertal

16) Contraceptive to be avoided in epilepsy

- a. OCPs
- b. Condom
- c. IUCD
- d. Mirena

17) Longest part of fallopian tube

- a. Ampulla
- b. Isthmus
- c. Intramural
- d. Fimbriae

18) In uncontrollable PPH, all of the following arteries can be ligated except

- a. Uterine artery
- b. Anterior division of internal iliac artery
- c. Posterior division of internal iliac artery
- d. Ovarian artery

19) Clue cells are

- a. Epithelial cells
- b. Lymphocytes
- c. Neutrophils
- d. Macrophages

20) Cervix to body ratio in a prepubertal girl is

- a. 1:1
- b. 2:1
- c. 1:2
- d. 3:1
