

MEDICAL**[MBBS 0625]****JUNE 2025****Sub. Code :6085****M.B.B.S. DEGREE EXAMINATION****(For the candidates admitted from the Academic Year 2019-2020)****THIRD PROFESSIONAL PART II SUPPLEMENTARY EXAMINATION****PAPER V – OBSTETRICS & GYNAECOLOGY*****Q.P. Code: 526085*****Time: 20 Minutes****Maximum : 20 Marks****Answer All Questions****Shade the correct answer with Black or Blue ink ball point pen****SET - B****III. Multiple Choice Questions:****(20 x 1 = 20)**

1. Post partum decidual secretions are known as
 - A) Lochia
 - B) Bleeding per vagina
 - C) Vasa previa
 - D) Decidua capsulari
2. During foetal life, Maximum growth is caused by
 - A) Growth hormone
 - B) Insulin
 - C) Cortisol
 - D) Thyroxine
3. Intermittent cell predominance is a cytology seen in
 - A) Pregnancy
 - B) Post Menstrual
 - C) Post ovulatory
 - D) Pre menstrual
4. Specific gravity of amniotic fluid is
 - A) 1.004 to 1.006
 - B) 1.006 to 1.008
 - C) 1.008 to 1.010
 - D) 1.016 to 1.018
5. Rate of amniotic fluid production is
 - A) 300 cc / hour
 - B) 500 cc/ hour
 - C) 700 cc/ hour
 - D) 900 cc /hour
6. Most common cause of thrombocytopenia in pregnancy is
 - A) Immune
 - B) Idiopathic
 - C) Infantile
 - D) Gestational
7. Alpha and Beta segments are not seen in
 - A) FSH
 - B) HCG
 - C) Prolactin
 - D) Insulin
8. Living ligature of uterus is
 - A) Endocentrium
 - B) Myometrium inner layer
 - C) Myometrium middle layer
 - D) Pericentrium
9. Ritgen manoeuvre is performed in shoulder dystocia
 - A) Delivery of head in breech presentation
 - B) Delivery of legs in breech presentation
 - C) Delivery of head in normal delivery
 - D) Delivery of legs in normal delivery

10. In BISHOP score, all are included except
A) Effacement of cervix B) Dilatation of cervix
C) Station of head D) Inter spinous diameter
11. Likely size of uterus 8 weeks post-partum is
A) 100 gms B) 300 gms C) 700 gms D) 900 gm
12. Most common cause of first trimester abortion is
A) Chromosomal abnormalities B) Syphilis
C) Rh incompatibility D) Cervical incompetence
13. According to Hellin's law, Chances of twins in pregnancy are
A) 1 in 80 B) 1 in 70 C) 1 in 60 D) 1 in 50
14. Placenta previa can cause all except:
A) Painless bleeding B) Causeless bleeding
C) Recurrent bleeding D) First trimester bleeding
15. All of the following can cause DIC except:
A) Diabetes mellitus B) Amniotic fluid embolism
C) Intra uterine death D) Abruptio placentae
16. Submento bregmatic diameter is
A) 9.5 cm B) 11.5 cm C) 13.5 cm D) 12.5 cm
17. Smallest diameter of the pelvis is
A) Interspinous diameter B) Diagonal conjugate
C) True conjugate D) Intertuberous diameter
18. Regimen followed in expectant management of placenta previa is
A) Liley's method B) Crede's method
C) Macaffee and Johnson's method D) Brechett Andrew's method
19. Clotting factor which is not increased in pregnancy is
A) Factor 2 B) Factor 7 C) Factor 10 D) Factor 11
20. Indication of induction of labour is
A) Placenta previa B) Gestational hypertension at term
C) Breech presentation D) Maternal heart disease

[MBBS 0625]