

## www.FirstRanker.com

www.FirstRanker.com

Code No: F3104

**R09** 

## JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD MCA I Semester Examinations, January - 2020 PROBABILITY AND STATISTICS

Time: 3hrs Max.Marks:60

Answer any five questions All questions carry equal marks

---

- 1.a) There are 25 cards numbered 1 to 25. Two cards are drawn. Find the probability that their sums is i) even ii) a multiple of 3.
  - b) There are three boxes. I contains- 10 light bulbs out of which 4 are defective II, contains- 6 light bulbs out of which 1 is defective, III contains- 8 light bulbs out of which 3 are defective. A box is chosen at random and a bulb is selected. If it is defective then find the probability that it is from:

i) Box- I

ii) Box-II

iii) Box-III.

[6+6]

2.a) A random variable X has the following distribution

X	1	2	3	4	5
P(X = x)	1	1	1	-1	1
	2	4	8	16	16

Find: i) The mean

ii) variance.

iii) P(1< X< 4)

- b) The probability of man hitting a target is 1/3. If he fires 6 times, find the probability of hitting: i) At the most 5 times ii) At least 5 times iii) Exactly once. [6+6]
- 3.a) Average number of accidents on any day on a national highway is 1.6. Determine the probability that the number of accidents is i) at least one ii) at the most one
  - The marks obtained by 1000 students is normally distributed with mean 78 % and Standard deviation 11%. Determine
    - How many get more than 90%.
    - How many students get between 75% and 90%.

[6+6]

- A population consists of five numbers 5,10,15,20 .Consider all Samples of size two which can be taken with replacement from this population. Find:
  - a) The population mean.
  - b) The population Standard deviation.
  - c) The mean of the sampling distribution of means.
  - d) Standard deviation of the sampling distribution of means.

[12]

- 5.a) A Random sample of 100 items is taken from a population whose standard deviation is 5.1 The mean of the sample is 21.6. Construct 95% confidence interval for the mean.
- b) A sample of 121 students is found to have a mean weight of 68 kgs. Can this be regarded as a sample from a population with mean weight 75 kgs and standard deviation 31 kgs?
  [6+6]





## www.FirstRanker.com

## www.FirstRanker.com

- 6.a) A simple sample of heights of 6400 Englishmen has a mean of 67.85 and Standard deviation of 2.56 inches. While a simple sample of heights of 1600 Australian has a mean of 68.55 and a S.D. of 2.52 inches Test the significance between the two mean.
  - b) 400 articles from a factory are examined and 3% are found to be defective. 600 similar articles from a second factory are found to be 2 % defective. Test the significance between the differences of two proportions at 5% level. [6+6]
- Two random sample are drawn from two normal populations are as follows.

Sample I	ı				27			
Sample II	16	16	20	27	26	25	21	

Test whether two populations have been drawn from the same normal population. [12]

 The marks obtained by 10 students in Mathematics and Statistics are given below. Find the Coefficient of correlation between the two subjects and the regression lines. [12]

Marks in Maths	75	30	60	80	53	35	15	40	38	48
Marks in Statistics	85	45	54	91	38	63	35	43	45	44



