



- Q.3 Most common site of cervix carcinoma is:
- CA endometrium
  - CA vagina
  - CA cervix
- Q.4 Narrowest Part of Fallopian Tube is:
- Interstitial portion
  - Isthmus
  - Infundibulum
  - Ampulla
- Q.5 Peg cells are seen in:
- Vagina
  - Vulva
  - Ovary
  - Tubes
- Q.6 Strawberry vagina is seen in:
- Candida albicans
  - H. vaginalis
  - Syphilis
  - Trichomonas vaginalis
- Q.7 Clue cells are seen in:
- Bacterial vaginosis
  - Candidiasis
  - Trichomoniasis
  - Gonorrhoea
- Q.8 Mirena is:
- Used in abortion
  - Antiprogesterone
  - Progesterone IUCD
  - Hormonal implant
- Q.9 Amount of hormone in MALA D is:
- 30ug
  - 50ug
  - 10ug
  - 80ug
- Q.10 Most common type of HPV causing CA cervix is:
- 16 & 18
  - 1 & 33
  - 6 & 11
  - 2 & 14
- Q.11 Cervix Carcinoma arises from:
- Squamocolumnar junction
  - Isthmus
  - Cervical lip
  - Infertility
  - Menstrual irregularity
- Q.12 Red degeneration of fibroid is most common in:
- 2<sup>nd</sup> Trimester
  - 3<sup>rd</sup> Trimester
  - Puerperium s
  - First Trimester
- Q.13 Absolute contraindication of Hormone Replacement therapy is:
- Thrombosis
  - Fibrocystic Disease
  - Fibroadenoma
  - Haemorrhage

- Q.14 Stage – II
- Q.15 Stage – III
- Q.16 Stage – IV
- Q.17 Most common ovarian tumour in pregnancy is:
- Mucinous cyst adenoma
  - Dermoid cyst
  - Metastasis
  - Dysgerminoma
- Q.18 Fern test is due to:
- Presence of NACL under progesterone effect
  - Presence of NACL under estrogenic effect
  - LH/FSH
  - Mucus secretion by glands
- Q.19 Which intrauterine device (IUD) has life span of 10 years:
- CUT 380 A
  - CUT 200
  - NOVA T
  - Multilod
- Q.20 During Pomroy's method of sterilization which portion of tube is ligated:
- Isthmus
  - Ampullary
  - Isthmo ampullary
  - Cornual
- Q.21 Polymenorrhoea means:
- Menses < 21 days
  - menses > 35 days
  - Painful menses
  - DUB
- Q.22 Treatment of DUB in young female is:
- Hormones
  - Radiotherapy
  - Hysterectomy
  - D & C
- Q.23 In fibroid which is not seen:
- Amenorrhoea
  - Pelvic Mass

P.T.O.