

**MBBS III (Third) Professional Part-2 Examination
2018-19**

Course Code: MBS403

Paper ID: 03118404

Obstetric and Gynaecology - I

Time: 2 Hours 10 Minutes

Max Marks: 30

Note: Attempt all questions. Draw proper diagrams to support your answer.

Part 'B'

1. Classify hypertensive disorders in pregnancy and how will you manage a case of primigravida with 36 weeks of gestation with severe preclampsia (7)
2. Define postpartum hemorrhage (PPH). Enumerate the causes of PPH. Discuss the management of atonic PPH. (7)

Part 'C'

3. Write short notes on the following: (4x4=16)
 - a) Non Stress Test (NST)
 - b) Caput succedaneum
 - c) Partogram.
 - d) Threatened preterm labour

MBBS III (Third) Professional Part-2 Examination 2018-19

Roll No.

Student's Signature

Student's Name

Invigilator's Signature

Course Code: MBS403

Paper ID: 03118404

Obstetric and Gynaecology - I

Part 'A'

Time: 20 Minutes

Max Marks: 10

- Note:**
1. Attempt all questions and return this part of the question paper to the invigilator after 20 Minutes.
 2. Please tick (✓) correct one only. Cutting, overwriting or any other marking are not allowed.
 3. For answering please use Ball- pen only.

Which of the following is seen in the infant of a diabetic mother:

- Hyper kalemia
- Hyper calcemia
- Macrocytic anaemia
- Polycythemia

Q.3 Twin peak sign seen in:

- Monochorionic diamniotic twins
- Dichorionic mono amniotic twins
- Conjoined twins
- Diamniotic dichorionic twins

Q.4 Which test differentiates maternal and fetal blood cells:

- APT test
- Kleihauer test
- Bubble test
- Lilys test

Q.5 Choriocarcinoma commonly metastasize to:

- Brain
- Lung
- Ovary
- Vagina

Q.6 All of the following drugs have been used for medical abortion except:

- Mefepristone
- Misoprostol
- Methotrexate
- Atosiban

Q.7 In anticipated preterm delivery the dose of dexamethasone given to mother is:

- 12mg 12hourly 2 doses
- 12mg 24hourly 4 doses
- 6mg 24hourly 2 doses
- 6mg 12 hourly 4 doses

Q.8 Spalding's sign is seen in:

- Fresh Still born
- Live born
- Premature
- Macerated still born

Q.9 Best method to deliver arms in breech:

Q.17 Shortest diameter of fetal skull is:

- Suboccipito-bregmatic
- Bitemporal
- Suboccipitofrontal
- Biparietal

Q.18 Induction of labour is usually avoided in:

- Malpresentation
- PIH
- Diabetes
- IUFD

Q.19 Most commonly used antihypertensive in pregnancy is:

- Nifedipine
- Methyldopa
- Frusemide
- Hydralazine

Q.20 Commonest cause of breech presentation is:

- Prematurity
- Diabetes mellitus
- Postmaturity
- Contracted pelvis

Q.10 Pain in early labour is Limited to dermatomes:

- T10-L₁
- T11-L₂
- L4-L5
- L2-L3

P.T.O.

Q.11 Early deceleration occur due to:

- Fetal head compression
- Umbilical cord compression
- Fetal hypoxia
- Congenital heart block

Q.12 Bishop's scoring includes all except:

- Cervical consistency
- Cervical length
- Station of head
- Cervical colour

Q.13 The therapeutic serum level of magnesium during magnesium sulphate therapy for eclampsia is:

- 4-7meq/L
- 7-10 meq/L
- 10-12 meq/L
- 12-15 meq/L

Q.14 Most common cause of secondary PPH is:

- Uterine atony
- Rupture uterus
- Retained placental bits
- Vaginal lacerations

Q.15 The period of gestation at which the crown rump length is used to estimate the gestational age is:

- 4-6 weeks
- 6-12 weeks
- 15-20 weeks
- 20-24 weeks

Q.16 Number of antenatal care in an uncomplicated pregnancy should be minimum:

- 4 times
- 8 times
- 12 times
- 16 times