

**Instructions:**

All questions are compulsory

1) MCQ question paper should be conducted and completed in first 30 min.

2) Fill (dark) the appropriate empty circle against the question number once only.

3) Use blue/black ball point pen only.

4) Each MCQ carries half mark.

5) Students will not be allotted mark if he/she overwrites/ strikes or put white ink on the cross once marked on MCQ.

Time : 30 Minutes

Maximum Marks : 8

Q. 1 Total MCQs : 16

16 X ½ = 8

- 1) Most bacteria are destroyed in the phagolysosome except  
a. Staphylococcus    b. Mycobacterium Lepra    c. Neisseria Gonorrhoeae    d. Streptococcus Pyogene
- 2) All are super antigens except :-  
a. H antigen    b. Staphylococcal enterotoxin    c. Exfoliative Toxin    d. Some viral protein
- 3) Molecular weight of IgA is :-  
a. 150000    b. 180000    c. 160000    d. 190000
- 4) Example of flocculation test are :-  
a. VDRI. & Kahn Test    b. VDRL & RPR test    c. RPR & Kahn    d. None of the above
- 5) Activation of C3 in alternate pathway occur  
a. Without help of C142    b. With the help of C2    c. With the help of C142    d. None of the above
- 6) To demonstrate fungal hyphae the concentration of KOH which is used is :-  
a. 20%    b. 10%    c. 40%    d. 30%
- 7) Cowdry type A inclusion bodies are seen in  
a. Rhabdo Virus    b. Adenovirus    c. Poliovirus    d. Herpes virus
- 8) All are RNA viruses except :-  
a. Adeno Virus    b. Toga Virus    c. Rhabdo Virus    d. Corona Virus
- 9) Amniotic sac inoculation is mainly used for isolation of :-  
a. Chlamydia    b. Rickettsiae    c. Influenza virus    d. Rabies Virus
- 10) Continuous cell lines are all except :-  
a. Hela    b. HEP-2    c. IIL-8    d. Vero
- 11) Biphasic (Saddle Back) fever is seen in :-  
a. Dengue    b. Chikungunya    c. Typhoid Fever    d. Rickettsial Fever
- 12) The infective dose of girdia lamblia is :-  
a. 1-10 Cysts    b. 10-100 Cysts    c. 100-1000 Cysts    d. None of the above
- 13) Schuffner's dots are present in :-  
a. P. Vivax    b. P. Malariae    c. P Falciparum    d. P. Ovale
- 14) In microfilarae of W bancrofti tail tip contains :-  
a. 2 nuclei    b. no nuclei    c. 4 nuclei    d. None of the above
- 15) In toxoplasma gondii's life cycle following are intermediate host except :-  
a. Cat    b. Human being    c. Rat    d. Cattle
- 16) Trichophyton produces infection of :-  
a. Hair & Skin    b. Skin & Nail    c. Hair, Skin & Nail    d. Only skin