



Instructions

- 1) All questions are compulsory
- 2) MCQ question paper should be conducted and completed in first 30 min.
- 3) Fill (dark) the appropriate empty circle against the question number once only.
- 4) Use blue/black ball point pen only.
- 5) Each MCQ carries half mark.
- 6) Students will not be allotted mark if he/she overwrites/ strikes or put white ink on the cross once marked on MCQ.

Time : 30 Minutes

Maximum Marks : 8

16 X ½ = 8

Q. 1

Total MCQs : 16

- 1 If a drug has short duration of action, the action can be prolonged by all except-
 a. Enteric coating b. Interfering with renal excretion c. Inhibiting drug metabolism d. Intravenous administration
- 2 Following are pro drugs except-
 a. Enalapril b. Ranitidine c. Levodopa d. Sulindac
- 3 Which of the following factor is most commonly related to the selection of dosage
 a. Age b. Sex c. Weight d. Body surface area
- 4 After prolonged treatment following drugs should be gradually withdrawn except-
 a. Clonidine b. Alpha methyl dopa c. Chlorthiazide d. Prednisolone
- 5 Following are the effect of the Acetylcholine except-
 a. Constriction of bronchi b. Increase of heart rate c. Increased peristalsis d. Miosis
- 6 Which of the following drug is most suitable for the treatment of Myasthenia Gravis-
 a. Physostigmine b. Atropine c. Pyridostigmine d. Pralidoxime
- 7 Noradrenaline release is controlled by-
 a. Presynaptic autoreceptors b. Postsynaptic receptors c. Nucleic receptors d. Silent receptors
- 8 Which antihypertensive agent is used for management of hypertensive emergencies-
 a. Prazosin b. Glyceryl trinitrate c. Verapamil d. Atenolol
- 9 Ankle oedema produced by Amlodipine is -
 a. Permanent b. Resistant to diuretic c. Relieved by diuretics d. Observed in all patients
- 10 Fenofibrates actions exclude-
 a. Reduce triglyceride level b. Increase HDL-C level c. Decrease lipoprotein lipase activity d. Decrease LDL level
- 11 Spironolactone and ACE inhibitors are not used concurrently because of danger of-
 a. Hyperglycemia b. Hyperkalemia c. Hypokalemia d. Hypoglycemia
- 12 All of the following antihistaminic has anticholinergic action except-
 a. Cetirizine b. Promethazine c. Diphenhydramine d. Cyproheptadine
- 13 Drugs used for prophylaxis of migraine are all except-
 a. β -blocker b. NSAID c. Tricyclic antidepressant d. Flunarizine a CCB
- 14 Regarding peripherally acting muscle relaxants all are correct except-
 a. Given IV b. Used for short term purposes c. Blocks neuromuscular transmission d. Causes CNS depression
- 15 Sucralfate acts by -
 a. Acting against H.Pylori b. Physical barrier c. As antacid d. Decreasing acid secretion
- Stimulant purgative is contraindicated in-
 a. Pregnancy b. Preparation of bowel surgery c. With anthelmintics d. Atonic constipation

