



Instructions:

- 1) All questions are compulsory
- 2) MCQ question paper should be conducted and completed in first 30 min.
- 3) Fill (dark) the appropriate empty circle against the question number once only.
- 4) Use blue/black ball point pen only.
- 5) Each MCQ carries half mark.
- 6) Students will not be affected mark if he/she overwrites/ strikes or put white ink on the cross once marked on MCQ.

Maximum Marks :

8

16 X 1/2 = 8

Time : 30 Minutes

Total MCQs : 16

Q. 1

- 1 Prostaglandin useful for the prevention of duodenal ulcer is ☒ a. Dinoprost ☐ b. Misoprostol ☐ c. Alprostadil ☐ d. Carboprost
- 2 Drugs used in prophylaxis of migraine are all except ☒ a. Propranolol ☐ b. Flunarizine ☐ c. Topiramate ☒ d. Levettiracetam
- 3 Analgesic that itself can cause headache as a side effect is ☒ a. Indomethacin ☐ b. Mephenamic acid ☐ c. Piroxicam ☐ d. Aspirin
- 4 Drug of choice for acute gout ☒ a. Colchicine ☐ b. Indomethacin ☐ c. Allopurinol ☐ d. Dexamethasone
- 5 Most cardiotoxic local anesthetic is ☐ a. Procaine ☒ b. Bupivacaine ☐ c. Prilocaine ☐ d. Tetracaine
- 6 Muscle relaxant of choice in liver disease is ☐ a. Atracurium ☒ b. Pipecuronium ☐ c. Rocuronium ☐ d. Vecuronium
- 7 Dissociative anesthesia is produced by ☒ a. Ketamine ☐ b. Fentanyl ☐ c. Propofol ☐ d. both a and b
- 8 Eutectic mixture of local anesthetic (EMLA) cream is ☒ a. Bupivacaine 2.0% + Prilocaine 2.5% ☐ b. Lidocaine 2.5% + Prilocaine 2.5% ☐ c. Lidocaine 2.5% + Prilocaine 5% ☐ d. Bupivacaine 0.5% + Lidocaine 2.5%
- 9 Treatment of choice for chancroid is ☒ a. Penicillin ☐ b. Chloramphenicol ☐ c. Tetracycline ☐ d. Erythromycin
- 10 Longest acting sulphonamide is ☒ a. Sulfadiazine ☐ b. Sulfadoxine ☐ c. Sulfamethoxazole ☐ d. Sulfamethiazole
- 11 ATT causing orange colored urine is ☒ a. Rifampicin ☐ b. Isoniazid ☐ c. Streptomycin ☐ d. Pyrazinamide
- 12 Zidovudine causes ☐ a. Neurotoxicity ☐ b. Nephrotoxicity ☒ c. Neutropenia ☐ d. Ototoxicity
- 13 Conversion of T₄ to T₃ is inhibited by all except ☐ a. Propranolol ☐ b. Propylthiouracil ☒ c. Amiodarone ☐ d. Methimazole
- 14 Contraceptive that should be avoided in epilepsy is ☒ a. Oral contraceptives ☐ b. Condoms ☐ c. Intrauterine contraceptive devices ☐ d. Post-coital pills
- 15 Insulin causes all of the following except ☒ a. Glycogenesis ☐ b. Glycolysis ☐ c. Lipogenesis ☐ d. Ketogenesis
- 16 Probenecid interacts with ☒ a. Streptomycin ☐ b. Ampicillin ☐ c. Vancomycin ☐ d. Erythromycin

