



- Instructions:
- All questions are compulsory
 - Draw diagrams wherever necessary
 - Answers of Questions and Sub questions must be written strictly according to the serial order of question paper.
 - MCQ has to be answered in theory answer book
 - Please write MCQ answer neatly and in serial order with black or blue pen in brackets for example:- 1. (a) 2. (b)
 - MCQ has to be answered only once, any kind for repetition or cutting or erasing or whitener will be consider as malpractice.
- Such answers will not be counted in the marks and action will be taken according to UFM rules of University

Q. 1 Total MCQs : 20

20 X 1/2 = 10

- All are differential diagnosis of thunderclap headache except
a. Encephalitis b. Ischemic stroke c. Subarachnoid hemorrhage d. Meningitis
- Which is not a cardinal feature of Parkinson's disease
a. Bradykinesia b. Rigidity c. Rest tremors d. Headache
- Intravenous thrombolysis in ischemic stroke can be done
a. In 8 hrs b. 12hrs c. 3-6hrs d. 12-24hrs
- Which of the following is not an indication of intravenous thrombolysis
a. Ischemic stroke b. Symptoms in 4.5 hrs c. Age < 18yrs d. CT scan showing no hemorrhage or edema
- Which drug is given to reduce serum triglycerides
a. Fibrates b. Statins c. Ezetimide d. Niacin
- Thrombolysis can be done in treatment of acute MI, if patient comes in
a. 3hrs b. 6hrs c. 12hrs d. 24hrs
- Which of the following is not recommended for the treatment of Raynaud's phenomenon?
a. Nifedipine b. Sildenafil c. Doxazosine d. Telmisartan
- Following are the risk factors involved in asthma
a. Gender b. Passive smoking c. Ethnicity d. All
- Which is not a complication of oral theophylline
a. Diuresis b. Nausea c. Bradycardia d. Seizures
- Factors increasing clearance of theophylline except
a. Smoking b. High protein diet c. Viral infection d. Drugs like rifampicin
- Which of the following is a cause of demyelinating neuropathy
a. CIDP b. Hyperglycemia c. Uremia d. Antiretroviral drugs
- CSF finding in guillian Barre syndrome
a. Normal cells with increased protein b. Increased protein with normal cells c. Increased cell with increased protein d. Normal cell with normal protein
- The dome of diaphragm is elevated in
a. Emphysema b. Typhoid fever c. COPD d. Liver abscess
- Tubercular ascitis has following feature except
a. Increased ADA b. Increased protein c. Hemorrhagic effusion d. High SAAG

13. The following are present in anterior mediastinum except
 a. Thymoma b. Teratoma c. Neuroblastoma d. Cyst
14. Which of the following is not a cause of low pleural fluid sugar
 a. Dressler's syndrome b. Tubercular effusion c. Erysipelas d. Malignant effusion
15. Fulminant hepatitis is common in
 a. HAV b. HBV c. HCV d. HEV
16. Which of the following is transmitted by fecal-oral route
 a. HBV b. HDV c. HCV d. HEV
17. Dane particle is related to
 a. HEV b. HANB c. HBV d. None
18. HCV is associated with
 a. LKM antibody b. Autoimmune cirrhosis c. Antimitochondrial antibody d. None

Q.2 Long Answer Question

4X 5 = 20

- Describe clinical feature and complication of rheumatic heart disease
- Diagnosis and treatment of acute bronchial asthma
- Describe cirrhosis of liver and its complication
- Diagnosis and management of acute meningitis

Q.3 Short Answer Question

4X 3 = 12

- Treatment of rheumatoid arthritis
- Complication of sickle cell disease
- Treatment of status epilepticus
- Clinical feature and treatment of iron deficiency anemia

Q.4 Very Short Answer Question

9X 2 = 18

- Four causes of ascitis
- Four causes of extensor plantar response
- Glass glow coma scale
- Management of acute myeloid leukemia
- Diagnosis of pulmonary tuberculosis
- Four causes of atrial fibrillation
- Differential diagnosis of diastolic murmur at mitral area
- Treatment of chronic hepatitis B
- Paraneoplastic syndrome