



Topic:- PHY MSC S2

1) An atomic transition line with wavelength 350 nm is observed to be split into three components in a spectrum of light from a sunspot. Adjacent components are separated by 1.7 pm. Determine the strength of the magnetic field in the sunspot

[Question ID = 6662]

1. 3 T [Option ID = 26642]
2. 0.03 T [Option ID = 26643]
3. 3.3 T [Option ID = 26644]
4. 0.3 T [Option ID = 26645]

Correct Answer :-

- 0.3 T [Option ID = 26645]

2) which one of the following is correct in respect of an electron and a proton having a same de-Broglie wavelength of 2 Å

[Question ID = 6663]

1. Both have same kinetic energy [Option ID = 26646]
2. Both have same velocity [Option ID = 26647]
3. Both have same momentum [Option ID = 26648]
4. The kinetic energy of proton is more than that of electron [Option ID = 26649]

Correct Answer :-

- Both have same momentum [Option ID = 26648]

3) If  $r_p$  &  $r_H$  are the radius and  $E_p$  &  $E_H$  are the energy of an electron in the  $n^{\text{th}}$  orbit of positronium atom and hydrogen atom respectively, then

[Question ID = 6664]

1.  $r_p = 2 r_H$  and  $E_p = E_H/2$

[Option ID = 26650]

2.  $r_p = 2 r_H$  and  $E_p = 2 E_H$

[Option ID = 26651]

3.  $r_p = 2 r_H$  and  $E_p = E_H/4$

[Option ID = 26652]

4.  $r_p = r_H$  and  $E_p = 2 E_H$

[Option ID = 26653]

Correct Answer :-

- $r_p = 2 r_H$  and  $E_p = E_H/2$

[Option ID = 26650]

4) An X-ray beam of wavelength 0.16 nm is incident on a set of planes of a certain crystal. The first Bragg reflection is observed for an incidence angle of  $30^\circ$ . What is the corresponding inter planar spacing?

[Question ID = 6669]

1. 0.16 nm [Option ID = 26670]
2. 0.67 nm [Option ID = 26671]
3. 1.02 nm [Option ID = 26672]
4. 0.89 nm [Option ID = 26673]

Correct Answer :-

- 0.16 nm [Option ID = 26670]

5) What is the velocity of conduction electron of silver having Fermi energy 5.52 eV

[Question ID = 6670]

1.  $1.39 \times 10^6$  m/s [Option ID = 26674]
2.  $2.39 \times 10^6$  m/s [Option ID = 26675]
3.  $0.89 \times 10^6$  m/s [Option ID = 26676]
4. 0 [Option ID = 26677]

Correct Answer :-

- $1.39 \times 10^6$  m/s [Option ID = 26674]

6) Given that a piece of n-type silicon contains  $8 \times 10^{21} \text{ m}^{-3}$  phosphorus impurity atoms, calculate the carrier concentration of silicon at room temperature. Given that the intrinsic electron concentration of silicon at room temperature is  $1.6 \times 10^{16} \text{ m}^{-3}$



10.  $10^{10} \text{ m}^{-3}$  [Option ID = 26671]  
 11.  $10^{11} \text{ m}^{-3}$  [Option ID = 26672]  
 12.  $10^{12} \text{ m}^{-3}$  [Option ID = 26673]  
 13.  $1.5 \times 10^{10} \text{ m}^{-3}$  [Option ID = 26680]  
 14.  $3.2 \times 10^{11} \text{ m}^{-3}$  [Option ID = 26681]

Correct Answer :-

- $3.2 \times 10^{10} \text{ m}^{-3}$  [Option ID = 26678]

7) The dispersion relation for a one-dimensional monoatomic lattice chain is given by the equation,  $\omega = \frac{2}{a} v_s \left| \sin\left(\frac{Ka}{2}\right) \right|$ , where, 'a' is the interatomic spacing,  $K = \frac{2\pi}{\lambda}$  and  $v_s$  has the dimension of velocity. The relation between the phase velocity  $V_p$  and group velocity  $V_g$  in the long wavelength limit is given by

[Question ID = 6674]

- $V_p = V_g$   
[Option ID = 26690]
- $V_p = 2V_g$   
[Option ID = 26691]
- $V_p = V_g/2$   
[Option ID = 26692]
- $V_p = V_g$   
[Option ID = 26693]

Correct Answer :-

- $V_p = V_g$   
[Option ID = 26690]

8) The largest wavelength of light falling on double slits separated by  $1.5 \mu\text{m}$ , for which there is a first order maximum is in the,

[Question ID = 6676]

- ultraviolet range [Option ID = 26698]
- visible range [Option ID = 26699]
- infrared range [Option ID = 26700]
- X-ray range [Option ID = 26701]

Correct Answer :-

- infrared range [Option ID = 26700]

9) In a multi-stage R-C coupled amplifier, the coupling capacitor

[Question ID = 6680]

- limits the low frequency response [Option ID = 26714]
- limits the high frequency response [Option ID = 26715]
- reduces the amplitude of input signal [Option ID = 26716]
- blocks d.c. component without affecting the frequency response [Option ID = 26717]

Correct Answer :-

- blocks d.c. component without affecting the frequency response [Option ID = 26717]

10) An AM transmitter is coupled to an aerial. The input current is found to be 5 A. With modulation the current value increases to 5.9 A. The depth of modulation is

[Question ID = 6681]

- 83.4% [Option ID = 26718]
- 88.6% [Option ID = 26719]
- 78.2% [Option ID = 26720]
- 74.3% [Option ID = 26721]

Correct Answer :-

- 88.6% [Option ID = 26719]

11) Hexadecimal equivalent of a digital number 10011101 is

[Question ID = 6683]

- H913 [Option ID = 26726]
- 9D [Option ID = 26727]
- AE [Option ID = 26728]
- 157 [Option ID = 26729]

Correct Answer :-

- 9D [Option ID = 26727]

12) If the doping concentration in a Si-Zener diode is increased, the Zener breakdown voltage

[Question ID = 6684]



Correct Answer :-

- Decreases [Option ID = 26730]

13) Which one of the following is an example of doubly magic nuclei

[Question ID = 6685]

1.  $^{18}\text{O}$

[Option ID = 26734]

2.  $^{48}\text{Ca}$

[Option ID = 26735]

3.  $^{124}\text{Sn}$

[Option ID = 26736]

4.  $^{204}\text{Pb}$

[Option ID = 26737]

Correct Answer :-

- $^{48}\text{Ca}$

[Option ID = 26735]

14) Which radiation has maximum ionization power?

[Question ID = 6686]

1. Alpha [Option ID = 26738]

2. Beta [Option ID = 26739]

3. Neutron [Option ID = 26740]

4. Gamma [Option ID = 26741]

Correct Answer :-

- Alpha [Option ID = 26738]

15) For beta-minus decay, which statement is TRUE?

[Question ID = 6688]

1. Daughter nuclide atomic mass ( $A_D$ ) is more than that of the parent nuclide atomic mass ( $A_P$ ) [Option ID = 26746]

2. Daughter nuclide atomic number ( $Z_D$ ) is same that of the parent nuclide atomic number ( $Z_P$ ) [Option ID = 26747]

3. Daughter nuclide neutron number ( $N_D$ ) is less than that of the parent nuclide neutron number ( $N_P$ ) [Option ID = 26748]

4. Daughter nuclide neutron number ( $N_D$ ) is same that of the parent nuclide neutron number ( $N_P$ ) [Option ID = 26749]

Correct Answer :-

- Daughter nuclide neutron number ( $N_D$ ) is less than that of the parent nuclide neutron number ( $N_P$ ) [Option ID = 26748]

16) The probability that student A solves the problem is  $1/2$ , and that of B is  $2/3$ . What is the probability that the problem is solved?

[Question ID = 6689]

1.  $4/6$

[Option ID = 26750]

2.  $1/3$

[Option ID = 26751]

3.  $5/6$

[Option ID = 26752]

4. none of these

[Option ID = 26753]

Correct Answer :-

- $5/6$

[Option ID = 26752]

17) Are the three points whose position vectors are  $2\mathbf{i}+3\mathbf{j}-4\mathbf{k}$ ,  $\mathbf{i}-2\mathbf{j}+3\mathbf{k}$  and  $-7\mathbf{j}+10\mathbf{k}$  collinear?

[Question ID = 6690]

1. Yes

[Option ID = 26754]

2. No



4. None of these

[Option ID = 26757]

Correct Answer :-

- Yes

[Option ID = 26754]

18) The number of independent fundamental solutions in n-th order ordinary differential equation is

[Question ID = 6692]

1.  $n-1$  [Option ID = 26762]
2.  $n$  [Option ID = 26763]
3.  $n+1$  [Option ID = 26764]
4.  $2n$  [Option ID = 26765]

Correct Answer :-

- $n$  [Option ID = 26763]

19) If  $z_1 = 2 - 3i$  and  $z_2 = 4 + 6i$  then find  $\frac{z_1}{z_2}$

[Question ID = 6693]

1.  $-5/26-6i/13$   
[Option ID = 26766]
2.  $-5/26+6i/13$   
[Option ID = 26767]
3.  $8+18i$   
[Option ID = 26768]
4.  $8-18i$   
[Option ID = 26769]

Correct Answer :-

- $-5/26-6i/13$

[Option ID = 26766]

20) The rank of the following matrix  $\begin{pmatrix} 1 & 5 & 1 \\ 2 & 1 & 1 \\ 3 & 6 & 2 \end{pmatrix}$  is

[Question ID = 6699]

1. 1  
[Option ID = 26790]
2. 2  
[Option ID = 26791]
3. 3  
[Option ID = 26792]
4. 4  
[Option ID = 26793]

Correct Answer :-

- 2

[Option ID = 26791]

21) Two Carnot engines X and Y are operating in series. The engine X receives heat at 1200 K and rejects to a reservoir at a temperature T. The second engine Y receives the heat rejected by X and in turn rejects to a heat reservoir at 300 K. Calculate the temperature T (in Kelvin) for the situation when the efficiency of the engines is same

[Question ID = 6704]

1. 600 K [Option ID = 26810]
2. 750 K [Option ID = 26811]
3. 0 [Option ID = 26812]
4. 450 K [Option ID = 26813]

Correct Answer :-

- 600 K [Option ID = 26810]

22) A square conducting loop of mass m, side l and resistance R is dropped into a region with a uniform horizontal magnetic



whose direction is perpendicular to the plane of the falling loop. The loop will reach a terminal velocity  $v$  given by

FirstRanker's choice

www.FirstRanker.com

www.FirstRanker.com

[Question ID = 7330]

1.  $v = \frac{mgR}{(Bl)^2}$

[Option ID = 29314]

2.  $v = \frac{2mgR}{B^2 l^2}$

[Option ID = 29315]

3.  $v = \frac{mgR}{2B^2 l^2}$

[Option ID = 29316]

4. None of these

[Option ID = 29317]

Correct Answer :-

•  $v = \frac{mgR}{(Bl)^2}$

[Option ID = 29314]

23) An ideal inductor, a resistor of resistance  $R$  Ohms and a capacitor with adjustable capacitance are connected in series to an alternating voltage with an effective value of  $V$  Volts and with frequency of  $f$  Hz. The current flowing through the circuit when the capacitance of the capacitor is set to  $C_1$  is the same as when the capacitance of the capacitor is set to  $C_2$ ,  $C_2 > C_1$ . The inductance of the inductor  $L$  is given by

[Question ID = 7331]

1.  $\frac{1}{8\pi^2 f^2} \frac{C_1 + C_2}{C_1 C_2}$

[Option ID = 29318]

2.  $\frac{1}{8\pi^2 f^2} \frac{C_1 C_2}{C_1 + C_2}$

[Option ID = 29319]

3.  $\frac{1}{2\pi f} \frac{C_1 C_2}{C_1 - C_2}$

[Option ID = 29320]

4.  $\frac{1}{2\pi^2 f^2 R (C_1 - C_2)} \frac{C_1 C_2}{C_1 + C_2}$

[Option ID = 29321]

Correct Answer :-

•  $\frac{1}{8\pi^2 f^2} \frac{C_1 + C_2}{C_1 C_2}$

[Option ID = 29318]

24) A cylinder of length  $L$  is made up of an inner core of steel of radius  $r_1$  and an outer sheath of copper of thickness  $r_1$ . The resistivities of steel and copper are  $\rho_1$  and  $\rho_2$  respectively. The total resistance of the cylinder is

[Question ID = 7332]

1.  $\frac{(\rho_1 \rho_2) L}{\pi r_1^2 (3\rho_1 + \rho_2)}$

[Option ID = 29322]

2.  $\frac{(3\rho_1 + \rho_2) L}{\pi r_1^2}$

[Option ID = 29323]

3.  $\frac{(\rho_1 + \rho_2) L}{4\pi r_1^2}$

[Option ID = 29324]

4. Cannot be determined from the information provided above

www.FirstRanker.com



- $\frac{(\rho_1 \rho_2) L}{\pi r_1^2 (3\rho_1 + \rho_2)}$

[Option ID = 29322]

25) A meter stick is at an angle of  $45^\circ$  to the  $x'$  axis in its rest frame. The rod moves with a speed of  $\frac{1}{\sqrt{2}}c$  along the  $+x$  direction w.r.t. a frame  $S$ . The length of the rod in  $S$  is

[Question ID = 7333]

1.  $\frac{\sqrt{3}}{2}$  meters

[Option ID = 29326]

2.  $\frac{\sqrt{5}}{3}$  meters

[Option ID = 29327]

3.  $\frac{\sqrt{2}}{3}$  meters

[Option ID = 29328]

4.  $\frac{2}{3}$  meters

[Option ID = 29329]

Correct Answer :-

- $\frac{\sqrt{3}}{2}$  meters

[Option ID = 29326]

26) An AC generator with output and frequency  $f$  is connected to the plates of an air filled parallel plate capacitor of plate area  $A$  and plate separation  $d$ . The maximum value of the displacement current is

[Question ID = 7334]

1.  $\frac{2\pi \epsilon_0 fVA}{d}$

[Option ID = 29330]

2.  $\frac{\epsilon fV}{d}$

[Option ID = 29331]

3.  $\frac{2\pi f \epsilon_0 A}{Vd}$

[Option ID = 29332]

4. Cannot be determined from the information provided

[Option ID = 29333]

Correct Answer :-

- $\frac{2\pi \epsilon_0 fVA}{d}$

[Option ID = 29330]

27) An electron enters a uniform magnetic field of flux density  $1.2 \text{ Wb/m}^2$ . Find the energy difference in (eV), between electrons having spins parallel and anti-parallel to the field. (Given:  $\mu_B = 9.3 \times 10^{-24} \text{ J/T}$ )

[Question ID = 7335]

1.  $3.95 \times 10^{-2} \text{ eV}$

[Option ID = 29334]

2.  $13.95 \times 10^{-5} \text{ eV}$





4.  $33.95 \times 10^{-5} \text{ eV}$

[Option ID = 29337]

Correct Answer :-

- $13.95 \times 10^{-5} \text{ eV}$

[Option ID = 29335]

28) Using the vector atom model, determine the possible values of the angular momentum of an electron in  $f$ - shell

[Question ID = 7336]

1.  $\frac{3\sqrt{7}}{2} \hbar, \frac{\sqrt{35}}{2} \hbar$

[Option ID = 29338]

2.  $\frac{2\sqrt{7}}{2} \hbar, \frac{\sqrt{25}}{2} \hbar$

[Option ID = 29339]

3.  $\frac{5\sqrt{7}}{2} \hbar, \frac{\sqrt{15}}{2} \hbar$

[Option ID = 29340]

4.  $\frac{\sqrt{7}}{2} \hbar, \frac{\sqrt{5}}{2} \hbar$

[Option ID = 29341]

Correct Answer :-

- $\frac{3\sqrt{7}}{2} \hbar, \frac{\sqrt{35}}{2} \hbar$

[Option ID = 29338]

29) The two eigenvalues of the matrix  $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$  are

[Question ID = 7337]

1. 2,0

[Option ID = 29342]

2. 1,1

[Option ID = 29343]

3. 1,2

[Option ID = 29344]

4. 0,1

[Option ID = 29345]

Correct Answer :-

- 2,0

[Option ID = 29342]

30) The commutator,  $\pi[x^2, p_x]$ , is equal to

[Question ID = 7338]

1.  $ihx$

[Option ID = 29346]

2.  $2ihx$

[Option ID = 29347]

3.  $2ihp_x$

[Option ID = 29348]

4. Zero

[Option ID = 29349]

Correct Answer :-

- $ihx$

[Option ID = 29346]

31) A particle of mass  $m$  is confined in the ground state of a one dimensional box extending from  $x = -2L$  to  $x = +2L$ . The wave function of the particle in this state is



[Question ID = 7339]

1.  $\frac{h^2 \pi^2}{32mL^2}$

[Option ID = 29350]

2.  $\frac{h^2 \pi^2}{2mL^2}$

[Option ID = 29351]

3.  $\frac{h^2 \pi^2}{4mL^2}$

[Option ID = 29352]

4.  $\frac{h^2 \pi^2}{16mL^2}$

[Option ID = 29353]

Correct Answer :-

•  $\frac{h^2 \pi^2}{32mL^2}$

[Option ID = 29350]

32) The normalized wave functions  $\psi_1$  and  $\psi_2$ , correspond to the ground state and the first excited states of a particle in a potential. The operator  $\hat{A}$  acts on the wave functions as  $\hat{A}\psi_1 = \psi_2$  and  $\hat{A}\psi_2 = \psi_1$ . The expectation value of the operator  $\hat{A}$  for the state  $\psi = (3\psi_1 + 4\psi_2)/5$  is

[Question ID = 7340]

1. 0.96 [Option ID = 29354]

2. -0.32 [Option ID = 29355]

3. 0 [Option ID = 29356]

4. 0.75 [Option ID = 29357]

Correct Answer :-

• 0.96 [Option ID = 29354]

33) The primitive translation vector of a two-dimensional lattice are  $a = 2i + j$ ,  $b = 2j$ . The primitive translation vector of its reciprocal lattice in x-direction is given by

[Question ID = 7341]

1.  $a^* = \pi i$

[Option ID = 29358]

2.  $a^* = 2\pi i$

[Option ID = 29359]

3.  $a^* = i$

[Option ID = 29360]

4.  $a^* = \pi j$

[Option ID = 29361]

Correct Answer :-

•  $a^* = \pi i$

[Option ID = 29358]

34) The mean drift speed  $v_d$  of an electron in an applied electric field E with electron density 'n' can be expressed as

[Question ID = 7342]

1.  $v_d = |\sigma E / ne|$

[Option ID = 29362]

2.  $v_d = |\sigma E / e|$





4. none of these

[Option ID = 29365]

Correct Answer :-

•  $v_d = |\sigma E / ne|$

[Option ID = 29362]

35) An un-damped oscillator has time period  $\tau_0 = 1.0$  sec. Now a little damping is added so that its time period changes to  $\tau_1 = 1.001$  sec. By what factor will the amplitude of oscillation decrease after 10 cycles?

[Question ID = 7343]

1.  $\approx 17$

[Option ID = 29366]

2.  $\approx 1$

[Option ID = 29367]

3.  $\approx 1/17$

[Option ID = 29368]

4. None of these

[Option ID = 29369]

Correct Answer :-

•  $\approx 17$

[Option ID = 29366]

36) A kilogram of water has a constant heat capacity of 4.2 kJ/K/kg over the temperature range 0°C to 100°C. The water was initially at 0°C and is brought into contact with a heat reservoir at 100°C. When the water is in thermal equilibrium with the heat reservoir calculate the change in entropy of the universe (Water + Reservoir).

[Question ID = 7344]

1. 184.8 J/K

[Option ID = 29370]

2. 2437.8 J/K

[Option ID = 29371]

3. 0 J/K

[Option ID = 29372]

4. 1310.8 J/K

[Option ID = 29373]

Correct Answer :-

• 184.8 J/K

[Option ID = 29370]

37) Two identical finite bodies of constant volume and of constant heat capacity at constant volume  $C_v$ , are used to drive a heat engine. Their initial temperatures are  $T_1$  and  $T_2$ . The maximum amount of work which can be obtained from the system is

[Question ID = 7345]

1.  $C_v \{ 2 (T_1 T_2)^{1/2} - (T_1 + T_2) \}$

[Option ID = 29374]

2.  $C_v (T_1 + T_2)$

[Option ID = 29375]

3.  $C_v (T_1 T_2)^{1/2}$

[Option ID = 29376]

4. 0

[Option ID = 29377]



38) For a system of bosons, we can write the Bose-Einstein distribution function as  $f(E_i) = \frac{1}{\exp(\alpha + \beta E_i) - 1}$ , Where,  $\beta = 1/k_B T$  and  $\alpha = \mu/k_B T$  ( $k_B$  = Boltzmann constant). If  $\mu$  represents the chemical potential, then which one of the following is true?

[Question ID = 7346]

1.  $\mu \leq 0$

[Option ID = 29378]

2.  $\mu \geq 0$

[Option ID = 29379]

3.  $\mu \leq 1$

[Option ID = 29380]

4.  $\mu \geq 1$

[Option ID = 29381]

Correct Answer :-

•  $\mu \geq 0$

[Option ID = 29379]

39) An ideal capacitor C is charged to a voltage  $V_0$  and connected at  $t = 0$  across an ideal inductor L (The circuit now consists of a capacitor and an inductor only). If the resonant frequency

$$\omega_0 = 1/\sqrt{LC}$$

, the voltage across the capacitor at time  $t > 0$  is given by

[Question ID = 7347]

1.  $V_0$  [Option ID = 29382]

2.  $V_0 \cos(\omega_0 t)$  [Option ID = 29383]

3.  $V_0 \sin(\omega_0 t)$  [Option ID = 29384]

4.  $V_0 e^{-\omega_0 t} \cos(\omega_0 t)$

[Option ID = 29385]

Correct Answer :-

•  $V_0 \cos(\omega_0 t)$  [Option ID = 29383]

40) Magnetic moment of proton ( $\mu_p$ ) in terms of nuclear magneton ( $\mu_N$ ) is

[Question ID = 7348]

1.  $\mu_p = 1.9\mu_N$

[Option ID = 29386]

2.  $\mu_p = 2.7\mu_N$

[Option ID = 29387]

3.  $\mu_p = 3.8\mu_N$

[Option ID = 29388]

4.  $\mu_p = 5.4\mu_N$

[Option ID = 29389]

Correct Answer :-

•  $\mu_p = 2.7\mu_N$

[Option ID = 29387]

41) Find the eigenvalues of  $A + 4I$  where  $I$  is identity matrix and  $A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$

[Question ID = 7349]

1. 1,3

[Option ID = 29390]

2. 5,7



4. None of these

[Option ID = 29393]

Correct Answer :-

- 5,7

[Option ID = 29391]

42) The limit  $\lim_{n \rightarrow \infty} \left(\frac{1}{n}\right)^{\frac{1}{n}}$  is

[Question ID = 7350]

1.  $\frac{1}{e}$

[Option ID = 29394]

2. 1

[Option ID = 29395]

3. 0

[Option ID = 29396]

4.  $e$

[Option ID = 29397]

Correct Answer :-

- 1

[Option ID = 29395]

43)  $\left(\frac{1+i}{\sqrt{2}}\right)^{49}$

is equal to

[Question ID = 7351]

1.  $\left(\frac{1+i}{\sqrt{2}}\right)$

[Option ID = 29398]

2.  $\left(\frac{2+98i}{\sqrt{2^{49}}}\right)$

[Option ID = 29399]

3.  $\left(\frac{1-i}{\sqrt{2}}\right)$

[Option ID = 29400]

4.  $\left(\frac{2-98i}{\sqrt{2^{49}}}\right)$

[Option ID = 29401]

Correct Answer :-

- $\left(\frac{1+i}{\sqrt{2}}\right)$

[Option ID = 29398]

44)  $\sin\left(\frac{\pi}{4} + i\right)$  is equal to

[Question ID = 7352]

1.  $\frac{\sqrt{2}}{4} \left(e - \frac{1}{e}\right) + \frac{\sqrt{2}}{4} \left(e - \frac{1}{e}\right) i$

[Option ID = 29402]

2.  $\frac{\sqrt{2}}{4} \left(e + \frac{1}{e}\right) + \frac{\sqrt{2}}{4} \left(e - \frac{1}{e}\right) i$



[Option ID = 29404]

4.  $\frac{\sqrt{2}}{4} \left( e - \frac{1}{e} \right) + \frac{\sqrt{2}}{4} \left( e + \frac{1}{e} \right) i$

[Option ID = 29405]

Correct Answer :-

•  $\frac{\sqrt{2}}{4} \left( e + \frac{1}{e} \right) + \frac{\sqrt{2}}{4} \left( e - \frac{1}{e} \right) i$

[Option ID = 29403]

45) Two students are working on a math problem. The first student has probability  $\frac{1}{2}$  of solving it and the second student has probability  $\frac{3}{4}$  of solving it. What is the probability that at least one of them solves the problem?

[Question ID = 7353]

1.  $\frac{3}{8}$

[Option ID = 29406]

2.  $\frac{5}{8}$

[Option ID = 29407]

3.  $\frac{7}{8}$

[Option ID = 29408]

4.  $\frac{9}{8}$

[Option ID = 29409]

Correct Answer :-

•  $\frac{7}{8}$

[Option ID = 29408]

46) Expansion of the function  $f(z) = \frac{1}{z^2 - 3z + 2}$  in the region defined by  $|z| > 2$  is

[Question ID = 7354]

1.  $z^{-2} + 3z^{-4} + 7z^{-6} + \dots$

[Option ID = 29410]

2.  $z^{-2} + 3z^{-3} + 7z^{-4} + \dots$

[Option ID = 29411]

3.  $z^{-1} + 3z^{-2} + 7z^{-3} + \dots$

[Option ID = 29412]

4.  $z^{-3} + 3z^{-4} + 7z^{-5} + \dots$

[Option ID = 29413]

Correct Answer :-

•  $z^{-2} + 3z^{-3} + 7z^{-4} + \dots$

[Option ID = 29411]

47) The Fourier transformation of the function  $f(x) = 1$  for  $|x| < a$

$= 0$  for  $|x| > a$

is

[Question ID = 7355]

1.  $\sqrt{\frac{2}{\pi}} \frac{\sin sa}{s}$



2.  $\sqrt{\frac{2}{\pi}} \frac{\cos sa}{s}$

[Option ID = 29415]

3.  $\sqrt{\frac{\pi}{2}} \frac{\sin sa}{s}$

[Option ID = 29416]

4.  $\sqrt{\frac{\pi}{2}} \frac{\cos sa}{s}$

[Option ID = 29417]

Correct Answer :-

•  $\sqrt{\frac{2}{\pi}} \frac{\sin sa}{s}$

[Option ID = 29414]

48) The Laplace transformation of the function  $f(t) = 2^t$  is

[Question ID = 7356]

1.  $\frac{\ln 2}{s - \ln 2}$

[Option ID = 29418]

2.  $\frac{1}{s - \ln 2}$

[Option ID = 29419]

3.  $\frac{1}{s \ln 2 - 1}$

[Option ID = 29420]

4.  $\frac{\ln 2}{s \ln 2 - 1}$

[Option ID = 29421]

Correct Answer :-

•  $\frac{1}{s - \ln 2}$

[Option ID = 29419]

49) Consider a collection of non-interacting particles, each of mass  $m$  in a volume where the gravitational force is a -ve ( $z$ -direction). Consider the system is in thermal equilibrium at a temperature  $T$ . Find the partition function

[Question ID = 7357]

1.  $Q_N = \left[ \frac{(kT)^3}{2\pi m g h^2} \right]^{N/2}$

[Option ID = 29422]

2.  $Q_N = \left[ \frac{2\pi m g h^2}{(kT)^3} \right]^{N/2}$

[Option ID = 29423]

3.  $Q_N = \left[ \frac{(kT)^3}{2\pi m g h^2} \right]^N$

[Option ID = 29424]

4.  $Q_N = \left[ \frac{2\pi m g h^2}{(kT)^3} \right]^N$

[Option ID = 29425]



[Option ID = 29422]

50) The quantum distribution function for any gas atom which follows MB, BE and FD statistics is given as a generalized single form

$$f_i = g_i / [\exp(\epsilon_i - \mu) / (kT + j)]$$

If the distribution function follows the MB statistics in a classical limit then what will be the condition of the following. Symbols have their usual meanings

[Question ID = 7358]

1.  $\frac{f_i}{g_i} \ll 1 ; j = 1$

[Option ID = 29426]

2.  $\frac{f_i}{g_i} \ll 1 ; j = 0$

[Option ID = 29427]

3.  $\frac{g_i}{f_i} \gg 1 ; j = 1$

[Option ID = 29428]

4.  $\frac{g_i}{f_i} \gg 1 ; j = -1$

[Option ID = 29429]

Correct Answer :-

•  $\frac{f_i}{g_i} \ll 1 ; j = 0$

[Option ID = 29427]