Firstranker's choice in PsychologyRanker.com

www.FirstRanker.com

Topic:- PSY PHD S2

- Which of the following category is not used for the interpretation of Rorschach inkblot test?
 [Question ID = 9156]
- 1. Location [Option ID = 36618]
- 2. Content [Option ID = 36619]
- 3. Dominance [Option ID = 36620]
- Determinants [Option ID = 36621]

Correct Answer :-

- Dominance [Option ID = 36620]
- 2) In a 2x2x2 ANOVA, the number of factors and levels respectively are:-

[Question ID = 9157]

- 1. 3, 3 [Option ID = 36622]
- 2. 2, 2 [Option ID = 36623]
- 3. 3, 2 [Option ID = 36624]
- 2, 3 [Option ID = 36625]

Correct Answer :-

- 3, 2 [Option ID = 36624]
- A rank-order test of a difference among three or more dependent groups is called:-

[Question ID = 9158]

- 1. Spearman's rho [Option ID = 36626]
- 2. Kruskal Wallis test [Option ID = 36627]
- 3. Friedman test [Option ID = 36628]
- 4. Repeated Measures ANOVA [Option ID = 36629]

Correct Answer :-

- Friedman test [Option ID = 36628]
- Company records showing present performance and promotion possibility of inside candidates for the most important positions are called

[Question ID = 9159]

1. Qualification inventories

[Option ID = 36630]

2. Position replacement charts

[Option ID = 36631]

3. Personnel replacement charts

[Option ID = 36632]

4. Expectancy charts

[Option ID = 36633]

Correct Answer :-

· Personnel replacement charts

[Option ID = 36632]

5) In a 2 x 3 ANOVA, with N = 10 in each treatment group, the SS_{error} is calculated to be 378, the value of Mean SS_{error} will be:-

[Question ID = 9160]

1. 6

[Option ID = 36634]

2.

[Option ID = 36635]

3. 8

[Option ID = 36636]

[Option ID = 36637]

www.FirstRanker.com

Correct Answer :-

your other categorical independent variable(s). Question ID = 9161] Analysis of Variance (Option ID = 36438] Analysis of Variance (Option ID = 36439] Multivariate Analysis of Variance (Option ID = 36440] Factorial Analysis of Covariance (Option ID = 36441) orrect Answer : Analysis of Covariance (Option ID = 36441) orrect Answer : Analysis of Covariance (Option ID = 36441) orrect Answer : Analysis of Covariance (Option ID = 36441) orrect Answer : Analysis of Covariance (Option ID = 36442) (Case-Control Design (Option ID = 36442) Repeated Cintral Trial (Option ID = 36441) orrect Answer : Case-Control Design (Option ID = 36441) orrect Answer : Case-Control Design (Option ID = 36442) In Linear regression, the predicted values of dependent variable fall: Question ID = 9163] On the line of best fit (Option ID = 36448) On corresponding values of independent variables on abscissa (Option ID = 36447) On corresponding values of independent variables on ordinate (Option ID = 36448) orrect Answer : On the line of best fit (Option ID = 36446) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express on the intercept of value (Option ID = 36451) Approximate of determination (Option ID = 36651) Approximate of determination (Option ID = 36651) Orrect Answer : Regression coefficient (Option ID = 36651) Variance (Option ID = 36651) Variance (Option ID = 36653) Orrect Answer : Regression coefficient (Option ID = 36653) On the intercept of determination (Option ID = 36653) Variance (Option ID = 36653) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express or coefficient (Option ID = 36653) Orrect Answer : Regression coefficient (Option ID = 36653) Variance (Option ID = 36653) The amount of increase in Y that, on the average accompanies a unit increase in X (when both measures are express or coefficient (Option ID = 36653) Orrect Answer : Regression coefficient (Option ID = 36653) Orrect Answ	i) In	www.FirstRanker.com www.FirstRanker.com the effect of an undesired continuous variable on a dependent variable is partialled out
Question ID = 9161] Analysis of Covariance (Option ID = 36638] Analysis of Covariance (Option ID = 36639] Michitariata Analysis of Variance (Option ID = 36649] Factorial Analysis of Covariance (Option ID = 36649] Factorial Analysis of Covariance (Option ID = 36649] Analysis of Covariance (Option ID = 36649] A retrospective study, called as	rom other categori	
Analysis of Corariance (Option ID = 36649) Michitariate Analysis of Variance (Option ID = 36640) Factorial Analysis of Corariance (Option ID = 36641) 7. A retrospective study, called as, would select people who had, or had not, experienced a heart attained then look backward in time to see whether they had been in the habit of taking aspirin in the past. Question ID = 9162] Case-Control Design (Option ID = 36642) Randomed Clinical Trial (Option ID = 36643) Carbort Study (Dittion ID = 36644) Repeated clinical Trial (Option ID = 36643) Control Study (Dittion ID = 36644) Repeated clinical Trial (Option ID = 36642) In Linear regression, the predicted values of dependent variable fallity on the line of best fit (Option ID = 36643) On the line of best fit (Option ID = 36644) On corresponding values of independent variables on abscissa (Option ID = 36647) On corresponding values of independent variables on ordinate (Option ID = 36648) On the line of best fit (Option ID = 36646) On the line of best fit (Option ID = 36646) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express or raw-score units) is called:- Question ID = 91641 Regression coefficient (Option ID = 36650) Regression coefficient (Option ID = 36650) Forect Answer :- Regression coefficient (Option ID = 36650) On Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeling with an emen, are used to determine:- Question ID = 91651 Item validity (Option ID = 36653) Multitat multimethed analysis (Option ID = 36660) Orceres validity (Option ID = 36660) Multitat multimethed analysis (Option ID = 36660) Orceres validation (Option ID = 36660) Orceres validation (Option ID = 36660) Orceres validation (Option ID = 36660)	_	
Multivariate Analysis of Variance (Option ID - 36640] Factorial Analysis of Covariance (Option ID - 36641) Pretect Answer : Analysis of Covariance (Option ID - 36639] A retrospective study, called as, would select people who had, or had not, experienced a heart atta nd then look backward in time to see whether they had been in the habit of taking aspirin in the past. Question ID - 9162] Case-Control Design (Option ID - 36642] Randomierd Cinical Trial (Option ID - 36643) Cohort Study (Option ID - 36644) Repeated Clinical Trial (Option ID - 36645) arrect Answer : Case-Control Design (Option ID - 36642) In Linear regression, the predicted values of dependent variable fall:-Question ID - 91631 On the line of beat fit (Option ID - 36649) On corresponding values of independent variables on abscissa (Option ID - 36648) On the intercept of Y- axis (Option ID - 36649) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express 1 raw-score units) is called:-Question ID - 91641 Regression coefficient (Option ID - 36650) Coefficient of determination (Option ID - 36650) Coefficient of determination (Option ID - 36650) Prairance (Option ID - 36650) On In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an term, are used to determine:- Question ID - 91651 Item validity (Option ID - 36650) Item validity (Option ID - 36650) Item validity (Option ID - 36653) Item validity (Option ID - 36653) Item validity (Option ID - 36653) Item watchilly (Option ID - 36653) Multitrat multimethed analysis (Option ID - 36660) Concurrent validity (Option ID - 36650) Multitrat multimethed analysis (Option ID - 36660) Concurrent validity (Option ID - 36660)		
Factorial Analysis of Covariance [Option ID = 36641] Princed Answer : Analysis of Covariance [Option ID = 36639] A retrospective study, called as, would select people who had, or had not, experienced a heart attand then look backward in time to see whether they had been in the habit of taking aspirin in the past. Question ID = 9162] Case-Control Besign [Option ID = 36642] Randomized Clinical Trial [Option ID = 36643] Cohort Study [Option ID = 36644] Repeated Clinical Trial [Option ID = 36645] Ornered Answer : Case-Control Design [Option ID = 36642] On Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On the line of best fit [Option ID = 36646] On corresponding values of independent variables on ordinate [Option ID = 36647] On corresponding values of independent variables on ordinate [Option ID = 36648] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36646] On the interest of ** axis [Option ID = 36650] Conflictent of determination [Option ID = 36650] National or or of estimate [Option ID = 36650] National or or of estimate [Option ID = 36650] On In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an ema, are used to determine: Question ID = 9165] Item walldity (Option ID = 36650] Item validity (Option ID = 36650] Item validity (Option ID = 36650] Number of the following is a method for validating a construct? Question ID = 9166] Concurrent validity (Option ID = 36650] Multirat mutimethod analysis (Option ID = 36660) Correct Answer :- Correct Answer :-		
Analysis of Covariance (Option ID = 36639) A retrospective study, called as, would select people who had, or had not, experienced a heart atta of then look backward in time to see whether they had been in the habit of taking aspirin in the past. Question ID = 9162] Case-Cantrol Design (Option ID = 36642) Randomized Clinical Trial (Option ID = 36643) Cabort Study (Option ID = 36644) Repeated Clinical Trial (Option ID = 36645) Carrect Answer: Case-Cantrol Design (Option ID = 36645) In Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On the line of best fit (Option ID = 36646) On corresponding values of independent variables on abscissa (Option ID = 36647) On the line of best fit (Option ID = 36646) On the line of best fit (Option ID = 36646) On the line of best fit (Option ID = 36646) On the line of best fit (Option ID = 36646) Or the line of best fit (Option ID = 36650) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express on raw-score units) is called:- Question ID = 9164] Regression coefficient (Option ID = 36650) Coefficient of determination (Option ID = 36650) Coefficient of determination (Option ID = 36650) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express on raw-score units) is called:- Question ID = 9164] Regression coefficient (Option ID = 36650) Option ID = 36650 Option ID = 36650 Option ID = 36650 Coefficient of determination (Option ID = 36650) Item validity (Option ID = 36650) Option ID = 36660) Concurrent validity (Option ID = 36660) Concurrent validity (Opti		
Augustion ID = 9162] Case-Control Design [Option ID = 36642] Case-Control Design [Option ID = 36642] Repeated Clinical Trial [Option ID = 36643] Cohort Study [Option ID = 36645] Provent Answer: Case-Control Design [Option ID = 36645] On the line of best fit [Option ID = 36642] On the line of best fit [Option ID = 36642] On corresponding values of independent variables on abscissa [Option ID = 36647] On corresponding values of independent variables on adscissa [Option ID = 36648] On the line of best fit [Option ID = 36646] On the line roy of Y: axis [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36650] Coefficient of determination [Option ID = 36650] Coefficient of determination [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36650] Or the line of best fit [Option ID = 36660] Or the l	orrect Answer :- Analysis of Covarianc	e [Option ID = 36639]
Augustion ID = 9162] Case-Control Design [Option ID = 36642] Case-Control Design [Option ID = 36642] Repeated Clinical Trial [Option ID = 36643] Cohort Study [Option ID = 36645] Provent Answer: Case-Control Design [Option ID = 36645] On the line of best fit [Option ID = 36642] On the line of best fit [Option ID = 36646] On corresponding values of independent variables on abscissa [Option ID = 36647] On corresponding values of independent variables on adscissa [Option ID = 36647] On corresponding values of independent variables on adscissa [Option ID = 36648] On the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or the line of best fit [Option ID = 36646] Or The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express in raw-score units) is called: Question ID = 9164] Regression coefficient [Option ID = 36650] Coefficient of determination [Option ID = 36652] Variance [Option ID = 36653] Item validity [Option ID = 36650] Lem validity [Option ID = 36655] Lem validity [Option ID = 36656] Lem validity [Option ID = 3665	\ A retrespective	study, called as would select people who had, or had not, experienced a heart atta
Question ID = 9162] Case-Control Design [Option ID - 36642] Randomized Clisical Trial [Option ID - 36643] Cohort Study [Option ID - 36644] Repeated Clisical Trial [Option ID - 36645] orrect Answer :- Case-Control Design [Option ID - 36642]) In Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On corresponding values of independent variables on abscissa [Option ID - 36647] On corresponding values of independent variables on ordinate [Option ID - 36647] On corresponding values of independent variables on ordinate [Option ID - 36648] On the line of best fit [Option ID - 36646] On the line of best fit [Option ID - 36646] On the line of best fit [Option ID - 36646] On the line of best fit [Option ID - 36646] The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express or raw-score units) is called:- Question ID = 9164] Regression coefficient [Option ID - 36650] Coefficient of determination [Option ID - 36650] Orrect Answer :- Regression coefficient [Option ID - 36650] Or Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an em, are used to determine:- Question ID = 9165] Item validity [Option ID - 36655] Item valiability [Option ID - 36655] Ornect Validity [Option ID - 36655] Multitarit multimethod analysis [Option ID - 36660] Concurrent validity [Option ID - 36658] Concurrent validity [Option ID - 36665] Increase validation [Option ID - 36660] Concurrent validity [Option ID - 36666]		
Case-Control Design [Option ID = 36643] Repeated Clinical Trial [Option ID = 36645] Cohort Study (Option ID = 36645] Repeated Clinical Trial [Option ID = 36645] Repeated Clinical Trial [Option ID = 36645] In Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On the line of best fit [Option ID = 36646] On corresponding values of independent variables on abscissa [Option ID = 36647] On corresponding values of independent variables on arbitrate [Option ID = 36648] On the intercept of Y- axis [Option ID = 36646] On the intercept of Y- axis [Option ID = 36646] The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express in raw-score units) is called:- Question ID = 9164] Regression coefficient [Option ID = 36650] Coefficient of determination [Option ID = 36650] Variance (Option ID = 36653) Direct Answer:- Regression coefficient [Option ID = 36650] Oil In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an item, are used to determine:- Question ID = 9165] Item validity (Option ID = 3665) Item validity (Option ID = 36657) Item validity (Option ID = 36655) Item validity (Option ID = 36658) Concurrent validity (Option ID = 36658)		
Cohort Study (Option ID = 36644) Repeated Clinical Trial (Option ID = 36645) In Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On the line of best fit (Option ID = 36642) On corresponding values of independent variables on abscissa (Option ID = 36647) On corresponding values of independent variables on ordinate (Option ID = 36648) On the intercet of Y- axis (Option ID = 36649) Orrect Answer: On the line of best fit (Option ID = 36646) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express raw-score units) is called:- Question ID = 9164] Regression coefficient (Option ID = 36650) Coefficient of determination (Option ID = 36651) Standard error of estimate (Option ID = 36652) Variance (Option ID = 36653) Orrect Answer: Regression coefficient (Option ID = 36650) O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an earn, are used to determine:- Question ID = 9165] Item validity (Option ID = 36656) Item validity (Option ID = 36655) Item reliability (Option ID = 36655) Item reliability (Option ID = 36655) Item reliability (Option ID = 36655) Item difficutly (Option ID = 36655) Item difficutly (Option ID = 36655) Item validity (Option ID = 36655) Item validity (Option ID = 36655) Multitrait multimethed analysis (Option ID = 36660) Concurrent validity (Option ID = 36651) Orrect Answer:- Item difficut (Option ID = 36658) Content validity (Option ID = 36659) Multitrait multimethed analysis (Option ID = 36660) Correct Answer:-	-	
Represted Clinical Trial [Option ID = 36645] orrect Answer: Case-Control Design [Option ID = 36642]) In Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On corresponding values of independent variables on abscissa [Option ID = 36647] On corresponding values of independent variables on ordinate [Option ID = 36648] On the intercept of Y- axis [Option ID = 36649] On the intercept of Y- axis [Option ID = 36649] On the line of best fit [Option ID = 36649] On the line of best fit [Option ID = 36648] On the line of best fit [Option ID = 36648] On the line of best fit [Option ID = 36648] On the line of best fit [Option ID = 36648] On the line of best fit [Option ID = 36650] Coefficient of determination [Option ID = 36650] Coefficient of determination [Option ID = 36651] Standard error of estimate [Option ID = 36652] Variance [Option ID = 36653] On In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etc., are used to determine: Question ID = 9165] Item validity [Option ID = 36650] Item validity [Option ID = 36651] Item validity [Option ID = 36651] orrect Answer: Item difficulty (Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 91661 Concurrent validity [Option ID = 36658] Content validity [Option ID = 36661] orrect Answer:		
Case-Control Design (Option ID = 36642)) In Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On the line of best fit (Option ID = 36646) On corresponding values of independent variables on abscissa (Option ID = 36647) On corresponding values of independent variables on ordinate (Option ID = 36648) On the intercept of Y- axis (Option ID = 36649) or rect Answer:- On the line of best fit (Option ID = 36646)) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express or raw-score units) is called:- Question ID = 9164] Regression coefficient (Option ID = 36650) Coefficient of determination (Option ID = 36650) Coefficient of determination (Option ID = 36651) Standard error of estimate (Option ID = 36651) Standard error of estimate (Option ID = 36651) orrect Answer:- Regression coefficient (Option ID = 36650) O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an actem, are used to determine:- Question ID = 9165] Item validity (Option ID = 36657) orrect Answer:- Item difficulty (Option ID = 36655) Item variability (Option ID = 36655) In Which of the following is a method for validating a construct? Question ID = 91661 Concurrent validity (Option ID = 36658) Content validity (Option ID = 36658) Content validity (Option ID = 36656) Concurrent validity (Option ID = 36656)		
in Linear regression, the predicted values of dependent variable fall:- Question ID = 9163] On the line of best fit (Option ID = 36646) On corresponding values of independent variables on abscissa (Option ID = 36647) On the corresponding values of independent variables on ordinate (Option ID = 36648) On the intercept of Y- axis (Option ID = 36649) Orrect Answer :- Question ID = 9164] Regression coefficient (Option ID = 36650) Correct Answer :- Regression coefficient (Option ID = 36650) Or in Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an atem, are used to determination (Dotton ID = 36650) Item validity (Option ID = 36657) Item validity (Option ID = 36657) Orrect Answer :- Regression to efficient (Option ID = 36650) Item validity (Option ID = 36657) Orrect Answer :- Item difficulty (Option ID = 36657) Orrect Answer :- Item difficulty (Option ID = 36655) Item variability (Option ID = 36657) Orrect Answer :- Item difficulty (Option ID = 36658) Orrect Answer :- Item difficulty (Option ID = 36658) Multitrait multity (Option ID = 36658) Ontent validity (Option ID = 36658) Outent validity (Option ID = 36658) Outent validity (Option ID = 36658) Multitrait multitraitent datalysis (Option ID = 36660) Cross-validation (Option ID = 36651) Orrect Answer :-	orrect Answer :-	
Question ID = 9163] On the line of best fit (Option ID = 36646) On corresponding values of independent variables on abscissa [Option ID = 36647] On corresponding values of independent variables on ordinate [Option ID = 36648] On the intercept of Y- axis [Option ID = 36649] on the intercept of Y- axis [Option ID = 36646] On the line of best fit [Option ID = 36646]) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express in raw-score units) is called:- Question ID = 9164] Regression coefficient [Option ID = 36650] Coefficient of determination [Option ID = 36652] Variance [Option ID = 36653] orrect Answer:- Regression coefficient [Option ID = 36650] O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an item, are used to determine:- Question ID = 9165] Item validity (Option ID = 36654) Item validity (Option ID = 36654) Item validity (Option ID = 36655) Item reliability (Option ID = 36656) Item variability (Option ID = 36655) Item validity (Option ID = 36658) Concurrent validity (Option ID = 36658) Concurrent validity (Option ID = 36658) Concurrent validity (Option ID = 36658) Content validity (Option ID = 36658) Content validity (Option ID = 36658) Content validity (Option ID = 36656) Cross-validation (Option ID = 36661) Orrect Answer:-	Case-Control Design	[Option ID = 36642]
On the line of best fit (Option ID = 36646) On corresponding values of independent variables on abscissa (Option ID = 36647) On the line of best fit (Option ID = 36649) Direct Answer :- On the line of best fit (Option ID = 36646) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express in raw-score units) is called:- Question ID = 9164] Regression coefficient (Option ID = 36650) Coefficient of determination (Option ID = 36650) Coefficient of determination (Option ID = 36651) Standard error of estimate (Option ID = 36652) Variance (Option ID = 36653) Direct Answer :- Regression coefficient (Option ID = 36650) O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an em, are used to determine:- Question ID = 9165) Item validity (Option ID = 36654) Item validity (Option ID = 36656) Item variability (Option ID = 36657) Direct Answer :- Item difficulty (Option ID = 36658) Content validity (Option ID = 36656) Multitrait multimented analysis (Option ID = 36660) Cross-validation (Option ID = 36661) Direct Answer :-		
On corresponding values of independent variables on abscissa (Option ID - 36647) On corresponding values of independent variables on ordinate (Option ID - 36648) On the intercept of Y- axis (Option ID - 36649) orrect Answer :- On the line of best fit (Option ID - 36646)) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express in raw-score units) is called:- Question ID = 9164] Regression coefficient (Option ID - 36650) Coefficient of determination (Option ID - 36651) Standard error of estimate (Option ID - 36652) Variance (Option ID - 36653) orrect Answer :- Regression coefficient (Option ID - 36650) O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an earm, are used to determine:- Question ID = 9165] Item validity (Option ID - 36654) Item validity (Option ID - 36655) Item reliability (Option ID - 36657) Item reliability (Option ID - 36655) Item reliability (Option ID - 36655) Item reliability (Option ID - 36655) Item variability (Option ID - 36655) Item variability (Option ID - 36658) Concurrent validity (Option ID - 36656) Corsen-validation (Option ID - 36660) Cross-validation (Option ID - 36661)		
On corresponding values of independent variables on ordinate [Option ID = 36648] On the intercept of Y- axis [Option ID = 36649] orrect Answer: Regression coefficient (Option ID = 36650) Or In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an item, are used to determine:- Question ID = 9164] Regression coefficient (Option ID = 36650) Or In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an item, are used to determine:- Question ID = 9165] Item validity (Option ID = 36655) Item reliability (Option ID = 36655) Item validity (Option ID = 36655) Item validity (Option ID = 36655) In Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity (Option ID = 36658) Content validity (Option ID = 36658) Content validity (Option ID = 36656) Authority (Option ID = 36651) Orrect Answer:- Item difficulty (Option ID = 36658) Content validity (Option ID = 36658) Content validity (Option ID = 36658) Content validity (Option ID = 36656) Orrect Answer:-		
On the intercept of Y- axis [Option ID = 36649] orrect Answer :- On the line of best fit [Option ID = 36646]) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express in raw-score units) is called:- Question ID = 9164] Regression coefficient [Option ID = 36650] Coefficient of determination [Option ID = 36651] Standard error of estimate [Option ID = 36652] Variance [Option ID = 36653] orrect Answer :- Regression coefficient [Option ID = 36650] O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an even, are used to determine:- Question ID = 9165] Item validity [Option ID = 36654] Item validity [Option ID = 36656] Item validity [Option ID = 36657] orrect Answer :- Item difficulty [Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-		
On the line of best fit [Option ID = 36646]) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express in raw-score units) is called:- Question ID = 9164] Regression coefficient [Option ID = 36650] Coefficient of determination [Option ID = 36651] Standard error of estimate [Option ID = 36652] Variance [Option ID = 36653] Direct Answer:- Regression coefficient [Option ID = 36650] O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an item, are used to determine:- Question ID = 9165] Item validity [Option ID = 36654] Item validity [Option ID = 36656] Item variability [Option ID = 36656] Item variability [Option ID = 36657] Direct Answer:- Item difficulty [Option ID = 36658] Concurrent validity [Option ID = 36658] Concurrent validity [Option ID = 36658] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] Direct Answer:-		
) The amount of increase in Y that, on the average, accompanies a unit increase in X (when both measures are express nav-score units) is called:— Question ID = 9164] Regression coefficient [Option ID - 36650] Coefficient of determination [Option ID - 36651] Standard error of estimate [Option ID - 36652] Variance [Option ID - 36653] Direct Answer:- Regression coefficient [Option ID - 36650] O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an em, are used to determine:- Question ID = 9165] Item validity [Option ID - 36654] Item reliability [Option ID - 36656] Item variability [Option ID - 36655] Item variability [Option ID - 36657] Direct Answer:- Item difficulty [Option ID - 36658] Concurrent validity [Option ID - 36658] Content validity [Option ID - 36658] Content validity [Option ID - 36658] Multitrait multimethod analysis [Option ID - 36660] Cross-validation [Option ID - 36661] Direct Answer:-		
n raw-score units) is called:- Question ID = 9164] Regression coefficient (Option ID = 36650] Coefficient of determination (Option ID = 36651) Standard error of estimate (Option ID = 36652) Variance (Option ID = 36653) orrect Answer :- Regression coefficient (Option ID = 36650) O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an term, are used to determine:- Question ID = 9165] Item validity (Option ID = 36654) Item difficulty (Option ID = 36656) Item reliability (Option ID = 36657) orrect Answer :- Item difficulty (Option ID = 36658) Concurrent validity (Option ID = 36658) Content validity (Option ID = 36658) Multitrait multimethod analysis (Option ID = 36660) Cross-validation (Option ID = 36661) orrect Answer :-	On the line of best fit	t (Option ID = 36646)
Regression coefficient [Option ID = 36650] O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an item, are used to determine;— Question ID = 9165] Item validity [Option ID = 36654] Item difficulty [Option ID = 36655] Item reliability [Option ID = 36657] orrect Answer:— Item difficulty [Option ID = 36657] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36658] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer:-		
Regression coefficient [Option ID = 36650] O) In Classical Test Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an item, are used to determine;— Question ID = 9165] Item validity [Option ID = 36654] Item difficulty [Option ID = 36655] Item reliability [Option ID = 36657] orrect Answer:— Item difficulty [Option ID = 36657] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36658] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer:-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est	it [Option ID = 36650] nination [Option ID = 36651] imate [Option ID = 36652]
tem, are used to determine:- Question ID = 9165] Item validity [Option ID = 36654] Item difficulty [Option ID = 36655] Item reliability [Option ID = 36656] Item variability [Option ID = 36657] orrect Answer:- Item difficulty [Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer:-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est Variance [Option ID =	it [Option ID = 36650] nination [Option ID = 36651] imate [Option ID = 36652]
tem, are used to determine:- Question ID = 9165] Item validity [Option ID = 36654] Item difficulty [Option ID = 36655] Item reliability [Option ID = 36656] Item variability [Option ID = 36657] orrect Answer:- Item difficulty [Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer:-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est: Variance [Option ID =	it [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] · 36653]
Question ID = 9165] Item validity [Option ID = 36654] Item difficulty [Option ID = 36655] Item reliability [Option ID = 36656] Item variability [Option ID = 36657] orrect Answer :- Item difficulty [Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien	it [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] id (Option ID = 36650)
Item difficulty [Option ID = 36655] Item reliability [Option ID = 36656] Item variability [Option ID = 36657] orrect Answer:- Item difficulty [Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer:-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est Variance [Option ID = Forrect Answer :- Regression coefficien O) In Classical Tes	t [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] it [Option ID = 36650] it [Option ID = 36650] It Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an
Item reliability [Option ID = 36656] Item variability [Option ID = 36657] orrect Answer :- Item difficulty [Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien O) In Classical Testern, are used to de	it [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] it [Option ID = 36650] it [Option ID = 36650] it Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine:-
Item variability [Option ID = 36657] orrect Answer :- Item difficulty [Option ID = 36655] 1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est: Variance [Option ID = Orrect Answer :- Regression coefficien O) In Classical Test term, are used to de Question ID = 9165 Item validity [Option	it [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] imate [Option ID = 36652] it [Option ID = 36650] It Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine:- i] ID = 36654]
1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficient Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien O) In Classical Testem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option	it [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] is 36653] it [Option ID = 36650] it Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine:- is ID = 36654] in ID = 36655]
1) Which of the following is a method for validating a construct? Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer:-	Question ID = 9164 Regression coefficient Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien O) In Classical Test tem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item reliability [Option	it [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] is 36653] It [Option ID = 36650] It Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine:- is ID = 36654] in ID = 36655] in ID = 36656]
Question ID = 9166] Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficient Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien O) In Classical Testem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item variability [Option	it [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] is 36653] It [Option ID = 36650] It Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine:- is ID = 36654] in ID = 36655] in ID = 36656]
Concurrent validity [Option ID = 36658] Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficient Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien O) In Classical Test tem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item variability	it [Option ID = 36650] Initiation [Option ID = 36651] Imate [Option ID = 36652] Initiation [Option ID = 36652] Initiation [Option ID = 36650] Initiation [Option ID = 36654] Initiation [Option ID = 36656] Initiation [Option ID = 36656] Initiation [Option ID = 36657] Initiation [Option ID = 36657] Initiation [Option ID = 36650] Initiation [Option ID = 36657] Initiation [Option ID = 36657] Initiation [Option ID = 36650] Initiation [Option ID = 36657] Initiation [Option ID = 36650] Initiation [Option ID = 36650] Initiation [Option ID = 36657] Initiation [Option ID = 36650] Initiation [O
Content validity [Option ID = 36659] Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est Variance [Option ID = Orrect Answer :- Regression coefficien O) In Classical Test tem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item variability [Option Item variability [Option Item difficulty [Option	int (Option ID = 36650) inination (Option ID = 36651) imate (Option ID = 36652) init (Option ID = 36652) init (Option ID = 36650) It (Option ID = 36650) It Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine:- i] ID = 36654] ID = 36655] ID = 36655] ID = 36657] In ID = 36655] In ID = 36655] In ID = 36655]
Multitrait multimethod analysis [Option ID = 36660] Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficient Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficient O) In Classical Test term, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item variability [Option Item difficulty [Option Item difficu	int (Option ID = 36650) inination (Option ID = 36651) imate (Option ID = 36652) int (Option ID = 36650) int (Option ID = 36650
Cross-validation [Option ID = 36661] orrect Answer :-	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est Variance [Option ID = Forrect Answer: Regression coefficien O) In Classical Test tem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item variability [Option Item difficulty [Option [Item difficulty [Item di	at [Option ID = 36650] mination [Option ID = 36651] mate [Option ID = 36652] at [Option ID = 36650] at [Option ID = 36650] at [Option ID = 36650] at Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine:- at [ID = 36654] at ID = 36655] at ID = 36656] at ID = 36655]
	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien O) In Classical Testerm, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item variability [Option Item difficulty [Option [Item difficulty [Item d	It (Option ID = 36650) Initiation (Option ID = 36651) Imate (Option ID = 36652) Initiate (Option ID = 36652) Initiate (Option ID = 36650) It Theory, p values, defined as the proportion of persons correctly responding to or agreeing with an etermine: Initiation ID = 36654) Initiation ID = 36655] Initiation ID = 36658]
Multitrait multimethod analysis [Option ID = 36660]	Question ID = 9164 Regression coefficient Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficient O) In Classical Test tem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item difficulty [Option Item variability [Option Item difficulty [Option Item difficul	int [Option ID = 36650] inination [Option ID = 36651] imate [Option ID = 36652] int [Option ID = 36650] int [Option ID = 36654] int [Option ID = 36655] int [Option ID = 36655] int [Option ID = 36655] int [Option ID = 36656] int [Option ID = 36658] int [Option ID = 36658] int [Option ID = 36659] int [Option ID = 36660]
	Question ID = 9164 Regression coefficien Coefficient of determ Standard error of est: Variance [Option ID = orrect Answer :- Regression coefficien O) In Classical Testem, are used to de Question ID = 9165 Item validity [Option Item difficulty [Option Item reliability [Option Item variability [Option Item difficulty [Option I) Which of the for Question ID = 9166 Concurrent validity [Option Multitrait multimetho Cross-validation [Option In Correct Answer :- Item difficulty [Option In Concurrent validity [Optio	It [Option ID = 36650] Intraction [Option ID = 36655] Intraction [Option ID = 36658] Intraction [Option ID = 36658] Intraction [Option ID = 36660] Intracti

Contracting [Option ID = 36663]
 Offshoring [Option ID = 36664]
 Recruiting [Option ID = 36665]



www.FirstRanker.com

13) An employee refused to obey a supervisors' order to meet with him unless a union representative was present at the meeting. As a result, the employee was discharged and subsequently filed a grievance protesting the discharge. This is a case of:-

[Question ID = 9168]

- 1. Insubordination [Option ID = 36666]
- 2. Lockout [Option ID = 36667]
- 3. Absenteeism [Option ID = 36668]
- 4. Injunction [Option ID = 36669]

Correct Answer :-

- Insubordination [Option ID = 36666]
- 14) In a ______ scale, stimuli (test items) are presented in order of increasing extremeness.

[Question ID = 9169]

- Thurstone [Option ID = 36670]
- Likert [Option ID = 36671]
- 3. Osgood [Option ID = 36672]
- Guttman [Option ID = 36673]

Correct Answer :-

- Guttman [Option ID = 36673]
- ______ calculates rater agreement in a fairly narrow circumstance. Specifically, two judges place a number of stimuli into nominal categories.

[Question ID = 9170]

- 1. Kendall's coefficient of concordance [Option ID = 36674]
- 2. Cohen's Kappa coefficient [Option ID = 36675]
- 3. Inter-observer coefficient [Option ID = 36676]
- 4. Spearman's rater agreement [Option ID = 36677]

Correct Answer :-

- Cohen's Kappa coefficient [Option ID = 36675]
- is the correlation between the extracted factor and variable.

[Question ID = 9171]

- 1. Communality [Option ID = 36678]
- 2. Eigen value [Option ID = 36679]
- 3. Factor loading [Option ID = 36680]
- 4. Specificity [Option ID = 36681]

Correct Answer :-

- Factor loading [Option ID = 36680]
- 17) Which of the following cannot be used to describe the spread of values in a distribution of data?

[Question ID = 9172]

- The range [Option ID = 36682]
- 2. Measures of variability [Option ID = 36683]
- 3. The standard deviation [Option ID = 36684]
- 4. The mean [Option ID = 36685]

Correct Answer :-

- The mean [Option ID = 36685]
- 18) In an experimental study, when the inferential statistics indicate that the experimental group's score differed significantly from the control group's score, it can be concluded that;-

[Question ID = 9173]

- 1. The difference between the groups can be attributed to chance [Option ID = 36686]
- 2. The difference between the groups can be attributed to the treatment [Option ID = 36687]
- 3. There are no differences between the groups [Option ID = 36688]
- 4. The difference between the groups arises from the individual difference among the participants. [Option ID = 36689]

Correct Answer :-

- The difference between the groups can be attributed to the treatment [Option ID = 36687]
- 19) The correlation between a natural dichotomous variable and continuous variable is called;-

[Question ID = 9174]

- Biserial correlation [Option ID = 36690]
- Point biserial correlation [Option ID = 36691]
- Tetrachoric correlation [Option ID = 36692]
 Phi- coefficient [Option ID = 36693]
- www.FirstRanker.com



20) Which statement is not true about Chomskywww.FirstRanker.com.ment?

www.FirstRanker.com

[Question ID = 9175]

1. There is a universal grammar.

[Option ID = 36694]

2. Language acquisition is genetically wired.

[Option ID = 36695]

3. There is an innate modular language.

[Option ID = 36696]

4. Language acquisition is based on reinforced learning

[Option ID = 36697]

Correct Answer :-

· Language acquisition is based on reinforced learning

[Option ID = 36697]

21) Which of the validity scales of MMPI-2 can be used to correct people's tendency to under report as they fear being judged or deny having any problems?

[Question ID = 9176]

- L-scale [Option ID = 36698]
- 2. TRIN scale [Option ID = 36699]
- 3. VRIN scale [Option ID = 36700]
- 4. K-scale [Option ID = 36701]

Correct Answer :-

K-scale [Option ID = 36701]

22) The smallest unit of speech that convey meaning is called:

[Question ID = 9177]

- Phonemes [Option ID = 36702]
- Lexemes [Option ID = 36703]
- Syntax [Option ID = 36704]
- 4. Morphemes [Option ID = 36705]

Correct Answer :-

Morphemes [Option ID = 36705]

23) Thomas Kuhn's major contribution was in delineating the role of:

[Question ID = 9178]

- 1. Values and consensus in science [Option ID = 36706]
- 2. Errors in science [Option ID = 36707]
- 3. Gender in science [Option ID = 36708]
- Money in science [Option ID = 36709]

Correct Answer :-

· Values and consensus in science [Option ID = 36706]

24) Paul Feyerabend speaks in favour of:-

[Question ID = 9179]

- A pluralistic vision of science [Option ID = 36710]
- 2. A universal vision of science [Option ID = 36711]
- 3. Western science [Option ID = 36712]
- Modern science [Option ID = 36713]

Correct Answer :-

A pluralistic vision of science [Option ID = 36710]

25) Feminist psychology is mainly concerned with:-

[Question ID = 9180]

- 1. Critiquing mainstream psychology [Option ID = 36714]
- 2. Creating employment for women psychologists [Option ID = 36715]
- 3. Attacking patriarchy [Option ID = 36716]
- 4. Lending women a voice and the construction of a female psychology [Option ID = 36717]

Correct Answer :-

Lending women a voice and the construction of a female psychology [Option ID = 36717]

www.FirstRanker.com

26) In qualitative research, one of the suggested approach to establish validity of research process is:-

[Option ID = 36718] 2. Homocidacity	www.FirstRanker.com	www.FirstRanker.com
[Option ID = 36719] 3. Triangulation		
[Option ID = 36720] 4. Concurrence		
[Option ID = 36721]		
Correct Answer :- Triangulation		
[Option ID = 36720]		
27) The subjectivity of researcher is the soci [Question ID = 9182] 1. Psycho-historical research [Option ID = 36722] 2. Narrative Analysis [Option ID = 36723] 3. Experimental studies [Option ID = 36724] 4. Discourse analysis [Option ID = 36725]	urce of contamination of data in	
Correct Answer :- • Experimental studies [Option ID = 36724]		
28) The objective of generalization of reserving [Question ID = 9183] 1. Phenomenological perspective [Option ID = 36726] 2. Thematic analysis [Option ID = 36727] 3. Qualitative research [Option ID = 36728] 4. Quantitative research [Option ID = 36729]	arch findings are better achieved in	
Correct Answer :- • Quantitative research [Option ID = 36729]		
29)behaviour is that aspect measures as vomiting, taking laxatives, diure [Question ID = 9184] 1. Purging [Option ID = 36730] 2. Regurgitation [Option ID = 36731] 3. Belching [Option ID = 36732] 4. Bruxism [Option ID = 36733]		ninates food from the body by such
Correct Answer :- • Purging [Option ID = 36730]		
30) The analytic process by which concepts research is called:- [Question ID = 9185] 1. Open coding [Option ID = 36734] 2. Axial coding [Option ID = 36735] 3. Network coding [Option ID = 36736] 4. Specific coding [Option ID = 36737]	to observed data and phenomena are	attached during grounded theory
Correct Answer :- Open coding [Option ID = 36734]		
31) An act in which a person who is no long [Question ID = 9186] 1. Mental representation (Option ID = 36738)	er present is copied by children who h	ave witnessed a similar act is called

- 2. Deferred imitation [Option ID = 36739]
- 3. Object permanence [Option ID = 36740]
- 4. Symbolic representation [Option ID = 36741]

Correct Answer :-

Deferred imitation [Option ID = 36739]

32) The process of relating codes (categories and concepts) to each other, via a combination of inductive and deductive reasoning is called

[Question ID = 9187]

- 1. Axial coding [Option ID = 36742]
- 2. Open coding [Option ID = 36743]

www.FirstRanker.com

www.FirstRanker.com

Axial coding [Option ID = 36742]

33) The symbolic interactionism paradigm is a theory that emphasize the:-

[Question ID = 9188]

1. Relationship among individuals in the society

[Option ID = 36746]

2. Exchange of meaning through language and symbols

[Option ID = 36747]

3. Way in which people make sense of their social world

[Option ID = 36748]

4. All of these

[Option ID = 36749]

Correct Answer :-

· All of these

[Option ID = 36749]

34) The attitude of attending systematically to the context of knowledge construction, especially to the effect of the researcher, at every step of research process is referred as

[Question ID = 9189]

- 1. Reflexivity [Option ID = 36750]
- Reflectivity [Option ID = 36751]
- 3. Refractivity [Option ID = 36752]
- Reliability [Option ID = 36753]

Correct Answer :-

· Reflexivity [Option ID = 36750]

35) When leaders go beyond their own self interest and focus on opportunities to help followers grow and develop is called _______ leadership.

[Question ID = 9190]

- 1. Authentic [Option ID = 36754]
- 2. Democratic [Option ID = 36755]
- 3. Servant [Option ID = 36756]
- 4. Transformational [Option ID = 36757]

Correct Answer :-

Servant [Option ID = 36756]

36) When bargaining is operating under the assumption that one or more can create a win-win situation, it is known as:

[Question ID = 9191]

- Distribution [Option ID = 36758]
- 2. Integrative [Option ID = 36759]
- 3. Retributive [Option ID = 36760]
- Summative [Option ID = 36761]

Correct Answer :-

Integrative [Option ID = 36759]

37) The benefit of using 2-way ANOVA over multiple 1-way ANOVA is

[Question ID = 9192]

- 1. Moderation effects can be analysed [Option ID = 36762]
- 2. The power of the test can be increased [Option ID = 36763]
- 3. Multiple comparison of means can be made [Option ID = 36764]
- The error variance can be reduced [Option ID = 36765]

Correct Answer :-

· Moderation effects can be analysed [Option ID = 36762]

38) The number of treatment groups in a 3 x 4 x 4 factorial ANOVA will be

[Question ID = 9193]

- 1. 11 [Option ID = 36766]
- 16 (Option ID = 36767)
 12 (Option ID = 36768)
- 4. 48 [Option ID = 36769]

www.FirstRanker.com

Correct Answer :-

1. State of perceptual anomaly [Option ID = 36790]

State of Déjà vu [Option ID = 36791]
 Dream-like ecstatic state [Option ID = 36792]



45) The human science movement was part of the
45) The human science movement was part of the
[Question ID = 9200] 1. Renaissance
[Option ID = 36794] 2. Naturwissenschaften
[Option ID = 36795] 3. Counter-enlightenment
[Option ID = 36796] 4. Scientific revolution
[Option ID = 36797]
Correct Answer :- • Counter-enlightenment
[Option ID = 36796]
46) While doing the qualitative research, the knowledge symbolize our sense of their meaning in movement, sound, color and shape and story is often referred as knowledge. [Question ID = 9201] 1. Experiential [Option ID = 36798] 2. Practical [Option ID = 36799] 3. Propositional [Option ID = 36800] 4. Presentational [Option ID = 36801]
• Presentational [Option ID = 36801]
47) Which of the following diagnostic criteria has been developed by National Institute of Mental Health, USA? [Question ID = 9202] 1. Research Diagnostic Criteria [Option ID = 36802] 2. Diagnostic Statistical manual [Option ID = 36803] 3. International Classification of Diseases (Mental and Behavioural Disorders) [Option ID = 36804] 4. Research Domain Criteria [Option ID = 36805] Correct Answer: • Research Domain Criteria [Option ID = 36806]
Research bornain Criteria (option to = 30003)
48) Robert Siegler uses the metaphor of
Correct Answer :- • Natural selection, strategy choice [Option ID = 36806]
49) Children have an impressive, as a result of which long before they start to talk, they acquire a stock of speech structures from which they later learn meanings. [Question ID = 9204] 1. Discriminatory learning practice [Option ID = 36810] 2. Statistical learning capacity [Option ID = 36811] 3. Pre-syntactic learning capacity [Option ID = 36812] 4. Perceptual discrimination capacity [Option ID = 36813]
Correct Answer :- • Statistical learning capacity [Option ID = 36811]
50) Which of the following statements about correlation coefficient is not true? [Question ID = 9205] 1. Correlation coefficient range from -1.00 to +1.00 [Option ID = 36814] 2. A correlation of23 is stronger than the correlation of +.45 [Option ID = 36815] 3. A correlation of90 and +.90 is equally strong [Option ID = 36815]

4. Negative correlation means one scores are higher and other scores are lower [Option ID = 36817]



www.FirstRanker.com

