

EAS-101

www.FirstRank www.First (Following Paper ID and Boll No. to be filled in youer.com Answer Book)

Paper ID : 199121

Roll No.	П	П	П	TT	Τ
,					_

B. Tech.

(SEM. I) THEORY EXAMINATION, 2015-16 ENGINEERING PHYSICS-I

[Time: 3 hours] [Total Marks: 100]

SECTION-A

- 1. Attempt all parts. All parts carry equal marks. Write answer of each part in short. (2x10=20)
 - (a) Is earth an inertial frame? Explain.
 - (b) What is significance of mass energy relation?
 - (c) Two independent sources cannot produce interference. Why?
 - (d) What happen when young doubel slit experiment immersed in water?
 - (e) What will be the effect on the intensity of principal maxima of diffraction pattern when single slit is replaced by double slit?

4500 (1) P.T.O.

MMM Files Ranker con

FR	FirstRanker.c	0
1500	Firstranker's choice www.FirstRan	nk
(2)	E wedge shape thin film seen by normally reflected sodium collight. In Newton's ring experiment the diameter of 4th and 12th dark ring are 0.400 cm and 0.700 cm respectively. Deduce the diameter of 20th ring. Obtain intensity expression for single slit Fraunhofer diffraction pattern. Light of wavelength 5500Å falls normally on a slit of width 22.0×10-5 cm. Calculate the	Discuss the formation of interference fringes due to a
EAS-101	sodium gr of 4th 00 cm unhofer Å falls late the	hie to a

4500

P.T.O.

11. (a) Explain the formation of spectrum by plane

diffraction grating.

(c) Find the mass and speed of electron with energy

of 2 MeV.

holography.

of image on hologram. Also describe application of

Describe the process of construction and reconstruction

Attempt any five questions from this section. (10x5=50)

Derive the expression for time dilation. Show that it is a

real effect.

Deduce an expression for variation of mass with

velocity. At what speed the mass of an object becomes

2.25 times its rest mass

SECTION-C

Attempt any two questions from this section. (15x2=30)10. (a) Show that space time or interval between two (b) Show the effect of introducing thin mica sheet in double slit experiment. the path of one of interfering beam in young's events remains invariant under Lorentz transformation.

MNMKIRSK

www.FirstRank

er.com



FirstRanker.com

Explain how wavelength of sodium light can be www.FirstRank calculated from legench biprisim experiment.

- If the population ratio between two energy states (c) in Ruby laser is 2×10⁻⁴⁰, emitting a light beam of wavelength 6943Å. Find the temperature of energy states.
- 12. (a) Discuss the difference between single mode fibre and multimode fibre.
 - Explain construction and working of Nicol prism. (b)
 - (c) A step index fibre has core refractive index 1.466, cladding refractive indes 1.460, compute the maximum radius allowed, if fibre support only one mode by using wavelength of 5000Å.

Physical Constants:

Mass of electron, $m_0 = 9.1 \times 10^{-31} \text{ kg}$

Mass of proton, $m_p = 1.67 \times 10^{-27} \text{ kg}$

 $= 3 \times 10^8 \text{ m/s}$ Speed of light, c

Planck's constant, $h = 6.63 \times 10^{-34} \text{ J/s}$ Charge on electron, $e = 1.6 \times 10^{-19} \text{ C}$

Boltzmann's constant = $1.38 \times 10^{-23} \text{m}^2 \text{kgs}^{-2} \text{K}^{-1}$

-x--

(4) 4500 **EAS-101**

www.FirsiRanker.com