



DU PhD in Chemistry

Topic:- DU\_118\_PHD\_CHEM

1) Which of the following statements about sulfur dioxide is true?

[Question ID = 677]

1. It forms a S-S dimer in condensed phase [Option ID = 2707]
2. Its anhydride of sulfuric acid [Option ID = 2706]
3. Its O-S-O angle is  $180^\circ$  [Option ID = 2708]
4. It is a product of the combustion of fossil fuels that contain sulfur [Option ID = 2705]

Correct Answer :-

- It is a product of the combustion of fossil fuels that contain sulfur [Option ID = 2705]

2) Which of the following is a strong acid in pure liquid HF

[Question ID = 683]

1.  $H_2O$  [Option ID = 2731]
2.  $NaF$  [Option ID = 2729]
3.  $CH_3COOH$  [Option ID = 2730]
4.  $SbF_5$  [Option ID = 2732]

Correct Answer :-

- $SbF_5$  [Option ID = 2732]

3) Each of the following molecules can act as a chelating agent EXCEPT [Question ID = 679]

1.  $HC(CH_2CH_2NH_2)_3$  [Option ID = 2716]
2.  $CH_3NHCH_2CH_2CH_3$  [Option ID = 2714]
3.  $N(CH_2CH_2NH_2)_3$  [Option ID = 2713]
4.  $H_2NCH_2CH_2NH_2$  [Option ID = 2715]

Correct Answer :-

- $CH_3NHCH_2CH_2CH_3$  [Option ID = 2714]

4) What is correct about h-index?

[Question ID = 758]

1. Alternative of impact factor [Option ID = 3031]
2. Based on most quoted papers [Option ID = 3030]
3. Quantify scientific productivity [Option ID = 3029]
4. All of these [Option ID = 3032]

Correct Answer :-

- All of these [Option ID = 3032]

5) The hyperfine electron spin resonance (e.s.r.) spectrum of the benzene radical has how many lines? [Question ID = 748]

1. 12 [Option ID = 2992]
2. 7 [Option ID = 2990]
3. 1 [Option ID = 2991]
4. 6 [Option ID = 2989]

Correct Answer :-

- 7 [Option ID = 2990]



6) The energy changes involving the core electrons of an atom or molecule are expressed in which region of the electromagnetic spectrum? [Question ID = 742]

1. Ultraviolet and Visible region [Option ID = 2967]
2. X-ray region [Option ID = 2968]
3. Radiofrequency region [Option ID = 2966]
4. Infra-red region [Option ID = 2965]

Correct Answer :-

- X-ray region [Option ID = 2968]

7) Find out the expected intensity ratio of M and M+1 signal for the Naphthalene molecular ion [Question ID = 726]

1. 99:1.1 [Option ID = 2903]
2. 1.1:99 [Option ID = 2904]
3. 9:01 [Option ID = 2901]
4. 1:9 [Option ID = 2902]

Correct Answer :-

- 9:01 [Option ID = 2901]

8) Cobalt-60 is used in radiation therapy of cancer and can be produced by the bombardment of Cobalt-59 with [Question ID = 692]

1. Alpha particles [Option ID = 2765]
2. Beta particles [Option ID = 2767]
3. Neutrons [Option ID = 2766]
4. Gamma rays [Option ID = 2768]

Correct Answer :-

- Neutrons [Option ID = 2766]

9) The standard emf of galvanic cell involving 3 moles of electrons in its redox reaction is 0.59 V. The equilibrium constant for the reaction of the cell is- [Question ID = 763]

1.  $10^{15}$  [Option ID = 3051]
2.  $10^{30}$  [Option ID = 3052]
3.  $10^{25}$  [Option ID = 3049]
4.  $10^{20}$  [Option ID = 3050]

Correct Answer :-

- $10^{30}$  [Option ID = 3052]

10) A characteristic common to polymers that can be made to conduct electricity such as polyacetylene, polypyrrole is: [Question ID = 685]

1. Conjugation throughout the polymeric chain. [Option ID = 2740]
2. A high degree of cross linking [Option ID = 2738]
3. A very low glass transition temperature [Option ID = 2737]
4. Presence of stereogenic centers of the same configuration [Option ID = 2739]

Correct Answer :-

- Conjugation throughout the polymeric chain. [Option ID = 2740]

11) Impact factor is [Question ID = 768]

1. Ratio between citations and recent citable items publish [Option ID = 3071]
2. All of these [Option ID = 3072]
3. Addition of citations and recent citable items publish [Option ID = 3069]
4. Ratio between recent citable items publish and citations [Option ID = 3070]

Correct Answer :-

- Ratio between recent citable items publish and citations [Option ID = 3070]

12) On the basis of oxidation-reduction potential, which of the following is most likely to occur? [Question ID = 693]

1.  $\text{Al(s)} + 3\text{NaNO}_3\text{(aq)} \rightarrow 3\text{Na(s)} + \text{Al(NO}_3)_3\text{(aq)}$  [Option ID = 2770]
2.  $\text{Ca(s)} + 2\text{NaNO}_3\text{(aq)} \rightarrow 2\text{Na(s)} + \text{Ca(NO}_3)_2\text{(aq)}$  [Option ID = 2772]
3.  $\text{Pb(s)} + 2\text{LiNO}_3\text{(aq)} \rightarrow 2\text{Li(s)} + \text{Pb(NO}_3)_2\text{(aq)}$  [Option ID = 2771]
4.  $\text{Zn(s)} + 2\text{AgNO}_3\text{(aq)} \rightarrow 2\text{Ag(s)} + \text{Zn(NO}_3)_2\text{(aq)}$  [Option ID = 2769]

Correct Answer :-

- $\text{Zn(s)} + 2\text{AgNO}_3\text{(aq)} \rightarrow 2\text{Ag(s)} + \text{Zn(NO}_3)_2\text{(aq)}$  [Option ID = 2769]

13) How many diastereoisomers are possible for the compound 2, 4 -diphenylcyclobutane-1, 3 di carboxylic acids. [Question ID = 725]

1. 6 [Option ID = 2899]
2. 5 [Option ID = 2898]
3. 8 [Option ID = 2900]
4. 4 [Option ID = 2897]

Correct Answer :-

- 5 [Option ID = 2898]

14) An increase in equivalent conductance of a strong electrolyte with dilution is mainly due to- [Question ID = 764]

1. Increase in ionic mobility of ions [Option ID = 3055]
2. Increase in number of ions [Option ID = 3054]
3. 100% ionization of electrolyte at normal dilution [Option ID = 3056]
4. Increase in both i.e. number of ions and ionic mobility of ions. [Option ID = 3053]

Correct Answer :-

- Increase in ionic mobility of ions [Option ID = 3055]

15) The solid state structures of the principal allotropes of elemental boron are made up of which of the following structural units [Question ID = 699]

1.  $\text{B}_4$  tetrahedra [Option ID = 2796]
2.  $\text{B}_6$  octahedra [Option ID = 2795]
3.  $\text{B}_8$  cubes [Option ID = 2794]
4.  $\text{B}_{12}$  icosahedra [Option ID = 2793]

Correct Answer :-

- $\text{B}_{12}$  icosahedra [Option ID = 2793]

16) The molecular geometry of thionyl chloride is best described as [Question ID = 688]

1. T-shaped [Option ID = 2752]
2. Tetrahedral [Option ID = 2751]
3. Trigonal pyramidal [Option ID = 2749]
4. Trigonal planar [Option ID = 2750]

Correct Answer :-

- Trigonal pyramidal [Option ID = 2749]

17) In a face-center cubic (FCC) type of crystal lattice, the number of atoms belonging exclusively to each unit cell within the lattice is/are: [Question ID = 754]

1. 2 [Option ID = 3014]
2. 1 [Option ID = 3013]
3. 3 [Option ID = 3015]
4. 4 [Option ID = 3016]

Correct Answer :-

- 4 [Option ID = 3016]

18) Among the following, the weakest oxidizing agent is [Question ID = 675]

1. Mg (s) [Option ID = 2698]
2.  $I_2$  (s) [Option ID = 2699]
3.  $H^+$  (aq) [Option ID = 2700]
4.  $MnO_4^-$  (aq) [Option ID = 2697]

Correct Answer :-

- Mg (s) [Option ID = 2698]

19) For a polymer, which of the following statement/s is/are true? [Question ID = 759]

1. Weight average molecular weight is almost always higher than the number average molecular weight [Option ID = 3035]
2. Formation of a polypeptide from its monomers (amino acids) is an example of addition polymerization [Option ID = 3034]
3. All of these [Option ID = 3036]
4. Vinyl polymerization is an example of condensation polymerization. [Option ID = 3033]

Correct Answer :-

- Weight average molecular weight is almost always higher than the number average molecular weight [Option ID = 3035]

20) Quantum dots are [Question ID = 762]

1. Three dimensional [Option ID = 3048]
2. One dimensional [Option ID = 3046]
3. Two dimensional [Option ID = 3047]
4. Zero dimensional [Option ID = 3045]

Correct Answer :-

- Zero dimensional [Option ID = 3045]

21) The unit of rate constant for a third order reaction is: [Question ID = 749]

1.  $s^{-1}$  [Option ID = 2993]
2.  $mol^{-1} dm^3 s^{-1}$  [Option ID = 2995]
3.  $mol^{-2} dm^6 s^{-1}$  [Option ID = 2996]
4.  $mol dm^{-3} s^{-1}$  [Option ID = 2994]

Correct Answer :-

- $mol^{-2} dm^6 s^{-1}$  [Option ID = 2996]

22) All the following elements have at least one isotope that is not radioactive EXCEPT [Question ID = 673]

1. Pb [Option ID = 2690]
2. O [Option ID = 2689]
3. Sn [Option ID = 2691]
4. No [Option ID = 2692]

Correct Answer :-

- No [Option ID = 2692]

23) The conditions for a species to follow Bose-Einstein statistics are; [Question ID = 736]

1. Particles are indistinguishable, with no restriction on filling up of energy levels [Option ID = 2944]
2. Particles are indistinguishable, with a restriction on filling up of energy levels [Option ID = 2943]
3. Particles are distinguishable, with a restriction on filling up of energy levels [Option ID = 2941]
4. Particles are distinguishable, with no restriction on filling up of energy levels [Option ID = 2942]

Correct Answer :-

- Particles are indistinguishable, with no restriction on filling up of energy levels [Option ID = 2944]

24) In the kinetic theory of collisions, the SI unit of collision number, in terms of m (meter) and s (second), is:

[Question ID = 761]

1.  $m^{-2}s^{-1}$  [Option ID = 3042]
2.  $m^4s^{-1}$  [Option ID = 3041]
3.  $m^2s^{-1}$  [Option ID = 3043]
4. None of these [Option ID = 3044]

Correct Answer :-

- None of these [Option ID = 3044]

25) Correct characteristics of the functional groups of adenine in DNA base pair are [Question ID = 706]

1. Both N(3) and C(6)NH<sub>2</sub> are hydrogen bond donors. [Option ID = 2824]
2. N(3) is a hydrogen bond acceptor and C(6)NH<sub>2</sub> is a hydrogen bond donor. [Option ID = 2821]
3. Both N(3) and C(6)NH<sub>2</sub> are hydrogen bond acceptors [Option ID = 2823]
4. N(1) is a hydrogen bond acceptor and C(6)NH<sub>2</sub> is a hydrogen bond donor. [Option ID = 2822]

Correct Answer :-

- N(1) is a hydrogen bond acceptor and C(6)NH<sub>2</sub> is a hydrogen bond donor. [Option ID = 2822]

26) The carbon monoxide molecule has an internuclear distance of 1.13 Angstroms. What is the moment of Inertia of this molecule? [Question ID = 740]

1.  $21.6 \times 10^{-47} \text{ kgm}^2$  [Option ID = 2960]
2.  $14.5 \times 10^{-47} \text{ kgm}^2$  [Option ID = 2957]
3.  $14.5 \times 10^{-47} \text{ kgm}^2$  [Option ID = 2958]
4.  $1.45 \times 10^{-47} \text{ kgm}^2$  [Option ID = 2959]

Correct Answer :-

- $14.5 \times 10^{-47} \text{ kgm}^2$  [Option ID = 2957]

27) Which of the following represent/s non-linear optical technique? [Question ID = 744]

1. Second Harmonic generation [Option ID = 2974]
2. Two-photon photoluminescence [Option ID = 2975]
3. Four-wave mixing [Option ID = 2973]
4. All of these [Option ID = 2976]

Correct Answer :-

- All of these [Option ID = 2976]


28) . Which of the following does not affect the intensity of spectral lines of a sample? [Question ID = 743]

1. Path length of a sample [Option ID = 2972]
2. Population of energy states [Option ID = 2970]
3. Heisenberg's Uncertainty principle [Option ID = 2971]
4. Concentration of a sample [Option ID = 2969]


Correct Answer :-

- Heisenberg's Uncertainty principle [Option ID = 2971]


29) Find out the major product of the following reaction




[Question ID = 716]




1. [Option ID = 2862]



2. [Option ID = 2863]




3. [Option ID = 2861]



4. [Option ID = 2864]

Correct Answer :-



\* [Option ID = 2864]

30) Provide the suitable reagents for this conversion:



[Question ID = 712]

 1.  $\text{NaNO}_2 / \text{H}_2\text{SO}_4 / \text{PCl}_3$  [Option ID = 2845]

 2.  $\text{H}_2\text{O}_2 / \text{OH}^-$ ,  $\text{HNO}_3 / \text{H}_2\text{SO}_4 / \text{PCl}_3$  [Option ID = 2846]

 3.  $\text{HNO}_3 / \text{H}_2\text{SO}_4 / \text{POCl}_3$  [Option ID = 2848]

 4. m-CPBA,  $\text{HNO}_3 / \text{H}_2\text{SO}_4 / \text{PCl}_3$  [Option ID = 2847]

Correct Answer :-

 \* m-CPBA,  $\text{HNO}_3 / \text{H}_2\text{SO}_4 / \text{PCl}_3$  [Option ID = 2847]

 31) Which of the following complexes does not contain a significant  $\pi$  component in the metal-ligand bonding?

[Question ID = 686]

 1.  $[\text{Co}(\text{NH}_3)_6]^{3+}$  [Option ID = 2743]

 2.  $[\text{Cr}(\eta\text{-C}_6\text{H}_6)]$  [Option ID = 2742]


 3.  $[\text{Co}(\text{CN})_3]^{3-}$  [Option ID = 2744]

4.  $[\text{Fe}(\text{CO})_5]$  [Option ID = 2744]





Correct Answer :-

\*  $[\text{Co}(\text{NH}_3)_6]^{3+}$  [Option ID = 2743]


32) The product obtained in the following conversion is:




[Question ID = 5633]



Correct Answer :-



33) In the multi-step synthesis given below, the overall yield for the formation of S from P is:



[Question ID = 730]

1. 40 % [Option ID = 2918]

2. 50 % [Option ID = 2920]

3. 72 % [Option ID = 2917]


4. 36 % [Option ID = 2919]

Correct Answer :-






• 36 % [Option ID = 2919]


34) The compound formed in the following reaction is:



[Question ID = 714]



Correct Answer :-


 35) A 499 mg sample of  $\text{CuSO}_4 \cdot n\text{H}_2\text{O}$  is heated to drive off the waters of hydration and then reweighed to give a final mass of 319 mg. Given the sample contains 2.0 mmol of Cu, what is the average number of waters of hydration,  $n$  in  $\text{CuSO}_4 \cdot n\text{H}_2\text{O}$ ?

[Question ID = 669]

1. 2 [Option ID = 2673]
2. 18 [Option ID = 2676]
3. 5 [Option ID = 2674]
4. 10 [Option ID = 2675]

Correct Answer :-

- 5 [Option ID = 2674]

 36) What is the orbital angular momentum quantum number  $l$  of the electron that is most easily removed when ground state aluminium is ionized?

[Question ID = 689]

1. 2 [Option ID = 2754]
2. 0 [Option ID = 2756]
3. 1 [Option ID = 2755]



4. 3 [Option ID = 2753]


Correct Answer :-

• 1 [Option ID = 2755]


37) The major product obtained in the following reaction is:




[Question ID = 718]



1. C) [Option ID = 2871]




2. B) [Option ID = 2870]



3. A) [Option ID = 2869]

4. D) both (B) and (C) [Option ID = 2872]


Correct Answer :-




• A) [Option ID = 2869]

38)


Predict the major product:




[Question ID = 709]




1. [Option ID = 2836]



2. [Option ID = 2833]




3. [Option ID = 2835]




4. [Option ID = 2834]

Correct Answer :-



[Option ID = 2836]

39) The IUPAC name of the compound given below is:



[Question ID = 705]

1. (2Z, 4Z)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 2819]
2. (2E, 4E)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 2817]
3. (2Z, 4E)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 2818]
4. (2Z, 4E)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 2820]

Correct Answer :-

- (2Z, 4E)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 2820]

40) Which of the following statements about complexes that form between metals  $M^{n+}$  and EDTA in aqueous solutions is true?

[Question ID = 680]

1. The presence of other complexing ligands in solution affects the equilibrium concentration of metal-EDTA complexes [Option ID = 2719]
2. Metal-EDTA complexes have an equilibrium concentration independent of pH [Option ID = 2718]
3. Metal-EDTA complexes are often 2:1 in stoichiometry [Option ID = 2717]
4. Metal-EDTA complexes are less stable than the corresponding metal-ammine complexes [Option ID = 2720]

Correct Answer :-

- The presence of other complexing ligands in solution affects the equilibrium concentration of metal-EDTA complexes [Option ID = 2719]

41) A 0.600 g sample of pure, weak diprotic acid gives end points at 20.0 mL and 40.0 mL when titrated with 0.100 M NaOH. What is the molar mass of the weak acid?

[Question ID = 671]

1. 150 g [Option ID = 2682]
2. 300 g [Option ID = 2684]
3. 120 g [Option ID = 2681]
4. 180 g [Option ID = 2683]

Correct Answer :-

- 300 g [Option ID = 2684]

42) The microwave spectrum of a rigid diatomic molecule shows first three lines at  $2.65682 \text{ cm}^{-1}$ ,  $5.31364 \text{ cm}^{-1}$ , and  $7.97046 \text{ cm}^{-1}$ . What is the rotational constant of this molecule?

[Question ID = 756]

1.  $1.82118 \text{ cm}^{-1}$  [Option ID = 3021]
2.  $3.64236 \text{ cm}^{-1}$  [Option ID = 3022]
3.  $1.32841 \text{ cm}^{-1}$  [Option ID = 3024]
4.  $0.91059 \text{ cm}^{-1}$  [Option ID = 3023]

Correct Answer :-

1.32841  $\text{cm}^{-1}$  [Option ID = 3024]

43) It takes 10 minutes for the concentration of a radioactive species to decay to its  $1/4^{\text{th}}$  value of its original concentration. What is the rate constant of this radioactive decay reaction?


[Question ID = 750]

1. 415.8  $\text{s}^{-1}$  [Option ID = 2999]
2. 865.8  $\text{s}^{-1}$  [Option ID = 3000]
3. 0.00231  $\text{s}^{-1}$  [Option ID = 2997]
4. 0.001155  $\text{s}^{-1}$  [Option ID = 2998]

Correct Answer :-

0.00231  $\text{s}^{-1}$  [Option ID = 2997]


44) The major product in the following reaction is:



[Question ID = 715]


1. [Option ID = 2857]
2. [Option ID = 2860]
3. [Option ID = 2858]
4. [Option ID = 2859]

Correct Answer :-




45)


The product obtained in the following reaction is




[Question ID = 719]




[Option ID = 2873]



[Option ID = 2876]



[Option ID = 2874]



[Option ID = 2875]

Correct Answer :-

46)  $\text{PbF}_2(\text{s})$  which is slightly soluble in water is dissolved in water to form a standard solution in equilibrium with solid  $\text{PbF}_2$ . Which of the following will cause additional  $\text{PbF}_2(\text{s})$  to dissolve?

[Question ID = 674]

1. Evaporating some water to decrease the volume of the solution. [Option ID = 2696]

2. Adding solid  $\text{PbF}_2$  [Option ID = 2695]

3. Adding  $\text{Pb}(\text{NO}_3)_2$  [Option ID = 2694]

4. Adding  $\text{HNO}_3$  [Option ID = 2693]

Correct Answer :-

Adding  $\text{HNO}_3$  [Option ID = 2693]

47) Arrange the following intermediates in the order of decreasing basicity (strongest to weakest):

(i)  $\text{H}_2\text{C}=\text{CH}^-$  (ii)  $\text{CH}_3\text{CH}_2^-$  (iii)  $\text{CH}_3\text{CH}_2\text{O}^-$  (iv)  $\text{HC}\equiv\text{C}^-$

[Question ID = 728]

1.  $\text{iii} > \text{ii} > \text{i} > \text{iv}$  [Option ID = 2910]

2.  $\text{iii} > \text{iv} > \text{i} > \text{ii}$  [Option ID = 2911]

3.  $\text{iv} > \text{i} > \text{ii} > \text{iii}$  [Option ID = 2909]

4.  $\text{ii} > \text{i} > \text{iv} > \text{iii}$  [Option ID = 2912]

Correct Answer :-

- ii > i > iv > iii [Option ID = 2912]

48) For EDTA titrations, the analyte solution and the titrant solution are both buffered at the same pH for which of the following reasons:

- I. Conditional formation constant is affected by pH.
- II. The fraction of EDTA in the fully deprotonated  $Y^{4-}$  form varies with pH.
- III. When EDTA is complexed with metal ions,  $H^+$  ions are formed as product.

[Question ID = 697]

1. III only [Option ID = 2788]
2. II only [Option ID = 2787]
3. I only [Option ID = 2785]
4. I, II and III [Option ID = 2786]

Correct Answer :-

- I, II and III [Option ID = 2786]

49) When  $Fe_2O_3$  is dissolved in 6M  $HNO_3$ , which iron containing species dominate in the solution?

[Question ID = 700]

1.  $Fe(OH)_3$  [Option ID = 2798]
2.  $Fe(OH)_4^-$  [Option ID = 2797]
3.  $Fe(H_2O)_6^{3+}$  [Option ID = 2800]
4.  $Fe(H_2O)_6^{2+}$  [Option ID = 2799]

Correct Answer :-

- $Fe(H_2O)_6^{2+}$  [Option ID = 2799]

50) In  $CrF_2(s)$ , the coordination of six  $F^-$ , around the Cr is a distorted octahedron with four short and two long Cr-F bonds. Which of the following best explains this observation?


[Question ID = 678]

1.  $Cr^{2+}$  has a low cationic charge [Option ID = 2711]
2. F has -1 anionic charge and highly electronegative [Option ID = 2709]
3. Spin-orbit coupling in  $Cr^{2+}$  [Option ID = 2712]
4. The Jahn-Teller effect [Option ID = 2710]


Correct Answer :-

- The Jahn-Teller effect [Option ID = 2710]


51) The major product formed in the following reaction is:




[Question ID = 723]




1. [Option ID = 2889]



2. [Option ID = 2890]




3. [Option ID = 2891]



4. [Option ID = 2892]


Correct Answer :-




\* [Option ID = 2889]

- 52) The compound showing the following spectral characteristic is  $^1\text{H}$  NMR ( $\delta$  in ppm): 4.65 (2H, singlet), 3.65 (4H, quartet), 1.25 (6H, triplet);  $^{13}\text{C}$  NMR ( $\delta$  in ppm) = 15, 63, 95; DEPT-135 ( $\delta$  in ppm): 15 (positive), 63 (negative), 95 (negative); DEPT-90 ( $\delta$  in ppm): 15 (no peak), 63 (no peak), 95 (no peak).


[Question ID = 727]



1. [Option ID = 2907]



2. [Option ID = 2906]




3. [Option ID = 2905]




4. [Option ID = 2908]

Correct Answer :-



\* [Option ID = 2905]

- 53) In low chloride ion concentration, the anticancer drug cis-platin hydrolysis to give a diaqua complex and this binds to DNA via adjacent guanine.



The coordinating atom of guanine to Pt(II) is

[Question ID = 707]



1. N9 [Option ID = 2828]
2. N7 [Option ID = 2827]
3. N1 [Option ID = 2825]
4. N3 [Option ID = 2826]

Correct Answer :-

- N7 [Option ID = 2827]

54) The molecular geometry of  $\text{IF}_5$  is


[Question ID = 672]

1. Bicapped prism [Option ID = 2688]
2. Square pyramidal [Option ID = 2686]
3. Trigonal planar [Option ID = 2685]
4. Bent [Option ID = 2687]

Correct Answer :-

- Square pyramidal [Option ID = 2686]

55) What is the principal product of the following reaction?



[Question ID = 704]

1. [Option ID = 2816]
2. [Option ID = 2814]
3. [Option ID = 2813]
4. [Option ID = 2815]

Correct Answer :-


- [Option ID = 2813]

56) The major product of the reaction given below is:







[Question ID = 708]




[Option ID = 2829]



[Option ID = 2832]




[Option ID = 2831]




[Option ID = 2830]

Correct Answer :-




[Option ID = 2831]


57) The product obtained in the following conversion is:




[Question ID = 717]




[Option ID = 2866]



[Option ID = 2865]




[Option ID = 2867]



[Option ID = 2868]


Correct Answer :-




[Option ID = 2866]

58)


The major product in the following reaction is:




[Question ID = 711]




1. [Option ID = 2843]



2. [Option ID = 2842]




3. [Option ID = 2844]



4. [Option ID = 2841]

Correct Answer :-




[Option ID = 2842]


59) The product obtained in the following conversion is:



[Question ID = 710]




1. [Option ID = 2838]



2. [Option ID = 2839]




3. [Option ID = 2837]



4. [Option ID = 2840]

Correct Answer :-



[Option ID = 2838]

 60) The ionic strength of an aqueous 0.10 M  $\text{Pb}(\text{NO}_3)_2$  solution is


[Question ID = 682]

1. 0.30 M [Option ID = 2728]
2. 0.25 M [Option ID = 2727]
3. 0.60 M [Option ID = 2725]
4. 0.10 M [Option ID = 2726]


Correct Answer :-

- 0.30 M [Option ID = 2728]


61) Find out the product of the following reaction




[Question ID = 772]




[Option ID = 3087]



[Option ID = 3088]




[Option ID = 3086]




[Option ID = 3085]

Correct Answer :-




[Option ID = 3086]


62) The major product formed in the following reaction is



[Question ID = 733]



[Option ID = 2932]



Correct Answer :-



- 63) The energies of activation for forward and reverse reactions for  $A_2 + B_2 \rightarrow 2AB$  are  $180 \text{ kJ mol}^{-1}$  and  $200 \text{ kJ mol}^{-1}$  respectively. The presence of a catalyst lowers the activation energy of both (forward and reverse) reactions by  $100 \text{ kJ mol}^{-1}$ . The enthalpy change of the reaction ( $A_2 + B_2 \rightarrow 2AB$ ) in the presence of catalyst will be (in  $\text{kJ mol}^{-1}$ ):


[Question ID = 767]

- 120 [Option ID = 3065]
- 280 [Option ID = 3066]
- 300 [Option ID = 3068]
- 20 [Option ID = 3067]

Correct Answer :-

- 20 [Option ID = 3067]

- 64) The amino acid constituents of artificial sweetener given below are



[Question ID = 734]

- L-Aspartic acid and L-tyrosine [Option ID = 2936]
- D-Glutamic acid and L-phenylglycine [Option ID = 2933]
- L-Glutamic acid and L-phenylalanine [Option ID = 2934]
- L-Aspartic and L-phenylalanine [Option ID = 2935]

Correct Answer :-

- L-Aspartic and L-phenylalanine [Option ID = 2935]

- 65) Graphite reacts with potassium to produce a compound with empirical formula  $KC_8$  of the following which is the best description of this structure:


[Question ID = 676]

1.  $K^+$  ion packed with  $C_2^{2-}$  ions [Option ID = 2702]
2. Negatively charged hexagonal carbon layers with intercalated  $K^+$  ions [Option ID = 2703]
3. An expanded diamond lattice with  $K^+$  ions in the tetrahedral holes [Option ID = 2704]
4.  $K^+$  -ion closed packed with polyhedral  $C_8^{2-}$  ions [Option ID = 2701]

Correct Answer :-

- Negatively charged hexagonal carbon layers with intercalated  $K^+$  ions [Option ID = 2703]

66) The IUPAC name for the following molecule is:



[Question ID = 729]

1. (2Z, 4Z)-3, 4-dibromo hepta-2, 4-diene [Option ID = 2914]
2. (2E, 4E)-3, 4-dibromo hepta-2, 4-diene [Option ID = 2915]
3. (2E, 4Z)-3, 4-dibromo hepta-2, 4-diene [Option ID = 2916]
4. (2Z, 4E)-3, 4-dibromo hepta-2, 4-diene [Option ID = 2913]

Correct Answer :-

- (2E, 4Z)-3, 4-dibromo hepta-2, 4-diene [Option ID = 2916]

67) Saturated solution of  $KNO_3$  is used to make 'salt bridge' because-


[Question ID = 765]

1.  $KNO_3$  is highly soluble in water [Option ID = 3060]
2. velocity of  $K^+$  is greater than that of  $NO_3^-$  [Option ID = 3057]
3. velocity of  $NO_3^-$  is greater than that of  $K^+$  [Option ID = 3058]
4. velocity of both  $K^+$  and  $NO_3^-$  are nearly the same [Option ID = 3059]


Correct Answer :-

- velocity of both  $K^+$  and  $NO_3^-$  are nearly the same [Option ID = 3059]


68) In the following reaction the major product formed is:




[Question ID = 724]




1. [Option ID = 2893]



2. [Option ID = 2895]




3. [Option ID = 2896]




4. [Option ID = 2894]

Correct Answer :-




5. [Option ID = 2893]


69) The product formed in the following reaction is:




[Question ID = 721]




1. [Option ID = 2881]



2. [Option ID = 2884]




3. [Option ID = 2882]



4. [Option ID = 2883]

Correct Answer :-



5. [Option ID = 2882]

- 70) In a zero-order reaction for every  $10^\circ\text{C}$  rise of temperature, the rate is doubled. If the temperature is increased from  $10^\circ\text{C}$  to  $100^\circ\text{C}$ , the rate of the reaction will become-

[Question ID = 766]

1. 512 times [Option ID = 3064]
2. 256 times [Option ID = 3063]
3. 128 times [Option ID = 3062]
4. 64 times [Option ID = 3061]

Correct Answer :-

- 512 times [Option ID = 3064]



The reaction is not balanced. If the reaction is balanced using the smallest whole number coefficients possible, the coefficients for  $\text{I}^-$  will be:




[Question ID = 670]

1. 2 [Option ID = 2678]
2. 5 [Option ID = 2680]
3. 1 [Option ID = 2677]
4. 3 [Option ID = 2679]

Correct Answer :-

- 5 [Option ID = 2680]

- 72) Arrange the following compounds in decreasing order of IR stretching frequency of  $\text{C}=\text{O}$



[Question ID = 731]

1.  $\text{iv} > \text{i} > \text{ii} > \text{iii}$  [Option ID = 2924]
2.  $\text{ii} > \text{i} > \text{iii} > \text{iv}$  [Option ID = 2923]
3.  $\text{iii} > \text{i} > \text{ii} > \text{iv}$  [Option ID = 2922]
4.  $\text{i} > \text{ii} > \text{iii} > \text{iv}$  [Option ID = 2921]

Correct Answer :-

- $\text{iii} > \text{i} > \text{ii} > \text{iv}$  [Option ID = 2922]



The correct balanced one will be:

[Question ID = 690]


1.  $\text{MnO}_4^- : \text{IO}_3^-$  is 1:1 [Option ID = 2760]
2.  $\text{MnO}_4^- : \text{Mn}^{2+}$  is 3:1 [Option ID = 2759]
3.  $\text{I}^- : \text{IO}_3^-$  in 3:1 [Option ID = 2757]
4.  $\text{MnO}_4^- : \text{I}^-$  in 6:5 [Option ID = 2758]



Correct Answer :-

 •  $\text{MnO}_4^- : \text{I}^-$  in 6:5 [Option ID = 2758]

74) In the following reaction sequence, the structure of the product is:



[Question ID = 720]

1. [Option ID = 2879]

2. [Option ID = 2878]


3. [Option ID = 2880]

4. [Option ID = 2877]

Correct Answer :-

• [Option ID = 2877]

75) The major product formed in the sulphuric acid mediated rearrangement of compound is:



[Question ID = 722]


1. [Option ID = 2886]

2. [Option ID = 2888]

3. [Option ID = 2885]

4. [Option ID = 2887]

Correct Answer :-



[Option ID = 2887]

76) What is the specific resistance (or resistivity) of a conductor with cross-sectional area 4 cm<sup>2</sup>, length 2cm and resistance 10 ohms?

[Question ID = 755]

1. 20 Siemens<sup>-1</sup>cm [Option ID = 3019]
2. 10 Siemens<sup>-1</sup>cm [Option ID = 3018]
3. None of the above [Option ID = 3020]
4. 40 Siemens<sup>-1</sup>cm [Option ID = 3017]

Correct Answer :-

1. 20 Siemens<sup>-1</sup>cm [Option ID = 3019]

77) The anhydride of Ba(OH)<sub>2</sub> is

[Question ID = 695]


1. Ba [Option ID = 2779]
2. BaO [Option ID = 2780]
3. BaOH [Option ID = 2778]
4. BaH<sub>2</sub> [Option ID = 2777]

Correct Answer :-


1. BaO [Option ID = 2780]

78) A compound with molecular formula C<sub>4</sub>H<sub>6</sub>O<sub>2</sub> shows band at 1770 cm<sup>-1</sup> in IR spectra and peaks at 178, 68, 28, 22 ppm in <sup>13</sup>C NMR spectrums. The correct structure of the compound is:


[Question ID = 703]




1. [Option ID = 2811]



2. [Option ID = 2809]




3. [Option ID = 2812]




4. [Option ID = 2810]

Correct Answer :-



[Option ID = 2811]



[Option ID = 2810]

79) An aqua's solution of an optically pure compound of conc. 100 mg in 1 ml of water and measured in sample of 5 cm length was found to be  $-3^\circ$  the specific rotation is

[Question ID = 732]

1.  $-6^\circ\text{C}$  [Option ID = 2927]
2.  $+6^\circ\text{C}$  [Option ID = 2928]
3.  $-60^\circ\text{C}$  [Option ID = 2926]
4.  $-30^\circ\text{C}$  [Option ID = 2925]

Correct Answer :-

- $-60^\circ\text{C}$  [Option ID = 2926]

80) A monoatomic gas following Fermi-Dirac statistics begins to follow Maxwell-Boltzmann statistics at: [Question ID = 735]

1. Low Temperature and low density [Option ID = 2937]
2. High Temperature and high density [Option ID = 2940]
3. Low Temperature and high density [Option ID = 2938]
4. High Temperature and low density [Option ID = 2939]

Correct Answer :-

- High Temperature and low density [Option ID = 2939]

81) The Dulong and Petit's Law says that the molar heat capacity of elements is: [Question ID = 741]

1.  $10 \text{ Cal mol}^{-1} \text{K}^{-1}$  [Option ID = 2964]
2.  $6 \text{ Cal mol}^{-1} \text{K}^{-1}$  [Option ID = 2961]
3.  $12 \text{ Cal mol}^{-1} \text{K}^{-1}$  [Option ID = 2963]
4.  $3 \text{ Cal mol}^{-1} \text{K}^{-1}$  [Option ID = 2962]

Correct Answer :-

- $6 \text{ Cal mol}^{-1} \text{K}^{-1}$  [Option ID = 2961]

82) What is the most common natural form in which fluorine is found on earth?

[Question ID = 684]

1. As a fluoride ion in various minerals [Option ID = 2735]
2. As weak acid HF (aq) [Option ID = 2734]
3. In various fluorocarbon compounds in the atmosphere. [Option ID = 2736]
4. As  $\text{XeF}_2$  (s) [Option ID = 2733]

Correct Answer :-

- As a fluoride ion in various minerals [Option ID = 2735]

83) What is the correct form of Stirling's approximation?

[Question ID = 738]

1.  $\ln x! = x \ln x - x$  [Option ID = 2950]
2.  $\ln x! = \ln x + x$  [Option ID = 2951]

3.  $\ln x! = x \ln x + x$  [Option ID = 2949]

4.  $\ln x! = x - x \ln x$  [Option ID = 2952]

Correct Answer :-

•  $\ln x! = x \ln x - x$  [Option ID = 2950]

84) What is the total energy of one mole of an ideal monoatomic gas in terms of Boltzmann's Constant (k), Avogadro's number (N) and temperature (T)

[Question ID = 739]

1.  $3 NkT$  [Option ID = 2953]
2.  $(3/2) NkT$  [Option ID = 2956]
3.  $(1/2) NkT$  [Option ID = 2955]
4.  $NkT$  [Option ID = 2954]

Correct Answer :-

- $(3/2) NkT$  [Option ID = 2956]

85) The following equation is associated with the relationship between the diffusion current and the concentration of the depolarizer used in polarography: [Question ID = 753]

1. Debye-Huckel equation [Option ID = 3009]
2. Stern-Volmer equation [Option ID = 3010]
3. Nyquist equation [Option ID = 3012]
4. Ilkovic equation [Option ID = 3011]

Correct Answer :-

- Ilkovic equation [Option ID = 3011]

86) Electronic transitions originating from the 1S energy level of the Hydrogen atom to higher levels belong to which series? [Question ID = 747]

1. Lyman Series [Option ID = 2985]
2. Brackett Series [Option ID = 2987]
3. Balmer Series [Option ID = 2986]
4. Pfund Series [Option ID = 2988]


Correct Answer :-

- Lyman Series [Option ID = 2985]

87) Which of the following reactions best classified as an oxidative addition? [Question ID = 701]

1.  $[\text{Cr}(\text{CO})_6] + \text{Br}^- \rightarrow [\text{Cr}(\text{CO})_5\text{Br}]^- + \text{CO}$  [Option ID = 2801]
2.  $[\text{Pt}(\text{NH}_3)\text{Cl}_3]^- + \text{NH}_3 \rightarrow \text{Pt}(\text{NH}_3)_2\text{Cl}_2 + \text{Cl}^-$  [Option ID = 2803]
3.  $[\text{Pt}\{\text{P}(\text{C}_2\text{H}_5)_3\}_2\text{HCl}] + \text{HCl} \rightarrow [\text{Pt}\{\text{P}(\text{C}_2\text{H}_5)_3\}_2(\text{H})_2\text{Cl}_2]$  [Option ID = 2802]
4.  $[\text{MnH}(\text{CO})_5] + \text{CF}_2 = \text{CF}_2 \rightarrow [\text{Mn}(\text{CF}_2\text{CF}_2\text{H})(\text{CO})_5]$  [Option ID = 2804]

Correct Answer :-



88) Which of the following is required for both paramagnetism and ferromagnetism? [Question ID = 698]

1. Super exchange [Option ID = 2791]
2. unpaired electrons [Option ID = 2792]
3. Low-spin electron configuration [Option ID = 2790]
4. Strong oxidizing conditions [Option ID = 2789]

Correct Answer :-

- unpaired electrons [Option ID = 2792]

89) Which of the following experimental techniques is not used to determine the average molecular weight of a polymer? [Question ID = 760]

1. Transmission electron microscopy [Option ID = 3039]
2. Equilibrium sedimentation [Option ID = 3038]
3. Intrinsic viscosity measurement [Option ID = 3040]
4. Dynamic light scattering [Option ID = 3037]

Correct Answer :-

- Transmission electron microscopy [Option ID = 3039]

90) Which of the following is NOT a known relatively stable compound of uranium? [Question ID = 687]

1.  $\text{UF}_6$  [Option ID = 2745]
2.  $\text{UO}_3$  [Option ID = 2748]
3.  $\text{UO}_2$  [Option ID = 2747]
4.  $\text{U}(\text{CH}_3)_2$  [Option ID = 2746]

Correct Answer :-

- $\text{U}(\text{CH}_3)_2$  [Option ID = 2746]

91) Which of the following compounds exist in stereoisomeric form? [Question ID = 681]

1.  $[\text{Pt}(\text{NH}_3)_3\text{Cl}]^+$  [Option ID = 2721]
2.  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  [Option ID = 2724]
3.  $[\text{Pt}(\text{NH}_3)\text{Cl}_3]^-$  [Option ID = 2722]
4.  $[\text{PtCl}_4]^{2-}$  [Option ID = 2723]

Correct Answer :-

- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  [Option ID = 2724]

92) Which of the following statement is not true? [Question ID = 745]

1. Methane is a spherical top molecule [Option ID = 2978]
2. Chloroform is a symmetric top molecule [Option ID = 2980]
3. Vinyl chloride is a symmetric top molecule [Option ID = 2979]
4. Water is an asymmetric top molecule [Option ID = 2977]

Correct Answer :-

- Vinyl chloride is a symmetric top molecule [Option ID = 2979]

93) Which of the following is a n-type semiconductor? [Question ID = 696]

1. Silicon carbide [Option ID = 2784]
2. Silicon [Option ID = 2781]
3. Arsenic doped silicon [Option ID = 2783]
4. Gallium doped silicon [Option ID = 2782]

Correct Answer :-

- Arsenic doped silicon [Option ID = 2783]

94) Which of the statement is not true? [Question ID = 746]

1. Franck Condon Principle states that during electronic transition the internuclear distance of a molecule does not change [Option ID = 2983]
2. The intensity of a fundamental vibrational transition is higher than that of a first overtone transition. [Option ID = 2984]
3. Morse equation represents the energy expression of a simple harmonic oscillator [Option ID = 2982]
4. The energy spacing between various vibrational levels are the same in a simple harmonic oscillator [Option ID = 2981]

Correct Answer :-

- Franck Condon Principle states that during electronic transition the internuclear distance of a molecule does not change [Option ID = 2983]



95) Which of the statement is true? [Question ID = 752]

1. The mean ionic activity coefficients of aqueous NaCl solution and aqueous KBr solution, both at low concentrations, are independent of their respective ionic strengths [Option ID = 3008]
2. The mean ionic activity coefficient of aqueous NaCl solution at low concentration decreases with increase in its ionic strength [Option ID = 3006]
3. The mean ionic activity coefficients of aqueous NaCl solution and aqueous KBr solution, both at low concentrations, vary differently upon increase of their respective ionic strengths [Option ID = 3007]
4. The mean ionic activity coefficient of aqueous NaCl solution at low concentration increases with increase in its ionic strength [Option ID = 3005]

Correct Answer :-

- The mean ionic activity coefficient of aqueous NaCl solution at low concentration decreases with increase in its ionic strength [Option ID = 3006]

96) The highest temperature that can be achieved due to a single normal mode of vibration in a solid crystal is known as: [Question ID = 757]

1. Debye Temperature [Option ID = 3026]
2. Theta Temperature [Option ID = 3027]
3. Curie Temperature [Option ID = 3025]
4. Flory Temperature [Option ID = 3028]

Correct Answer :-

- Debye Temperature [Option ID = 3026]

97) Which is not a scientific site? [Question ID = 691]

1. Research Gate [Option ID = 2763]
2. Scopus [Option ID = 2761]
3. Web of Science [Option ID = 2762]
4. Google Plus [Option ID = 2764]

Correct Answer :-

- Google Plus [Option ID = 2764]

98) According to the Michaelis Menten equation for unimolecular reactions: [Question ID = 751]

1. The rate is first order at low pressure, but becomes zero order at high pressure [Option ID = 3003]
2. The rate is zero order at both low and high pressures [Option ID = 3002]
3. The rate is zero order at low pressure, but becomes first order at high pressure [Option ID = 3004]
4. The rate is first order at both low and high pressures [Option ID = 3001]

Correct Answer :-

- The rate is first order at low pressure, but becomes zero order at high pressure [Option ID = 3003]

99) The +1 oxidation state is more stable than +3 oxidation state for which of the following Group 13 element [Question ID = 694]

1. In [Option ID = 2775]
2. B [Option ID = 2773]
3. Al [Option ID = 2774]
4. Tl [Option ID = 2776]

Correct Answer :-

- Tl [Option ID = 2776]

100) In how many ways can 10 distinguishable particles be placed in 3 boxes, so that there are 3 particles in first box, 5 in second and 2 in third? [Question ID = 737]

1. None of these [Option ID = 2948]
2. 1520 ways [Option ID = 2946]
3. 3260 ways [Option ID = 2947]
4. 2520 ways [Option ID = 2945]

Correct Answer :-

- 2520 ways [Option ID = 2945]

