### www.FirstRanker.com

## QUESTION BANK IN PHYSICS (B.TECH FIRST YEAR)

# SUPERCONDUCTIVITY

- Explain Meissner effect, type-I and type-II superconductors. (4) {JUN 15 [GNE]}
- What is isotope effect? (2) {JUN 15 [PTU]}
- 3. Give a brief account of BCS theory of superconductivity. (3) {JUN 15 [PTU]}
- Give a brief account of occurrence of superconductivity using BCS theory. (3) {JUN 15 [PTU]}
- What is the wavelength of an electromagnetic photon, which can break a Cooper pair in a material having critical temperature of 4K? (2) {Dec 14 [GNE]}
- Derive London's equations and give their significance.(4) {Dec 14 [GME]}
- Derive London's equations and show that these equations can account for perfect diamagnetism property of an ideal superconductor. (4) {JUN 14 [GNE]}
- 8. What is the effect of magnetic field on superconductivity? Given a type-I superconductor with  $T_C = 7K$  and slope  $\frac{dH_C}{dt} = -5 \times 10^{-4} A/m$  at  $T_C$ . Estimate its critical field at 6K. (4) {JUN 14 [GNE]}
- What do you mean by Meissner effect? (2) {JUN 14 PPU]} \
- A superconducting state behaves according to which type of magnetic material in presence of applied magnetic field having magnitude less than critical value. (2) {JUN 14 [PTU]}
- 11. What do you understand by superconducting state? Under what conditions one can achieve it? (4) {Dec 2013 [PTU]]
- Why are type I superconductors poor current carrying conductors? (2) {Dec 2013 [PTU]}
- What is Meissner effect? (2) {Dec 2013 [PTU]}
- Derive London equation and discuss how its solution led to Meissner effect. (4)
  [Dec 2013 [PTU]]
- 15. The penetration depth of mercury at 3.5K is about 750A. What will be the penetration depth at 0K, if the critical temperature for mercury is 4.2K? (4) {Dec 2013 [PTU]}
- Enumerate the factors affecting superconductivity. (2) {Dec 2013 [GNE]}
- 17. The critical magnetic field for a superconductor at absolute zero is 9×10<sup>4</sup> Am<sup>-1</sup> and at 6K is 5×10<sup>4</sup> Am<sup>-1</sup>. Find the critical temperature and energy required to break Cooper pair at absolute zero. (4) {Dec 2013 [GNE]}
- Derive London's equations and hence explain Meissner's effect and flux penetration. (4) {Dec 2013 [GNE]}
- What is Cooper pair? (2) {Jun 2013 [GNE]}
- Deduce London equations and define London penetration depth. (4) {Jun 2013 [GNE]}
- What is Meissner Effect? Explain type-I and type-II superconductors. (4) {Dec 2012 [GNE]}
- 22. For a specimen of V<sub>3</sub>Ga, the critical fields are 1·4×10<sup>5</sup> A/m & 4·2×10<sup>5</sup> A/m at 14K and 13K respectively. Calculate the transition temperature and critical fields at 0K and 4·2K. (2) {Dec 2012 [GNE]}
- Outline some experimental facts about superconductivity. (4) {Dec 2012}



#### www.FirstRanker.com

## QUESTION BANK IN PHYSICS (B.TECH FIRST YEAR)

- What arte type-II superconductors? (2) {June 2012}
- 25. What is the physical phenomenon behind superconductivity? How successful is this in today's context? (4) {June 2012}
- Elaborate the main features of BCS theory. (4) {June 2012}
- 27. What are important features of BCS theory? (2) {Dec 2011}
- 28. What is the physical mechanism behind Meissner Effect? (3) {Dec 2011}
- Discuss London's theory of superconductivity. (5) {Dec 2011}
- 30. What are the conditions for a material to be superconductor? (2) [June 2011]
- Explain BCS theory of superconductivity. (4) {June 2011}
- 32. What are London Equations? Find the expression for penetration depth of a superconductor. (4) {June 2011}
- 33. What is Meissner Effect? (2) {Dec 2010}
- Explain the difference between type-I and type-II superconductors. (3) {Dec 2010}
- 35. Give the salient features of BCS theory of superconductors. (3) {Dec 2010}
- Superconductors are perfectly diamagnetic. Explain. (2) [Dec 2010]
- What is Cooper pair? (2) {June 2010}
- Discuss the important differences between type-I and type-II superconductors with the help of example and plots of magnetization (M) Vs magnetic field (H).
   (3) {June 2010}
- What is Meissner Effect? Further explain the effect of magnetic field on the superconducting state. (3) {June 2010}
- 40. Define London Penetration depth and write its expression. (2) {June 2010}
- 41. Draw graphs for hard and soft superconductors. (2) {Dec 2009}
- Explain BCS theory of superconductivity. (5) {Dec 2009}
- Write down the relation between critical field and critical temperature in superconductors. (2) [June 2009]
- 44. Plot the graphs for type-I and type-II superconductors. (2) {June 2009}
- Derive & explain the London equations and calculate the expression for the Penetration Depth. (8) (June 2009)
- 46. What do you mean by field penetration in the superconductors? (2) {Dec 2008}
- 47. What do you mean by coherence length in context with superconductors? (2) {Dec 2008}
- 48. Define Levitation effect and explain the various factors that can destroy the superconductivity. (4) {Dec 2008}
- Explain the BCS theory of superconductivity. (4)
- Why superconductors are perfectly diamagnetic in nature? (2) {May 2008}
- 51. What is critical field? Write down the expression for H<sub>c</sub> and differentiate between type-I and type-II superconductors. (4) {May 2008}
- Derive first London's equation and give its physical significance. (4) {May 2008}
- 53. State Meissner effect of superconductivity. (2) {Dec 2007}
- 54. What is London's penetration depth? How does it vary with temperature? (4) {Dec 2007}
- 55. Define Cooper Pair. Calculate the wavelength of a photon, which will be required to break a Cooper Pair in a superconductor (Zr) for which T<sub>C</sub> = 0·56K. (4) {Dec 2007}





### QUESTION BANK IN PHYSICS (B.TECH FIRST YEAR)

- Write the formula for variation of magnetic field intensity with temperature. (2)
- 57. What is superconductivity? What are the differences between type-I and type-II superconductors? A type-I superconductor with  $T_c = 7K$  has

slope 
$$\frac{dB_C}{dT} = -25mTK^{-1}$$
 at  $T_C$ . Estimate its critical field at 6K. (8) {May 2007}

- 58. What are Cooper Pairs? (2) {Dec 2006}
- What are type I and type II superconductors? Explain. (3) {Dec 2006}
- 60. For Hg (mercury), the critical temperature at which the superconductivity ensues with zero applied magnetic fields is 4.15K. The critical applied magnetic field at which superconductivity will not take place at any temperature is 0.041T. Find the applied magnetic field that will stop the superconductivity at  $2 \cdot 2K$ . (3) Dec
- What is Meissner Effect? (2) {May 2006}
- What do you understand by type-I and type-II superconductors? (6) {May 2006}
- Discuss London's theory of superconductivity. (2) May 2006
- 64. What are Cooper pairs? (2) {Dec2005}
- 65. What do you understand by type-I and type-II superconductors? Give BCS theory of superconductivity. (8)
- 66. What is the effect of magnetic field on superconductivity? (2) {May 2005}
- 67. What is Meissner Effect? Show how London equations lead to this effect. (1,4) {May 2005}
- 68. A type-I superconductor with  $T_c = 7K$  has slope  $\frac{dB_c}{dT} = -25mTK^{-1}$  at  $T_c$ . Estimate its critical field at 6K. (3) {May 2005}
- What is Meissner effect? (2) {Dec 2004}
- 70. What is superconductivity? What are the differences between type-I and type-II

superconductors? A type-I superconductor with 
$$T_C = 7K$$
 has slope  $\frac{dB_C}{dT} = -25mTK^{-1}$  at  $T_C$ . Estimate its critical field at 6K. (2,3,3) {Dec 2004}

- 71. Metals, which are very good conductors at normal temperatures do not show superconducting behaviour. Why? (2) {May 2004}
- Distinguish between type-I and type-II superconductors. Briefly discuss the BCS theory of superconductivity. (5) {May 2004}
- State and explain Meissner Effect. How do London equations account for this effect? (3) {May 2004}
- 74. What are Cooper pairs? (2) {Dec2003}

