

KCM 8406

Second Year MBBS Examination

II MBBS Pharmacology Paper 2

Time: 2 hours**Date: 01-04-2015****Max Marks: 40**

Instructions:

1. Answer to the points.
2. Figure to the right indicates marks.
3. Use separate answer books for each section.
4. Draw diagrams wherever necessary.
5. Write legibly.

Section 1

1.MCQ (10)

- a. Drug used for the treatment of antibiotic induced diarrhea is (a) Tetracycline (b) Chloramphenicol (c) Erythromycin (d) Metronidazole

- b. Which of the following drugs is an example of long acting beta agonist (LABA)? (a) Terbutaline (b) Adrenaline (c) Salmeterol (d) Salbutamol
- c. Drug of choice for warfarin overdose is (a) Protamine sulphate (b) Naloxone (c) Physostigmine (d) Vitamin,
- d. Which of the followings is a long acting insulin? (a) Insulin lispro (b) Insulin detemir (c) Insulin aspart (d) Insulin glilisin
- e. All of the + following are side effect of of glucocorticoid EXCEPT (a) Hypoglycemia (b) Delayed healing of wounds (c) Cataract (d) Myopathy
- f. Which of the followings is a penicillinase resistant penicillins? (a) Ampiciln (b) Cloxacilin (c) Amoxidiltin (d) Bacamplillin
- g. Rifampicin is indicated in all of the following conditions EXCEPT- (a) Tuberculosis (b) Leprosy (c) Lepra reaction (d) Brucellosis
- h. Most potent drug for treatment of leprosy is (a) Dapsone (b) Rifampicin (c) Clofazimine (d) Minocycline
- i. The most important dose limiting adverse

effect of Colchicine is (a) Sedation (b) Kidney disease (c) Diarrhea (d) Muscle paralysis

- j. Which drug combination is NOT suitable for the treatment of HIV-AIDS? (a) Zidovudine + Lamivudine + Nevirapine (b) Zidovudine Stavudine + Efavirenz (c) Zidovudine H₂amivudine + Efavirenz (d) Zidovudine + Tenofovir Lamivudine

Section 2

2. Discuss (Any Two of the Three) (10)

- a. Pharmacotherapy of asthma
- b. Treatment of diabetic ketoacidosis
- c. Pharmacotherapy of leprosy.

3. Write Short notes (Any four of the five) (10)

- a. Albendazole
- b. Oral iron Preparations
- c. Low Molecular Weight Heparin (LMWH)
- d. Adverse effects of antipsychotics
- e. Metronidazole.

4. Discuss (Any two of the Three) (10)

- a. Antiplatelet drugs
- b. Drugs used in the treatment of osteoporosis
- c. Drugs used in the treatment of UTI (Urinary tract infection)

www.FirstRanker.com