


ROUTES OF DRUG ADMINISTRATION



SYSTEMIC ROUTES

[DRUG IS ABSORBED IN BLOOD STREAM & DISTRIBUTED ALL OVER, INCLUDING SITE OF ACTION THROUGH CIRCULATION]

① ORAL → OLDEST & COMMONEST MODE.

→ BOTH SOLID & LIQUID DOSAGE FORMS CAN BE GIVEN ORALLY.

ADVANTAGES :

- SAFER ✓
- MORE CONVENIENT ✓
- DOES NOT NEED ASSISTANCE ✓
- NON-INVASIVE ✓
- OFTEN PAINLESS ✓
- CHEAPER ✓

LIMITATIONS : • ACTION IS SLOW ∴ NOT SUITABLE IN EMERGENCIES.

• MAY CAUSE NAUSEA & VOMITING

• CAN'T BE USED FOR UNCOOPERATIVE / UNCONSCIOUS / VOMITING PATIENT

• ABSORPTION OF DRUG IS VARIABLE ✓
(e.g. STREPTOMYCIN IS NOT ABSORBED)

[• SOME ARE DESTROYED BY DIGESTIVE JUICES (e.g. INSULIN) OR IN LIVER (e.g. GTN; TESTOSTERONE)

FIRST PASS METABOLISM

IS ~~WHICH~~ OCCURS.

∴ BIOAVAILABILITY VARIES.

② RECTAL → CERTAIN IRRITANT & UNPLEASANT DRUGS CAN BE PUT INTO RECTUM AS SUPPOSITORIES OR RETENTION ENEMA FOR SYSTEMIC EFFECT.

→ CAN BE USED IN PATIENT HAVING RECURRENT VOMITING OR IS UNCONSCIOUS.

→ DRUG ABSORBED INTO EXTERNAL HAEMORRHOIDAL VEINS (about 50%) BYPASS LIVER BUT NOT THAT ABSORBED INTO INTERNAL HAEMORRHOIDAL VEINS.

→ DIAZEPAM SOLUTION & PARACETAMOL SUPPOSITORY ARE RAPIDLY ABSORBED FROM RECTUM IN CHILDREN

LIMITATIONS: • ABSORPTION IS SLOW & IRREGULAR ;
OFTEN UNPREDICTABLE

Essentials of Medical Science

• INCONVENIENT & EMBARRASSING ✓

• RECTAL INFLAMMATION FROM IRRITANT DRUGS ✓

→ DIAZEPAM ✓
INDOMETHACIN ✓
PARACETAMOL ✓
ERGOTAMINE ✓
etc. } - Can be given
RECTALLY.

~~NON-ORAL TABLET~~
~~PARENTERAL ROUTES~~

③ SUBLINGUAL (BUCCAL) → TABLET / PELLET CONTAINING DRUG IS PLACED UNDER TONGUE OR CRUSHED IN THE MOUTH & SPREAD OVER BUCCAL MUCOSA

- ONLY LIPID SOLUBLE & NON-IRRITATING DRUGS CAN BE ADMINISTERED BY THIS ROUTE.
- ABSORPTION IS RELATIVELY RAPID. ✓
- ONE CAN SPIT THE DRUG AFTER DESIRED EFFECT IS OBTAINED. ✓

ADVANTAGE: • LIVER IS BYPASSED ✓

∴ DRUGS WITH HIGH FIRST PASS METABOLISM CAN BE ABSORBED DIRECTLY INTO SYSTEMIC CIRCULATION]

EXAMPLES: • GTN ✓

• BUPRENORPHINE ✓

• DESAMINO-OXYTOCIN ✓

NON-INJECTABLE PARENTERAL DRUGS

① NASAL → MUCOUS MEMBRANE OF NOSE CAN READILY ABSORB MANY DRUGS.

→ DIGESTIVE JUICES & LIVER ARE BYPASSED

ONLY CERTAIN DRUGS CAN BE USED BY THIS ROUTE.

Examples : • GnRH AGONISTS
• CALCITONIN ✓
• DESMOPRESSIN ✓

Essentials of Medical Science

② INHALATION → VOLATILE LIQUIDS & GASES ARE GIVEN BY THIS ROUTE FOR SYSTEMIC ACTION

→ ABSORPTION TAKES PLACE FROM VAST SURFACE OF ALVEOLI

∴ ACTION IS VERY RAPID.

→ [WHEN ADMINISTRATION IS DISCONTINUED
DRUG DIFFUSES BACK & RAPIDLY ELIMINATED
IN EXPIRED AIR.

→ IRRITANT VAPOURS CAUSES INFLAMMATION OF
RESPIRATORY TRACT & INCREASE SECRETION

→ EXAMPLE : GENERAL ANAESTHETICS.

③ CUTANEOUS → HIGHLY LIPID SOLUBLE DRUGS GIVEN BY THIS ROUTE

- DRUG IS APPLIED OVER SKIN FOR SLOW & PROLONGED ABSORPTION
- LIVER IS BYPASSED. =
- ABSORPTION OF DRUG CAN BE ENHANCED BY RUBBING THE PREPARATION. =

TRANS DERMAL THERAPUTIC SYSTEM (TTS) : //

Essentials of Medical Science

- I. DRUG IS ADMINISTERED IN THE FORM OF A PATCH OR OINTMENT THAT DELIVERS THE DRUG INTO CIRCULATION FOR SYSTEMIC EFFECT.
- II. DRUG IS DELIVERED AT A CONSTANT RATE INTO SYSTEMIC CIRCULATION VIA STRATUM CORNEUM.
- III. LAYERS :
 - BACKING FILM ①
 - DRUG RESERVOIR ②
 - RATE CONTROLLING MICROPORE MEMBRANE ③
 - ④ • ADHESIVE LAYER [WITH PRIMING DOSE]
 - ⑤ • OUTER LAYER [PEELED OFF BEFORE APPLYING ON SKIN]

IV. DRUG IS DELIVERED AT SKIN SURFACE BY DIFFUSION FOR PERCUTANEOUS ABSORPTION INTO CIRCULATION.

V. MICROPORE MEMBRANE IS SUCH THAT RATE OF DRUG DELIVERY TO SKIN SURFACE IS LESS THAN SLOWEST RATE OF ABSORPTION FROM SKIN.

∴ DRUG IS DELIVERED AT A CONSTANT & PREDICTABLE RATE IRRESPECTIVE OF SITE OF APPLICATION.

VI. COMMON SITES OF APPLICATION: CHEST, ABDOMEN, UPPER ARM, LOWER BACK, BUTTOCK OR MASTOID REGION

VII. EXAMPLES: ✓ SCOPOLAMINE PATCH FOR SIALORRHOEA & MOTION SICKNESS.
(HYOSCINE)

✓ GTN PATCH FOR PROPHYLAXIS OF ANGINA

✓ OESTROGEN PATCH FOR HRT.


✓ CLONIDINE PATCH FOR HYPERTENSION


VIII. ADVANTAGES: ✓ BIOAVAILABILITY IS 100%.
✓ QUICK ONSET OF ACTION
LARGE

- VIII. ADVANTAGES :
- SELF-ADMINISTRATION POSSIBLE ✓
 - PATIENT COMPLIANCE IS BETTER ✓
 - SYSTEMIC SIDE EFFECTS ARE LESS ✓
 - FIRST PASS METABOLISM IS BYPASSED ✓
 - DURATION OF ACTION IS PROLONGED ✓
 - PROVIDES A CONSTANT PLASMA CONCENTRATION OF DRUG ✓


- IX. DISADVANTAGES :
- EXPENSIVE ✓
 - LOCAL IRRITATION MAY CAUSE ITCHING ✓
 - PATCH MAY FALL OFF UNNOTICED ✓

Essentials of Medical Science





INJECTABLE PARENTERAL ROUTES



① SUBCUTANEOUS (S.C) → DRUG IS INJECTED INTO THE SUB SUBCUTANEOUS TISSUE OF THIGH, ABDOMEN, ARM.

EXAMPLE - ADRENALINE ; INSULIN etc.

- ADVANTAGES:
- SELF ADMINISTRATION IS POSSIBLE ✓
 - DEPOT PREPARATIONS CAN BE INSERTED INTO SUBCUTANEOUS TISSUE FOR PROLONGED ACTION. ✓
 - eg. NORPLANT FOR CONTRACEPTION. ✓

- DISADVANTAGES:
- IRRITANT DRUGS CAN NOT BE INJECTED
 - DRUG ABSORPTION IS SLOW
 - ∴ CAN'T BE GIVEN IN EMERGENCY.

② INTRADERMAL → DRUG IS INJECTED INTO LAYERS OF SKIN RAISING A BLEB. ✓

→ PAINFUL & A SMALL AMOUNT OF DRUG CAN BE ADMINISTERED. ✓

Example, BCG VACCINATION. ✓
&
DRUG SENSITIVITY TESTS. ✓

③ INTRAVENOUS (i/v) → DRUG IS INJECTED & AS A BOLUS OR INFUSED SLOWLY OVER HOURS INTO A SUPERFICIAL VEIN. ✓

→ DRUG REACHES DIRECTLY INTO BLOOD STREAM & EFFECTS ARE PRODUCED IMMEDIATELY. ✓
∴ USED IN EMERGENCY. ✓

→ ADVANTAGES :

- 100% BIOAVAILABILITY. ✓
- HIGHLY IRRITANT DRUGS CAN BE ADMINISTERED. ✓
∴ THEY GET DILUTED IN BLOOD. ✓
- QUICK ONSET OF ACTION. ✓
- LARGE VOLUME OF FLUID CAN BE ADMINISTERED. ✓
Eg. i/v FLUIDS IN SEVERE DEHYDRATION. ✓
- CONSTANT PLASMA LEVEL OF DRUG CAN BE MAINTAINED BY i/v INFUSION. ✓

→ DISADVANTAGES :

- LOCAL IRRITATION CAN CAUSE THROMBOPHLEBITIS. ✓

- SELF ADMINISTRATION NOT POSSIBLE USUALLY. ✓
- DEPOT PREPARATIONS CAN NOT BE GIVEN. ✓
- EXTRAVASATION OF SOME DRUGS (eg. Adrenaline) CAN CAUSE INJURY, NECROSIS & SLOUGHING OF TISSUES. ✓

→ PRECAUTIONS :

- DRUG SHOULD BE INJECTED SLOWLY. ✓
- BEFORE INJECTING, MAKE SURE THAT TIP OF NEEDLE IS IN VEIN. ✓

④ INTRAMUSCULAR (i/m) → DRUGS ARE INJECTED INTO
LARGE MUSCLES.

Eg. DELTOID, GLUTEUS MAXIMUS &
VASTUS LATERALIS.

→ ADVANTAGES : • MILD IRRITANTS CAN BE INJECTED ✓
∴ MUSCLE IS LESS RICHLY SUPPLIED
WITH SENSORY NERVES.

Essentials of Medical Science

- LESS PAINFUL ✓
- DEPOT INJECTIONS CAN BE GIVEN. ✓

→ DISADVANTAGES : • ABCESS FORMATION CAN OCCUR.
• SELF ADMINISTRATION NOT POSSIBLE
∴ DEEP PENETRATION IS NEEDED ✓
• SHOULD BE AVOIDED IN ANTICOAGULANT
TREATED PATIENTS ∴ IT CAN
PRODUCE LOCAL HEMATOMA.

Essentials of Medical Science

