

B.Tech II Year I Semester (R13) Supplementary Examinations June 2017

MATHEMATICS – II

(Common to CE and ME)

Time: 3 hours

Max. Marks: 70

PART – A

(Compulsory Question)

1 Answer the following: (10 X 02 = 20 Marks)

(a) What is the symmetric matrix? Give proper example.

(b) Determine the rank of
$$\begin{pmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 0 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & -2 & 0 \end{pmatrix}$$

(c) What is the formula for Newton's backward interpolation formula?

(d) Use the method of false position to find the root of the equation $x^3 - 18 = 0$, given it lies between 2 and 3. Write down the procedure.

(e) What is the formula of Simpson's 1/3 rule?

(f) Solve by Taylor's series method the equation $\frac{dy}{dx} = \log(xy)$ for $y(x)$, given $y(1) = 2$.

(g) Write Linear Property of Fourier transform.

(h) What is the formula of Fourier cosine transform?

(i) Derive a partial differential equation by eliminating the arbitrary function from the relation:

$$2z = \frac{x^2}{a^2} + \frac{y^2}{b^2}$$

(j) What is the One Dimensional Wave Equation (Vibration of a stretched string)?

PART – B

(Answer all five units, 5 X 10 = 50 Marks)

UNIT – I

2 Find the characteristic equation of the matrix $A = \begin{bmatrix} 4 & 3 & 1 \\ 2 & 1 & -2 \\ 1 & 2 & 1 \end{bmatrix}$ hence find A^{-1} .

OR

3 Find a matrix P which transform the matrix $A = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix}$ to diagonal form.

UNIT – II

4 Find the real root of the equation $x^4 - x - 9 = 0$ by Newton-Raphson method, correct to three places of decimal.

OR

5 Given the values:

x	5	7	11	13	17
f(x)	150	392	1492	2366	5202

Evaluate $f(9)$ use Lagrange's formula.

Contd. in page 2

UNIT – III

6 Use Simpson's $1/3^{\text{rd}}$ rule to find $\int_0^{0.6} e^{-x^2} dx$, by taking seven ordinates.

OR

7 Using Runge-Kutta Method of fourth order, find an approximate value of y when $x = 0.2$, given that $\frac{dy}{dx} = \frac{y^2 - x^2}{y^2 + x^2}$ with $y(0) = 1$.

UNIT – IV

8 Obtain Fourier series expansion for $f(x)$ defined as follows:

$$f(x) = x + \pi, \quad 0 \leq x \leq \pi$$

$$f(x) = -\pi - x, \quad -\pi \leq x < 0$$

OR

9 Find Fourier transform of $f(x) = \begin{cases} 1 & \text{for } |x| < 1 \\ 0 & \text{for } |x| > 1 \end{cases}$ and hence find $\int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$

UNIT – V

10 Form the partial differential equation (by eliminating the arbitrary constants a, b) from:

$$(x-a)^2 + (y-b)^2 + z^2 = k^2$$

OR

11 Using the method of separation of variables, solve $\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u$ where $u(x, 0) = 6e^{-3x}$.
