

Code No: 09A1BS04

R09**Set No. 2****I B.Tech Examinations, December 2010****MATHEMATICAL METHODS****Common to BME, IT, ICE, E.COMP.E, ETM, EIE, CSE, ECE, EEE****Time: 3 hours****Max Marks: 75****Answer any FIVE Questions****All Questions carry equal marks**

1. Verify Cayley Hamilton theorem and find the inverse of $\begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$. [15]

2. (a) Find a real root of the equation, $x \sin x + \cos x = 0$ using regula falsi method.
 (b) Find $y(32)$ if $y(10)=35.3$, $y(15) = 32.$, $y(20) = 29.2$, $y(25) = 26.1$, $y(30) = 23.2$, $y(35) = 20.5$ using Newton's forward interpolation formula. [8+7]

3. (a) Compute the first and second derivatives at $x=2.03$ of the following table:

x	1.96	1.98	2.00	2.02	2.04
y	0.7825	0.7739	0.7651	0.7563	0.7473

- (b) From the following table of values of x and y find $\frac{dy}{dx}$ at $x=0.5$

x	0.35	0.40	0.45	0.5	0.55	0.6	0.65
y	1.521	1.506	1.488	1.467	1.444	1.418	1.389

[8+7]

4. (a) Find the maximum and minimum values of $f = 3x^2 + 5y^2 + 3z^2 - 2xy + 2zx - 2yz$ subject to $x + y + z = 1$. Also find the point at which the maximum and minimum exists.

- (b) Find the nature of the quadratic form $10x^2 + 2y^2 + 5z^2 - 4xy - 10zx + 6yz$. [8+7]

5. Find $y(0.5)$, $y(1)$ and $y(1.5)$ given that $\frac{dy}{dx} = 4 - 2x$ and $y(0) = 2$ with $h = 0.5$ using modified Euler's method. [15]

6. If $f(x) = x$ for $0 < x < \frac{\pi}{2}$
 $= \pi - x$ for $\frac{\pi}{2} < x < \pi$. then prove that

(a) $f(x) = \frac{4}{\pi} \left[\sin x - \frac{1}{3^2} \sin 3x + \frac{1}{5^2} \sin 5x - \dots \right]$.

(b) $f(x) = \frac{\pi}{4} - \frac{2}{\pi} \left[\frac{1}{1^2} \cos 2x + \frac{1}{3^2} \cos 6x + \frac{1}{5^2} \cos 10x + \dots \right]$. [8+7]

7. (a) Solve $\frac{p}{x^2} + \frac{q}{y^2} = 1$.

(b) Solve $p^2 + q^2 = z^2(x^2 + y^2)$. [7+8]

8. (a) Find the Rank of the Matrix, by reducing it to the normal form. $\begin{bmatrix} 2 & 1 & -3 & -6 \\ 2 & -3 & 1 & 2 \\ 1 & 1 & 1 & 2 \end{bmatrix}$

Code No: 09A1BS04

R09**Set No. 2**

- (b) Find whether the following system of equations are consistent. If so solve them. $5x + 3y + 7z = 0$, $3x + 26y + 2z = 9$, $7x + 2y + 10z = 5$. [7+8]

FIRSTRANKER

Code No: 09A1BS04

R09**Set No. 4****I B.Tech Examinations, December 2010****MATHEMATICAL METHODS****Common to BME, IT, ICE, E.COMP.E, ETM, EIE, CSE, ECE, EEE****Time: 3 hours****Max Marks: 75****Answer any FIVE Questions****All Questions carry equal marks**

1. Find the eigen values and the corresponding eigen vectors of $\begin{bmatrix} 1 & 3 & 7 \\ 1 & 2 & 3 \\ 1 & 2 & 1 \end{bmatrix}$. [15]
2. Reduce the quadratic form to the canonical $3x^2 - 2y^2 - z^2 - 4xy + 8xz + 12yz$. [15]
3. (a) Solve $(x^3 + 3xy^2)p + (y^3 + 3x^2y)q = 2(x^2 + y^2)z$.
 (b) Solve $x(y^2 - z^2)p - y(z^2 + x^2)q = z(x^2 + y^2)$. [7+8]
4. (a) Find the Rank of the Matrix, by reducing it to the normal form. $\begin{bmatrix} 1 & -1 & 2 & 5 \\ 2 & 1 & 4 & 3 \\ 1 & -1 & -3 & 5 \end{bmatrix}$.
 (b) Solve the following tridiagonal system $3x - y = 5$, $x + 2y - 2z = 6$, $4y + 3z = 1$. [8+7]
5. Evaluate $\int_0^1 \frac{1}{1+x} dx$
 (a) By Trapezoidal rule and Simpson's $\frac{1}{3}$ rule.
 (b) Using Simpson's $\frac{3}{8}$ rule. [8+7]
6. If $f(x) = \begin{cases} 1 & \text{in } 0 < x < \Pi/2 \\ -1 & \text{in } \Pi/2 < x < \Pi \end{cases}$ Expand $f(x)$ in a series of cosines. [15]
7. Find $y(.1)$ and $y(.2)$ using Runge-Kutta fourth order formula given that $\frac{dy}{dx} = x^2 - y$ and $y(0) = 1$. [15]
8. (a) Find a real root of the equation $xe^x = \cos x$ by Newton Raphson method.
 (b) The amount A of a substance remaining in a reaction system after an interval of time t in a certain chemical experiment is given by the following data.
- | | | | | |
|---|------|------|-----|------|
| t | 2 | 5 | 8 | 14 |
| A | 94.8 | 87.9 | 813 | 68.7 |
- Find value of A at $t = 11$. [8+7]

Code No: 09A1BS04

R09**Set No. 1****I B.Tech Examinations, December 2010****MATHEMATICAL METHODS****Common to BME, IT, ICE, E.COMP.E, ETM, EIE, CSE, ECE, EEE****Time: 3 hours****Max Marks: 75****Answer any FIVE Questions****All Questions carry equal marks**

1. Form the partial differential equations

(a) $z = f(x-iz) + g(x-iz)$

(b) $z = y^2 + 2f\left(\frac{1}{x} + \log y\right)$

(c) $F(xy + z^2, x + y + z) = 0$. [5+5+5]

2. (a) Find a real root of the equation $e^x \sin x = 1$ using Newton Raphson method(b) Find $y(10)$, Given that $y(5) = 12$, $y(6) = 13$, $y(9) = 14$, $y(11) = 16$ using Lagrange's formula. [8+7]3. (a) Find the maximum and minimum values of $f = 2x^2 + 2y^2 + 2z^2 - 2xy + 2zx - 2yz$ subject to $x + y + z = 1$. Also find the point at which the maximum and minimum exists.(b) Find the nature of the quadratic form $3x^2 + 2y^2 + 3z^2 - 2xy - 2yz$. [8+7]4. Verify Cayley Hamilton theorem and find the inverse of $\begin{bmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{bmatrix}$. [15]5. (a) Find the Rank of the Matrix, by reducing it to the normal form. $\begin{bmatrix} 2 & 1 & 3 & 4 \\ 0 & 3 & 4 & 1 \\ 2 & 3 & 7 & 5 \\ 2 & 5 & 11 & 6 \end{bmatrix}$ (b) Solve the following equations by expressing the coefficient matrix as a product of a lower triangular and upper triangular matrices. $x + y - z = 5$, $2x + y + 2z = 5$, $3x + 2y - 4z = 7$. [7+8]6. Given $y' = x + \sin y$ and $y(0) = 1$ compute $y(0.2)$ and $y(0.4)$ with $h = 0.2$ using Euler's modified method. [15]7. (a) Find the half-range sine series of $f(x) = 1$ in $[0, l]$.(b) Find the half-range cosine and sine series for $f(x) = x$ in $(0, l)$. [7+8]8. Evaluate $\int_0^6 \frac{1}{(1+x)} dx$ by using(a) Simpson's $\frac{1}{3}$ Rule.

Code No: 09A1BS04

R09

Set No. 1

(b) Trapezoidal Rule.

(c) Simpson's $\frac{3}{8}$ Rule.

[5+5+5]

FIRSTRANKER

Code No: 09A1BS04

R09**Set No. 3****I B.Tech Examinations, December 2010****MATHEMATICAL METHODS****Common to BME, IT, ICE, E.COMP.E, ETM, EIE, CSE, ECE, EEE****Time: 3 hours****Max Marks: 75****Answer any FIVE Questions****All Questions carry equal marks**

1. (a) Find the Rank of the Matrix ,by reducing it to the normal form.

$$\begin{bmatrix} 2 & 3 & -1 & -1 \\ 1 & -1 & -2 & -3 \\ 3 & -1 & 3 & -2 \\ 6 & 3 & 0 & -7 \end{bmatrix}$$

- (b) Solve the following tridiagonal system

$$2x - y = 3, x + 4y + 3z = 3, 2y + 4z = 6.$$

[8+7]

2. (a) Express
- $f(x) = \frac{\pi^2}{12} - \frac{x^2}{4}$
- as a Fourier in
- $-\pi < x < \pi$
- .

- (b) Find the Fourier Series to represent the function
- $f(x) = \sin x$
- in
- $-\pi < x < \pi$
- .

[8+7]

3. (a) Find the constants a and b by the method of least squares such that
- $y = ae^{bx}$

x	2	4	6	8	10
y	4.077	11.084	30.128	81.897	222.62

- (b) Find the curve of best fit of the type
- $y = ae^{bx}$
- to the following data by method of least squares.

x	1	5	7	9	12
y	10	15	12	15	21

[8+7]

4. Given
- $\frac{dy}{dx} = \frac{y-x}{y+x}$
- and
- $y(0)=1$
- . Compute
- $y(0.1)$
- in steps of 0.02 using Euler's modified method. [15]

5. Reduce the quadratic form to the canonical form
- $2x^2 + 2y^2 + 2z^2 - 2xy + 2zx - 2yz$
- . [15]

6. (a) Solve
- $2(z+px+qy)=p^2y$
- .

- (b) Solve
- $(p^2 + q^2)y=qz$
- .

[8+7]

7. Diagonalize the matrix
- $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$
- . [15]

8. (a) Find a real root of the equation ,
- $x^3 - 9x + 1 = 0$
- using regula falsi method.

- (b) Prove that
- $(E^{1/2} + E^{-1/2})(1 + \Delta)^{1/2} = 2 + \Delta$
- .

[8+7]
