Development of the Vascular System

VASCULOGENESIS formation of the blood vessels from blood island
 ANGIOGENESIS Sprouting of vessels from the preexisting vessels

Once the process of vasculogenesis establishes a primary vascular bed, which include dorsal aorta and cardinal veins, additional vasculature is added by angiogenesis



www.FirstRanker.com

ARETERIAL SYSTEM

PHARYNGEAL ARCHES





they are not present simultaneously all the time.



AORTIC^{TR}ARCHES^{TSTRanker.com}

- Are the arteries that arise from the aortic sac and supply the pharyngeal arches.
- The aortic sac contributes a branch to each new arch as it forms, thus giving rise to total of five pairs of arteries.



- The aortic arches terminate in the right and left dorsal aortae which which later fuse in the caudal region to form single vessel.
- Division of the truncus arteriosus by the aortopulmonary septum divides the outflow channel of the heart into the ventral aorta and the pulmonary artery.
- The aortic sac then forms right and left horns, which subsequently give rise to the brachiocephalic artery and the proximal segment of the arch of aorta respectively.



CHANGES IN AORTIC ARCHES



www.FirstRanker.com www.FirstRanker.com artery

OTHER CHANGES IN THE ARCH SYSTEM

- The dorsal aorta located between the entrance of the 3rd and 4th arches, known as the carotid duct, is obliterated.
- The right dorsal aorta disappears between the origin of the 7th intersegmental artery and the junction with the left dorsal aorta.



www.FirstRanker.com

- Cephalic folding, growth of the forebrain, and elongation of the neck cause the heart to to descend into the thoracic cavity.
- Carotid and brachiocephalic arteries elongate considerably.
- Origin of the left subclavian artery shifts higher up until it it comes close to the originf of the left common carotid artery.



- The course of the recurrent laryngeal nerve (6th pharyngeal arch) becomes different on the right and left sides.
- When the heart descends, they hook around the 6th aortic arches and then ascend again to the larynx, thus thus accounting for their recurrent course.



WWW.FirstRanker.com DERIVATIVES OF THE AORTIC ARCHES

1	Maxillary arteries		
2	Hyoid and stapedial art	eries	
∎ 3	Common carotid and fin internal carotid arteries	rst part	of the
4	Left side - Arch of the aorta from the left common carotid to the left subclavian arteries Right side - Right subclavian artery (proximal		
6	Left side - Left pulmona	arv arter	v and ductus
_ 0	arteriosus		y and daotao
	Right side - Right pulme	onary a	rtery

INTERSEGMENTAL ARTERIES

Approx. 30 segmental branches of dorsal aorta carry blood to somites and their derivatives.

Persisting segmental arteries —

- Intercostal
- Lumbar
- Common iliac
- Lateral sacral
- Median sacral

UNPAIRED BRANCHES FROM THE DORSAL AORTA

Caudal end of the dorsal aorta becomes the median sacral artery

The vitelline arteries, initially a number of paired vessels supplying the yolk sac, gradually fuse and forms arteries located in the dorsal mesentery of the gut.

In the adult they are represented by the celiac, superior mesenteric, and inferior mesenteric arteries. These vessels supply derivatives of the foregut, midgut, and hindgut respectively



PAIRED UMBILICAL ARTERIES

The **umbilical arteries**, initially paired ventral branches of the dorsal aorta. During the 4th week, each artery acquires a secondary connection with the dorsal branch of the aorta, the **common iliac artery**, and loses its original origin.

Carry deoxygenated blood to the placenta.

After birth:

- The proximal portions become the internal iliac artery and superior vesical artery.
- Distal parts degenerate and become the median umbilical ligaments.



www.FirstRanker.com



CORONARY ARTERIES

- Two sources
- angioblast
- > epicardium
- Neural crest cells
- Connection of coronary arteries to Aorta occurs by in growth of arterial endothelial cells into aorta



CONGENITAL ANOMALIES

1. PATENT DUCTUS ARTERIOSUS

is one of the most frequently occurring abnormalities of the great vessels, especially in premature infants. There is a large difference between aortic and pulmonary pressures and the increased blood flow through the ductus prevents prevents its closure.



2) COARCTATION OF AORTA

- Aortic lumen below the origin of the left sub-clavian artery is significantly narrowed.
- Is of two types:
 - Preductal : ductus arteriosus persists
 - Postductal : ductus arteriosus is usually obliterated





Coarctation of the Aorta



VENOUS SYSTEM

- At 5th week three pairs of major veins can be be seen
 - Vitelline or omphalomesenteric veins, carrying blood from the yolk sac to the sinus venosus
 - Umbilical veins, originating in the chorionic villi and carrying oxygenated blood to the embryo.
 - Cardinal veins, draining the body of the embryo proper.



www.FirstRanker.com

VITELLINE VEINS

- Before entering the sinus venosus, the vitelline veins form a plexus around the duodenum and pass through the septum transversum.
- The liver cords growing into the septum interrupt the course of veins and an extensive vascular network is formed known as the hepatic sinusoids.



- With reduction of left sinus horn, blood from the left side of the liver is rechanneled toward the right resulting in the enlargement of the right vitelline vein (right hepatocardiac channel).
- Ultimately, the right hepatocardiac channel forms the hepatocardiac hepatocardiac portion of the inferior vena cava.
- The proximal part of the left vitelline vein disappears.
- The anastomotic network around the duodenum develops into a single single vessel, the portal vein.
- The superior mesenteric vein, which drains the primary intestinal loop loop is derived from the right vitelline vein. The distal portion of the left vitelline vein also disappears.



Derivatives of Vitelline veins

Inferior Vena Cava
Hepatic veins
The portal vein
Superior Mesenteric Vein



UMBILICAL VEINS

- The proximal part of both umbilical veins and the remainder of the right umbilical vein disappear, so that the left vein is the only one to carry blood from the placenta to the liver.
- With increase of the placental circulation, a direct communication is formed between the left umbilical vein and the right hepatocardiac channel, the ductus venosus.
- This vessel bypasses the sinusoidal plexus of the liver.
- After birth, the left umbilical vein and the ductus venosuss are obliterated and form the ligamentum teres hepatic and ligamentum venosum, respectively.

CARDINAL VEINS

During the 4th week, the cardinal veins form the main venous drainage system of embryo.

This system consists of:

- Anterior cardinal veins which drain the cephalic part of the embryo
- Posterior cardinal veins which drain the remaining part of the body of the embryo.
- The anterior and posterior veins join before entering the sinus horn and form the short common cardinal cardinal veins.

ANTERIOR CARDINAL VEINS

 The anterior cardinal vein become connected by By ahlouique anastonio sisting the 8th week.
 Shunts blood from left to right.

Anastomosis between the Anterior Cardinal vein viewelops into left Briachjocephalic vein

Right anterior cardinal vein and the right common common cardinal vein and the SVC.



During the 5th to 7th week, a number of additional veins are formed.

- Subcardinal veins which mainly drain the kidneys

- Sacrocardinal veins which drain the lower extremities
- Supracardinal veins which drain the body wall by way of the intercostal veins, thus taking over the functions of the posterior posterior cardinal veins.



Posterior Cardinal vein



Major tributaries of the posterior cardinal veins
Subcardinal veins
Supracardinal veins
Sacrocardinal veins

Formation of the vena cava system is characterized by the appearance of anastomoses between left and right in such a manner that the blood from the left left is channeled to the right side.



- The anastomosis between the cardinal veins develops into the left brachiocephalic vein.
- Most of the blood from the left side of the head and the left upper extremity is is then channeled to the right.
- The terminal portion of the left posterior cardinal vein entering into the left left brachiocephalic vein is retained as a small vessel, the left superior intercostal vein.
- This vessel receives blood from the 2nd and 3rd intercostal spaces.
- The superior vena cava is formed by the right common cardinal vein and and the proximal portion of the right anterior cardinal vein.





Formation of Inferior vena cava

- The anastomosis between the subcardinal veins forms the left renal vein.
- When this communication has been established, the left subcardinal vein disappears and only its distal portion remains as the left gonadal vein.
- Hence the right subcardinal vein becomes the main drainage channel and develops into the renal segment of the inferior vena cava.



- The anastomosis between the sacrocardinal veins forms the left common iliac vein.
- The right sacrocardinal vein becomes the sacrocardinal segment of the inferior vena cava.
- When the renal segment of the inferior vena cava connects with the hepatic segment, which is derived from the right vitelline vein, the inferior ior vena cava is complete.
- It thus consists of a hepatic segment, renal segment and a sacrocardinal segment.

Azygos and Hemiazygos veins

- With obliteration of the major portion of the posterior cardinal veins, the supracardinal veins gain in importance.
- The 4th to 11th right intercostal veins empty into the right supracardinal vein, which together with a portion of the posterior cardinal vein, forms the azygos vein.

On the left, the 4th to 7th intercostal veins enter into the left supracardinal vein, and the left supracrdinal vein empties into the azygos vein and is then known as the itemizzogos vein

www.FirstRanker.con

CONGENITAL ANOMALIES

DOUBLE INFERIOR VENA CAVA

ABSENCE OF INFERIOR VENA CAVA



www.FirstRanker.com

LEFT SUPERIOR VENA CAVA DOUBLE SUPERIOR VENA CAVA



www.FirstRanker.com

CIRCULATORY CHANGES AT BIRTH

FETAL CIRCULATION



BEFORE BIRTH



1. Closure of the umbilical arteries

 It is accomplished by contraction of smooth musculature in their walls and is probably caused by thermal and mechanical stimuli and change in oxygen tension. The distal parts of the umbilical arteries form the medial umbilical ligament, while the proximal portions remain remain open as the superior vesical arteries.

2. Closure of the umbilical vein and ductus venosus

 This occurs shortly after that of umbilical arteries. After obliteration, on umbilical vein forms the ligamentum teres hepatic in the lower margin margin of the falciform ligament. The obliterated ductus venosus forms the ligamentum venosum.

3. Closure of the ductus arteriosus

 It occurs by contraction of its muscular wall almost immediately after free birth and is mediated by bradykinin, released from the lungs during initial inflation. The obliterated ductus arteriosus forms the ligamentum arteriosum.

4. Closure of the foramen ovale

 This is caused by an increased pressure in the left atrium combined with data as a pressure on the right of the two first breath the sentum septum p primum is pressed against the septum secondum. Constant apposition gradually leads to fusion of the two septa in about a year.

AFTER BIRTH



LYMPHATIC SYSTEM

The lymphatic system begins its development later than the cardiovascular cardiovascular system and does note 5th appear until the 5th week of gestation.

The origin of lymphatic vessels is not clear, but they may form from mesenchyme in situ or may arise ass sad like outgrowths from the endothelium of veins.

As a result, six primary lymph sacs are formed:

- Two jugular, at the junction of the subclavian and anterior anterior anterior
- Two iliac, at the junction of the iliac and posterior cardinal veins
- One retroperitoneal, near the root of the mesentery
- The cisterna chyli, dorsal to the

retroperitoneal sac.



Numerous channels connect the sacs with each other and also drain lymph from the limbs, body wall, and head and neck. Two main channels, the right and left thoracic ducts, join the jugular sacs with the cisterna chyli and soon an anastomosis forms between these ducts. The thoracic duct then develops from the distal portion of the right thoracic duct, the

www.FirstRanker.com www.FirstRanker.com



www.FirstRanker.com

NINE WEEKS

FETAL PERIOD



The right lymphatic duct is derived from the cranial portion of the right thoracic thoracic duct.

Both ducts maintain their original connections with the venous system and empty into the junction of the internal jugular and subclavian veins.