#### Column-I Column-II a. Pistils fused Gametogenesis together b. Formation of

www.FirstRanker.com

(ii) Pistillate

c. Hyphae of higher Syncarpous Ascomycetes

d. Unisexual female flower

(iv) Dikaryotic

#### Codes:

gametes

а b d c (1)(i) (ii) (iv) (iii) (iii) (i) (iv) (ii) (3) (iv) (iii) (ii) (i) (4) (ii) (i) (iv) (iii)

3. In majority of angiosperms

(1) reduction division occurs the megaspore mother cells

(2) a small central cell is present in the embryo sac

(3) egg has a filiform apparatus

(4) there are numerous antipodal cells

4. Pollination in water hyacinth and water lily is brought about by the agency of

(1) birds

(2) bats

(3) water

(4) insects or wind

5. The ovule of an angiosperm is technically equivalent to

(1) megaspore mother cell

(2) megaspore

(3) megasporangium

(4) megasporophyll

JMD/E3

from one linkage group to another is called

(1) translocation

crossing-over www.FirstRanker.com

(3) inversion (4) duplication

8. The equivalent of a structural gene is

(1) operon

(2) recon

(3) muton

(4) cistron

9. A true breeding plant is

(1) near homozygous and produces offspring of its own kind

(2) always homozygous recessive in its genetic constitution

(3) one that is able to breed on its own

produced due to cross-pollination among unrelated plants

10. Which of the following rRNAs acts as structural RNA as well as ribozyme in bacteria?

(1) 23 S rRNA

(2) 5.8 S rRNA

(3) 5 S rRNA

(4) 18 S rRNA

11. Stirred-tank bioreactors have been designed

(1) availability of oxygen throughout the process

(2) ensuring anaerobic conditions in the culture vessel

(3) purification of product

(4) addition of preservatives to the product

www.FirstRanker.com

- 23. Methanogens belong to
  - (1) Dinoflagellates
  - (2) Slime moulds

### www.FirstRanker.com

(3) Eubacteria

Archaebacteria

- 24. Select the wrong statement.
  - (1) Diatoms are chief producers in the oceans.
  - (2) Diatoms are microscopic and float passively in water.
  - (3) The walls of diatoms are easily destructible.
  - (4) 'Diatomaceous earth' is formed by the cell walls of diatoms.
- 25. The label of a herbarium sheet does not carry information on
  - (1) local names
  - (2) height of the plant
  - (3) date of collection
  - name of collector
- **26.** Conifers are adapted to tolerate extreme environmental conditions because of
  - thick cuticle
    - (2) presence of vessels
    - (3) broad hardy leaves
    - (4) superficial stomata
- 27. Which one of the following statements is wrong?
  - (1) Agar-agar is obtained from *Gelidium* and *Gracilaria*.
  - (2) Laminaria and Sargassum are used as food.
    - (3) Algae increase the level of dissolved oxygen in the immediate environment.
    - (4) Algin is obtained from red algae, and carrageenan from brown algae.

groundnut, radish, gram and turnip have stamens with different lengths in their flowers?

www.FirstRanker.com

- (1) Five
- (2) Six
- (3) Three
- JAY Four
- 30. Radial symmetry is found in the flowers of
  - (1) Pisum
  - (2) Cassia
  - 13↑ Brassica
    - (4) Trifolium
- 31. Free-central placentation is found in
  - JY Brassica
  - (2) Citrus
    - (3) Dianthus
  - JAT Argemone
- 32. Cortex is the region found between
  - (1) endodermis and pith
  - (2) endodermis and vascular bundle
  - (3) epidermis and stele
    - (4) pericycle and endodermis
- 33. The balloon-shaped structures called tyloses
  - are extensions of xylem parenchyma cells into vessels
  - (2) are linked to the ascent of sap through xylem vessels
  - (3) originate in the lumen of vessels
  - (4) characterize the sapwood

istaiu. have their

ers of

(2) Methanogens-Prokaryotes

(3) Gas vacuoles-Green bacteria

36. Select the wrong statement.

Cyanobacteria lack flagellated cells.

(2) Mycoplasma wall-less microorganism.

(3) Bacterial cell wall is made up of peptidoglycan.

(4) Pili and fimbriae are mainly involved in motility of bacterial cells.

cell organelle containing hydrolytic enzymes is

ribosome

(2) mesosome

(3) lysosome

(4) microsome

38. During cell growth, DNA synthesis takes place in

(1) G<sub>2</sub> phase

(2) M phase

(3) S phase

(4) G<sub>1</sub> phase

39. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins?

(1) Pyruvic acid

12 Acetyl CoA

(3) Glucose-6-phosphate

(4) Fructose 1,6-bisphosphate

JMD/E3

41. You are given a tissue with its potential for differentiation in an artificial culture. Which of the following pairs of hormones would you

Large central vacuoles—AWWW.FirstRanker.como the mediumwuFirstRankerscomell as roots?

(1) Auxin and abscisic acid

(2) Gibberellin and abscisic acid

(3) IAA and gibberellin

Auxin and cytokinin

42. Phytochrome is a

(1) lipoprotein

(2) chromoprotein

(3) flavoprotein

(4) glycoprotein

43. Which is essential for the growth of root tip?

ULY Ca

(2) Mn

(3) Zn

(4) Fe

44. The process which makes major difference between C<sub>3</sub> and C<sub>4</sub> plants is

(1) photorespiration

(2) respiration

(3) glycolysis

(4) Calvin cycle

45. Which one of the following statements is not correct?

In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem.

(2) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes.

(3) Offspring produced by the asexual reproduction are called clone.

(4) Microscopic, motile asexual reproductive structures are called zoospores.

[ P.T.O.

481 LNG-20 (4) Multiload 375 vasectomy?

- 48. Which of the following is incorrect regarding www.FirstRanker.com
  - (1) Vasa deferentia is cut and tied
  - (2) Irreversible sterility
  - (3) No sperm occurs in seminal fluid
  - A No sperm occurs in epididymis
- 49. Embryo with more than 16 blastomeres formed due to in vitro fertilization is transferred into
  - (1) fimbriae
  - (2) cervix
  - (3) uterus
  - (4) Tallopian tube
- 50. Which of the following depicts the correct pathway of transport of sperms?
  - (1) Rete testis → Vas deferens → Efferent ductules → Epididymis
  - (2) Efferent ductules → Rete testis → Vas deferens → Epididymis
  - (a) Rete testis  $\rightarrow$  Efferent ductules  $\rightarrow$ Epididymis → Vas deferens
  - (4) Rete testis → Epididymis → Efferent ductules  $\rightarrow$  Vas deferens
- 51. Match Column-II with Column-II and select the correct option using the codes given below:

- a. Mons pubis
- (i) Embryo formation
- b. Antrum
- (ii) Sperm
- Trophectoderm
- (iii) Female external genitalia
- d. Nebenkern
- (iv) Graafian follicle

# Codes:

	а	b	С	d
(1)	(iii)	(i)	(iv)	(ii)
(2)	(i)	(iv)	(iii)	(ii)
(3)	(iii)	(iv)	(ii)	(i)
(4)	(iii)	(iv)	(i)	(ii)

JMD/E3

6

(3) www.FirstRanker.com

probability of their son being colour-blind is

- (1) 0.75(2) 1
- 54. Genetic drift operates in
- (1) non-reproductive population
  - (2) slow reproductive population
  - (3) small isolated population
  - (4) large isolated population
- 55. In Hardy-Weinberg equation, the frequency of heterozygous individual is represented by
  - (1) pq
- (2)  $q^2$
- (3)  $p^2$
- 56. The chronological order of human evolution from early to the recent is
  - (1) Ramapithecus → Homo Australopithecus → Homo erectus
  - (2) Australopithecus → Homo habilis Ramapithecus → Homo erectus
  - (3) Australopithecus → Ramapithecus Homo habilis  $\rightarrow$  Homo erectus
  - . If Ramapithecus ightarrow Australopithecus ightarrowHomo habilis  $\rightarrow$  Homo erectus
- 57. Which of the following is the correct sequence of events in the origin of life?
  - I. Formation of protobionts
  - II. Synthesis of organic monomers
  - III. Synthesis of organic polymers
  - IV. Formation of DNA-based genetic systems
  - (1) II, III, I, IV
  - (2) II, III, IV, I
- (3) I, II, III, IV
  - (4) I, III, II, IV



blind is		
	59. DNA-dependent RNA polymerase catalyzes	(1) (i) (iv) (ii) (iii)
	transcription on one strand of the DNA	(2) (iii) (iv) (i) (ii) (ii) (3) (iii) (i) (ii) (ii)
	which is called the www.FirstRa	nker. (3) (ii) (i) www.FirstRanker.com
	(1) alpha strand	(iii) (i) (iv) (ii) <b>64.</b> Biochemical Oxygen Demand (BOD) may <b>not</b>
	(2) antistrand	be a good index for pollution for water bodies
	13) template strand	receiving effluents from
	(4) coding strand	(1) petroleum industry
	<b>60.</b> Interspecific hybridization is the mating of	(2) sugar industry
		(3) domestic sewage (4) dairy industry
equency	superior males and females of different breeds	65. The principle of competitive exclusion was
nted by	•	stated by
	(2) more closely related individuals within same breed for 4-6 generations	(1) MacArthur
		(2) Verhulst and Pearl
	(3) animals within same breed without having common ancestors	(3) C. Darwin
volution	_	(4) G. F. Gause
volution	(4) two different related species	66. Which of the following National Parks is
bilis →	61. Which of the following is correct regarding	home to the famous musk deer or hangul? (1) Eaglenest Wildlife Sanctuary, Arunachal
S	AIDS causative agent HIV?	Pradesh
bilis →	(1) HIV is unenveloped retrovirus.	(2) Dachigam National Park, Jammu &
	(2) HIV does not escape but attacks the	Kashmir
ecus →	acquired immune response.	(3) Keibul Lamjao National Park, Manipur
	(3) HIV is enveloped virus containing one	(4) Bandhavgarh National Park, Madhya Pradesh
cus →	molecule of single-stranded RNA and one	67. A lake which is rich in organic waste may
	molecule of reverse transcriptase.	result in
correct	(4) HIV is enveloped virus that contains two identical molecules of single-stranded	(1) increased population of fish due to lots of
life?	RNA and two molecules of reverse	nutrients
	transcriptase.	(2) mortality of fish due to lack of oxygen
		(3) increased population of aquatic organisms due to minerals
	<b>62.</b> Among the following edible fishes, which one is a marine fish having rich source of	drying of the lake due to algal bloom
	omega-3 fatty acids?	68. The highest DDT concentration in aquatic
systems	(1) Mrigala	food chain shall occur in
	(2) Mackerel	(1) crab
	• •	De eel
	(3) Mystus	(3) phytoplankton
	Mangur Mangur	(4) seagull
	JMD <b>/E3</b>	7 [ P.T.O.
	· · · · · · · · · · · · · · · · · · ·	·

- a. Familyb. Order(i) Diptera(ii) Arthropoda
- . Class (iii) Muscidae

(iv)

d. Phylum www.FirstRanker.com

## Codes:

(4) (iii)

a b c d
(1) (iv) (iii) (ii) (i)
(2) (iv) (ii) (i) (iii)
(3) (iii) (i) (iv) (ii)

71. Choose the correct statement.

(ii)

(1) All reptiles have a three-chambered heart.

(i)

(2) All Pisces have gills covered by an operculum.

(3) All mammals are viviparous.

- (4) All cyclostomes do not possess jaws and paired fins.
- 72. Study the four statements (A-D) given below and select the two correct ones out of them:
  - A. Definition of biological species was given by Ernst Mayr.
  - B. Photoperiod does not affect reproduction in plants.
  - C. Binomial nomenclature system was given by R. H. Whittaker.
  - D. In unicellular organisms, reproduction is synonymous with growth.

The two correct statements are

(1) A and D

(2) A and B


B and C

(4) C and D

- 73. In male cockroaches, sperms are stored in which part of the reproductive system?
  - (1) Testes
  - (2) Vas deferens
  - 131 Seminal vesicles
  - (4) Mushroom glands
- 74. Smooth muscles are
  - H involuntary, cylindrical, striated
  - (2) voluntary, spindle-shaped, uninucleate
  - (3) involuntary, fusiform, non-striated
  - (4) voluntary, multinucleate, cylindrical

folding of most proteins?

- (1) Hydrophobic interaction
- (2) Ester bonds WWW.FirstRanker.com Hydrogen bonds
  - (4) Electrostatic interaction
- **77.** Which of the following describes the given graph **correctly**?




- Endothermic reaction with energy A in absence of enzyme and B in presence of enzyme
- Exothermic reaction with energy A in absence of enzyme and B in presence of enzyme
  - (3) Endothermic reaction with energy A in presence of enzyme and B in absence of enzyme
  - (4) Exothermic reaction with energy A in presence of enzyme and B in absence of enzyme
- **78.** When cell has stalled DNA replication fork, which checkpoint should be predominantly activated?
  - (1) M
  - (2) Both G<sub>2</sub>/M and M
  - (3)  $G_1/S$
  - (4)  $G_2/M$

nensional	d. Zygotene (iv) Chromosomes align at	(3) Erythrocytes
	equatorial plate	(4) Leucocytes
	Codes :	(esaconame a pepwwwoFirstRankeracomainly
	a b c d www.i ii stitaii (1) (ii) (iv) (iii) (i)	on hepatocytes, adipocytes and enhances cellular glucose uptake and utilization.
:he given	(2) (iv) (iii) (ii) (i)	(1) Secretin (2) Gastrin
	(iii) (iv) (ii) (i)	(3) Insulin (4) Glucagon
	(4) (i) (iv) (ii) (iii),  80. Which hormones do stimulate the	87. Osteoporosis, an age-related disease of skeletal system, may occur due to
	production of pancreatic juice and	(1) decreased level of estrogen
	bicarbonate? (1) Cholecystokinin and secretin	(2) accumulation of uric acid leading to
	(2) Insulin and glucagon	inflammation of joints
	(3) Angiotensin and epinephrine	<ul><li>(3) immune disorder affecting neuro- muscular junction leading to fatigue</li></ul>
	(4) Gastrin and insulin	(4) high concentration of Ca <sup>++</sup> and Na <sup>+</sup>
	81. The partial pressure of oxygen in the alveoli	88. Serum differs from blood in
	of the lungs is	(1) lacking clotting factors
	(1) less than that in the blood	(2) lacking antibodies
roduct	(2) less than that of carbon dioxide	(3) lacking globulins
	(3) equal to that in the blood	(4) Tacking albumins
rgy A in	more than that in the blood	89. Lungs do not collapse between breaths and
sence of	82. Choose the correct statement.	some air always remains in the lungs which
gy A in	Photoreceptors in the human eye are depolarized during darkness and become	can never be expelled because
sence of	hyperpolarized in response to the light	(1) there is a positive intrapleural pressure
4 :	stimulus.	pressure in the lungs is higher than the atmospheric pressure
rgy A in	(2) Receptors do not produce graded	(3) there is a negative pressure in the lungs
	potentials.	
gy A in	(3) Nociceptors respond to changes in pressure.	pulling at the lung walls
sence of	(4) Meissner's corpuscles are thermo-	90. The posterior pituitary gland is not a 'true'
ion fork,	receptors.	endocrine gland because
ninantly	83. Graves' disease is caused due to	it is under the regulation of hypo-
	hyposecretion of adrenal gland	thalamus (2) it secretes enzymes
	hypersecretion of adrenal gland	(3) it is provided with a duct
	(3) hyposecretion of thyroid gland	(4) it only stores and releases hormones
	(4) hypersecretion of thyroid gland	(1) It only occurs and reconstruction
	JMD <b>/E3</b>	9 [ P.T.O.
		·
		,
!		




- (1)  $\frac{2}{k} = \frac{3}{k_1 + k_2 + k_3} + \frac{1}{k_4}$
- (2)  $\frac{1}{k} = \frac{1}{k_1} + \frac{1}{k_2} + \frac{1}{k_3} + \frac{3}{2k_4}$
- (3)  $k = k_1 + k_2 + k_3 + 3k_4$
- (4)  $k = \frac{2}{3}(k_1 + k_2 + k_3) + 2k_4$
- **92.** The potential difference  $(V_A V_B)$  between the points A and B in the given figure is



- (1) +6 V
- (2) +9 V
- (3) -3 V
- (4) +3 V
- 93. A filament bulb (500 W, 100 V) is to be used in a 230 V main supply. When a resistance R is connected in series, it works perfectly and the bulb consumes 500 W. The value of R is
  - (1) 26 Ω
- (2)  $13 \Omega$
- (3) 230  $\Omega$
- (4) 46 Ω
- 94. A long wire carrying a steady current is bent into a circular loop of one turn. The magnetic field at the centre of the loop is B. It is then bent into a circular coil of n turns. The magnetic field at the centre of this coil of n turns will be
  - 2nB
- $\cdot$ (2)  $2n^2B$
- (3)/ nB
- (4)  $n^2 B$

JMD/E3

- 96. An electron is moving in a circular path under the influence of a transverse magnetic field of 3.57 × 10<sup>-2</sup> T. If the value of e/mww/firstRanker.comfrequency of revolution of the electron is
  - (1) 62·8 MHz
- (2) 6.28 MHz
- (3) 1 GHz
- (4) 100 MHz
- **97.** Which of the following combinations should be selected for better tuning of an *L-C-R* circuit used for communication?
  - (1)  $R = 15 \Omega$ , L = 3.5 H,  $C = 30 \mu F$
  - (2)  $R = 25 \Omega$ , L = 1.5 H,  $C = 45 \mu F$
  - (3)  $R = 20 \Omega$ , L = 1.5 H,  $C = 35 \mu F$
  - (4)  $R = 25 \Omega$ , L = 2.5 H,  $C = 45 \mu F$
- **98.** A uniform magnetic field is restricted within a region of radius r. The magnetic field changes with time at a rate  $\frac{d\vec{B}}{dt}$ . Loop 1 of radius R > r encloses the region r and loop 2 of radius R is outside the region of magnetic field as shown in the figure below. Then the e.m.f. generated is



- (1)  $-\frac{d\vec{B}}{dt}\pi R^2$  in loop 1 and zero in loop 2
- (2)  $-\frac{d\vec{B}}{dt}\pi r^2$  in loop 1 and zero in loop 2
- (3) zero in loop 1 and zero in loop 2
- (4)  $-\frac{d\vec{B}}{dt}\pi r^2$  in loop 1 and

 $-\frac{d\vec{B}}{dt}\pi r^2$  in loop 2

circular nsverse e value lency of

should 1 L-C-R

within ic field

op 1 of

loop 2 agnetic nen the

pop 2

юр 2

loop 2

a 220 V source. When the capacitor is 50% charged, the peak value of the displacement | 105. A linear aperture whose width is 0.02 cm is current is

- (1) 4·4 A
- (3) 2·2 A
- (4) 11 A

101. Two identical glass ( $\mu_g = 3/2$ ) equiconvex lenses of focal length f each are kept in contact. The space between the two lenses is filled with water ( $\mu_w = 4/3$ ). The focal length of the combination is

- (1) 4f/3
- (2) 3f/4
- (3) f/3
- (4) f

102. An air bubble in a glass slab with refractive index 1.5 (near normal incidence) is 5 cm deep when viewed from one surface and 3 cm deep when viewed from the opposite face. The thickness (in cm) of the slab is

- (1) 12
- (2) 16
- (3) 8
- (4) 10

103. The interference pattern is obtained with two coherent light sources of intensity ratio n. In the interference pattern, the ratio

$$\frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}}$$

will be

- $(1) \frac{\sqrt{n}}{(n+1)^2}$

(4) concave, -0.25 diopter

placed immediately in front of a lens of focal (2) 11 1/2 W.W. FirstRanker.comength 60 www.feirstRanker.comminated normally by a parallel beam of wavelength  $5 \times 10^{-5}$  cm. The distance of the first dark band of the diffraction pattern from the centre of the screen is

- (1) 0·20 cm
- (2) 0·15 cm
- (3) 0·10 cm
- (4) 0.25 cm

**106.** Electrons of mass m with de-Broglie wavelength  $\lambda$  fall on the target in an X-ray tube. The cutoff wavelength  $(\lambda_0)$  of the emitted X-ray is

(1) 
$$\lambda_0 = \frac{2m^2c^2\lambda^3}{h^2}$$

- (2)  $\lambda_0 = \lambda$
- (3)  $\lambda_0 = \frac{2mc\lambda^2}{h}$
- $(4) \quad \lambda_0 = \frac{2h}{mc}$

107. Photons with energy 5 eV are incident on a cathode C in a photoelectric cell. The maximum energy of emitted photoelectrons is 2 eV. When photons of energy 6 eV are incident on C, no photoelectrons will reach the anode A, if the stopping potential of A relative to C is


- (1) -1 V
- (2) -3 V
- (3) + 3 V
- (4) + 4 V

JMD/E3


11

[ P.T.O.

- 109. The half-life of a radioactive substance is 30 minutes. The time (in minutes) taken between 40% decay WWW 5 its Rankenuc m same radioactive substance is
  - (1) 45
- (2) 60
- (3) 15
- (4) 30
- 110. For CE transistor amplifier, the audio signal voltage across the collector resistance of  $2 k\Omega$  is 4 V. If the current amplification factor of the transistor is 100 and the base resistance is  $1 k\Omega$ , then the input signal voltage is
  - (1) 30 mV
- (2) 15 mV
- (3) 10 mV
- (4) 20 mV
- 111. The given circuit has two ideal diodes connected as shown in the figure below. The current flowing through the resistance  $R_1$  will be



- (1) 1·43 A
- (2) 3·13 A
- (3) 2·5 A
- (4) 10·0 A
- 112. What is the output Y in the following circuit, when all the three inputs A, B, C are first 0 and then 1?




- (1) 1, 0
- (2) 1, 1
- (3) 0, 1
- (4) 0, 0

- 114. Two cars P and Q start from a point at the same time in a straight line and their position. First Panker by  $and x_Q(t) = ft t^2$ . At what time do the cars have the same velocity?
  - (1)  $\frac{a+f}{2(1+b)}$
- $2) \frac{f-a}{2(1+b)}$
- $(3) \ \frac{a-f}{1+b}$
- 4)  $\frac{a+f}{2(b-1)}$
- 115. In the given figure,  $a = 15 \text{ m/s}^2$  represents the total acceleration of a particle moving in the clockwise direction in a circle of radius R = 2.5 m at a given instant of time. The speed of the particle is



- (1) 5.7 m/s
- (2) 6·2 m/s
- (3) 4.5 m/s
- (4) 5·0 m/s
- 116. A rigid ball of mass m strikes a rigid wall at 60° and gets reflected without loss of speed as shown in the figure below. The value of impulse imparted by the wall on the ball will be



- (1)  $\frac{mV}{2}$
- (2)  $\frac{mV}{3}$
- 31 mV
- $(4) \ 2mV$

at the their  $at + bt^2$ 

ie cars

esents ving in radius

e. The

wall at speed ilue of ie ball


- (2)  $160 \text{ m s}^{-1}$
- (3)  $100 \text{ m s}^{-1}$
- (4)  $80 \text{ m s}^{-1}$


- 118. Two identical balls A and B having velocities of 0.5 m/s and -0.3 m/s respectively collide elastically in one dimension. The velocities of Band A after the collision respectively will be
  - (1) -0.3 m/s and 0.5 m/s
  - (2) 0.3 m/s and 0.5 m/s
  - (3) -0.5 m/s and 0.3 m/s
  - (4) 0.5 m/s and -0.3 m/s
- 119. A particle moves from a point  $(-2\hat{i} + 5\hat{j})$  to  $(4\hat{j}+3\hat{k})$  when a force of  $(4\hat{i}+3\hat{j})$  N is applied. How much work has been done by the force?
  - (1) 5 J
  - (2) 2 J
  - (3) 8 J
  - (4) 11 J
- 120. Two rotating bodies A and B of masses mand 2m with moments of inertia  $I_A$  and  $I_B(I_B > I_A)$  have equal kinetic energy of rotation. If  $L_A$  and  $L_B$  be their angular momenta respectively, then
  - $(1) L_B > L_A$
  - (2)  $L_A > L_B$
  - $(3) L_A = \frac{L_B}{2}$
  - $(4) L_A = 2L_B$


- 122. A light rod of length l has two masses  $m_1$  and www.FirstRanker.comttached twww.FirstRanker.coment of inertia of the system about an axis perpendicular to the rod and passing through the centre of mass is
  - (1)  $(m_1 + m_2)l^2$
- $(3) \ \frac{m_1 m_2}{m_1 + m_2} l^2$


(4) 1:5

- 123. Starting from the centre of the earth having radius R, the variation of g (acceleration due to gravity) is shown by









- 124. A satellite of mass m is orbiting the earth (of radius R) at a height h from its surface. The total energy of the satellite in terms of  $g_0$ , the value of acceleration due to gravity at the earth's surface, is
  - (1)  $\frac{2mg_0R^2}{-}$

JMD/E3

13

[ P.T.O.

 $\rho_1 > \rho_2 > \rho_3$ ), having the surface tension T, rise to the same height 131. A given sample of all literal gas occup. in three identical capillaries. The angles of contact  $\theta_1$ ,  $\theta_2$  and  $\theta_2$  obey

- (1)  $\frac{\pi}{2} < \theta_1 < \theta_2 < \theta_3 < \pi$
- (2)  $\pi > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2}$
- (3)  $\frac{\pi}{2} > \theta_1 > \theta_2 > \theta_3 \ge 0$
- (4)  $0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$
- 127. Two identical bodies are made of a material for which the heat capacity increases with temperature. One of these is at 100 °C, while the other one is at 0 °C. If the two bodies are brought into contact, then, assuming no heat loss, the final common temperature is
  - (1) less than 50 °C but greater than 0 °C
  - (2) 0 °C
  - (3) 50 °C
  - (4) more than 50 °C
- 128. A body cools from a temperature 3T to 2Tin 10 minutes. The room temperature is T. Assume that Newton's law of cooling is applicable. The temperature of the body at the end of next 10 minutes will be

- 129. One mole of an ideal monatomic undergoes a process described by equation  $PV^3$  = constant. The heat capacity of the gas during this process is
  - (1) 2R
- (3)  $\frac{3}{2}R$
- (4)  $\frac{5}{2}R$

- a volume V at a pressure P and absolu temperature T. The mass of each molecu www.geirstRamker.com following give the density of the gas
  - (1) P/(kTV)
- (2) mkT
- (3) P/(kT)
- (4) Pm/(kT)
- 132. A body of mass m is attached to lower end of a spring whose upper is fixed. The spring has negligible ma When the mass m is slightly pulled do and released, it oscillates with a time pe of  $3 \, \text{s}$ . When the mass m is increased by 1 kg, the time period of oscillation becomes 5 s. The value of m in kg is
  - 16

- 133. The second overtone of an open o pipe has the same frequency as the overtone of a closed pipe L metre long. length of the open pipe will be
- (2) 4L
- (3) L
- (4) 2L
- 134. Three sound waves of equal ampli have frequencies (n-1),  $\overline{n}$ , (n+1). superimpose to give beats. The number beats produced per second will be
  - (1) 3
- (2) 2
- (3) 1
- (4) 4
- 135. An electric dipole is placed at an angle with an electric field intensity  $2 \times 10^{\overline{5}}$  M experiences a torque equal to 4 N n charge on the dipole, if the dipole ler 2 cm, is
  - (1) 5 mC
  - (2) 7 µC
  - (3) 8 mC
  - (4) 2 mC

5 Occupie id absolute h molecule owing gives

ed to the upper end zible mass. ulled down time period increased oscillations kg is

pen organ as the first e long. The

amplitudes n+1). They number of be

angle of 30° 10<sup>5</sup> N/C. It 4 N m. The ple length is

have electron density along the axes?

$$d_{z^2}, d_{x^2-y^2}$$

(2)  $d_{xy}$ ,  $d_{x^2-y^2}$ 

(3)  $d_{z^2}$ ,  $d_{xz}$ 

(4)  $d_{xz}$ ,  $d_{uz}$ 

138. The correct geometry and hybridization for XeF<sub>4</sub> are

- (1) planar triangle,  $sp^3d^3$
- (2) square planar,  $sp^3d^2$
- (3) octahedral,  $sp^3d^2$
- trigonal bipyramidal, sp3d

139. Among the following, which one is a wrong statement?

- (1) SeF<sub>4</sub> and CH<sub>4</sub> have same shape.
- (2) I<sub>3</sub> has bent geometry.

PH<sub>5</sub> and BiCl<sub>5</sub> do not exist.

- (4)  $p\pi$ - $d\pi$  bonds are present in SO<sub>2</sub>.
- 140. The correct increasing order of trans-effect of the following species is

(1) 
$$Br^- > CN^- > NH_3 > C_6H_5^-$$

(2) 
$$CN^- > Br^- > C_6H_5^- > NH_3$$

(3) 
$$NH_3 > CN^- > Br^- > C_6H_5^-$$

$$CN^- > C_6H_5^- > Br^- > NH_3$$

141. Which one of the following statements related to lanthanons is incorrect?

- (1) All the lanthanons are much more reactive than aluminium.
- (2) Ce (+4) solutions are widely used as oxidizing agent in volumetric analysis.
- (3) Europium shows +2 oxidation state.
- (4) The basicity decreases as the ionic radius decreases from Pr to Lu.

JMD/E3

(L) Chloroethene

www.FirstRanker.com

- (3) Chlorobenzene
- (4) Bromobenzene

144. In which of the following molecules, all atoms are coplanar?

(1) 
$$CH_3$$
  $C=C$   $CN$  (2)  $CH_3$   $CH$ 



145. Which one of the following structures represents nylon 6,6 polymer?

(1) 
$$\begin{pmatrix} H_2 & H_2$$

(2) 
$$\left(\begin{array}{c} H_2 \\ H_2 \\ H_2 \\ \end{array}\right) \left(\begin{array}{c} H_2 \\ N + (CH_2)_6 - NH \\ O \\ \end{array}\right)_n$$

(3) 
$$\begin{pmatrix} H_2 & H_2 & H_2 \\ C & C & C \\ NH_2 & CH_3 \end{pmatrix}_{66}$$

$$(4) \quad \begin{pmatrix} H_2 & H_2 & H_2 \\ C & C & C \\ NH_2 & NH_2 \end{pmatrix}_{66}$$

[ P.T.O.

15

(2) 2 and 5

(3) 2 and 3 www.FirstRanker.com

(4) 3 and 4

147. Which of the following compounds shall **not** produce propene by reaction with HBr followed by elimination or direct only elimination reaction?

(2) 
$$H_3C - C - CH_2Br$$

(3) 
$$H_2C \xrightarrow{CH_2} CH_2$$

**148.** Which one of the following nitro-compounds **does not** react with nitrous acid?

(1) 
$$H_3C$$
  $-C-NO_2$   $H_3C$ 

$$(2) \quad \overset{\text{CH}_3}{\underset{\text{O}}{\text{H}}} \overset{\text{CH}_3}{\underset{\text{NO}_2}{\text{NO}_2}}$$

$$\begin{array}{ccc} H_3C & H_2 \\ & & C \\ & & & \\$$

JMD/E3

(4) DNA → Carbohydrates → Proteins

150. The Way First Ranker Grown der of names four aldoses with configuration given belo

respectively, is

(1) L-erythrose, L-threose, D-erythros D-threose

D-erythrose, D-threose, L-erythrose, L-threose

(3) L-erythrose, L-threose, L-erythros D-threose

(4) D-threose, D-erythrose, L-threose L-erythrose

151. In the given reaction

the product P is

www.FirstRahker.com

- (2) Both II and III www.FirstRanker.com (3) III only

(4) Both I and III

156. The correct order of strengths of the carboxylic acids

is

- (1) III > II > I
- (2) II > I > III
- (3) I > II > III
- (4) II > III > I
- 157. The compound that will react most readily with gaseous bromine has the formula
  - (1)  $C_4H_{10}$
  - (2)  $C_2H_4$
  - (3)  $C_3H_6$
  - (4)  $C_2H_2$


[ P.T.O.

153. Consider the reaction

$$CH_3CH_2CH_2Br + NaCN \rightarrow CH_3CH_2CH_2CN + NaBr$$

This reaction will be the fastest in

- (1) N, N'-dimethylformamide (DMF)
- (2) water
- (3) ethanol
- (4) methanol
  - 154. The correct structure of the product Aformed in the reaction



is









- solution of AgNO<sub>3</sub> with electrolytic conductivity of  $5.76 \times 10^{-3} \text{ S cm}^{-1}$  at 298 K is www.FirstRanker.com
  - (1)  $0.086 \text{ S cm}^2/\text{mol}$
  - (2) 28.8 S cm<sup>2</sup>/mol
  - (3)  $2.88 \text{ S cm}^2/\text{mol}$
  - (4) 11.52 S cm<sup>2</sup>/mol
- 160. The decomposition of phosphine (PH<sub>3</sub>) on tungsten at low pressure is a first-order reaction. It is because the
  - (1) rate is independent of the surface coverage
  - (2) rate of decomposition is very slow
  - (3) rate is proportional to the surface coverage
  - (4) rate is inversely proportional to the surface coverage
- 161. The coagulation values in millimoles per litre of the electrolytes used for the coagulation of As<sub>2</sub>S<sub>3</sub> are given below:
  - I. (NaCl) = 52,
- II.  $(BaCl_2) = 0.69$ ,
- III. (MgSO<sub>4</sub>) = 0.22

The **correct** order of their coagulating power is

- (1) III > II > I
- (2) III > I > II
- (3) I > II > III
- (4) II > I > III
- 162. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of 3 amperes is
  - (1) 220 minutes
  - (2) 330 minutes
  - (3) 55 minutes
  - (4) 110 minutes

(2) 
$$\Delta S = RT \ln \left( \frac{p_i}{p_i} \right)$$
  
www.FirstRanker.com

(3) 
$$\Delta S = nR \ln \left( \frac{p_f}{p_i} \right)$$

(4) 
$$\Delta S = nR \ln \left( \frac{p_i}{p_f} \right)$$

- 165. The van't Hoff factor (i) for a dilute aqueou solution of the strong electrolyte bariu hydroxide is
  - (1) 2
- (2) 3
- (3) 0
- (4) 1
- 166. The percentage of pyridine  $(C_5H_5N)$  th forms pyridinium ion  $(C_5H_5N^+H)$  in a 0·10 aqueous pyridine solution  $(K_b)$  is
  - (1) 0.77%
  - (2) 1.6%
  - (3) 0.0060%
  - (4) 0.013%
- 167. In calcium fluoride, having the fluor structure, the coordination numbers calcium ion (Ca<sup>2+</sup>) and fluoride ion (F<sup>-</sup>) are
  - (1) 8 and 4
  - (2) 4 and 8
  - (3) 4 and 2
  - (4) 6 and 6
- 168. If the  $E_{cell}^{\circ}$  for a given reaction has a negarable, which of the following gives correct relationships for the values of and  $K_{eq}$ ?
  - (1)  $\Delta G^{\circ} < 0$ ;  $K_{eq} > 1$
  - (2)  $\Delta G^{\circ} < 0$ ;  $K_{eq} < 1$
  - (3)  $\Delta G^{\circ} > 0$ ;  $K_{eq} < 1$
  - (4)  $\Delta G^{\circ} > 0$ ;  $K_{eq} > 1$

	would be $(1) 1.6 \times 10^{-11} M$ $(2) zero$ WWW.First	176. The suspension of slaked lime in water is known as www.FirstRanker.com (1) milk of lime
	(3) $1.26 \times 10^{-5} M$ (4) $1.6 \times 10^{-9} M$	<ul><li>(2) aqueous solution of slaked lime</li><li>(3) limewater</li><li>(4) quicklime</li></ul>
ate aqueous lyte barium	171. Suppose the elements X and Y combine to form two compounds XY <sub>2</sub> and X <sub>3</sub> Y <sub>2</sub> . When 0·1 mole of XY <sub>2</sub> weighs 10 g and 0·05 mole of X <sub>3</sub> Y <sub>2</sub> weighs 9 g, the atomic weights of X and Y are	en nitrogen in NO <sub>2</sub> <sup>+</sup> , NO <sub>3</sub> and NH <sub>4</sub> <sup>+</sup> respectively
$C_5H_5N$ ) that in a $0.10~M$	(1) 20, 30     (2) 30, 20       (3) 40, 30     (4) 60, 40	(2) $sp^2$ , $sp$ and $sp^3$ (3) $sp$ , $sp^3$ and $sp^2$
·	172. The number of electrons delivered at the cathode during electrolysis by a current of 1 ampere in 60 seconds is (charge on electron = $1.60 \times 10^{-19}$ C)	of 178. Which of the following fluoro-compounds is most likely to behave as a Lewis base?
the fluorite numbers for on (F <sup>-</sup> ) are	(1) $3.75 \times 10^{20}$ (2) $7.48 \times 10^{23}$ (3) $6 \times 10^{23}$ (4) $6 \times 10^{20}$	(1) CF <sub>4</sub> (2) SiF <sub>4</sub> (3) BF <sub>3</sub> (4) PF <sub>3</sub> 32+4 1  179. Which of the following pairs of ions is
	<ul> <li>73. Boric acid is an acid because its molecule</li> <li>(1) accepts OH<sup>-</sup> from water releasing proton</li> <li>(2) combines with proton from water molecule</li> </ul>	isoelectronic and isostructural? (1) $SO_3^{2-}$ , $NO_3^{-}$ (2) $ClO_3^{-}$ , $SO_3^{2-}$ (3) $CO_3^{2-}$ , $NO_3^{-}$ (4) $ClO_3^{-}$ , $CO_3^{2-}$ (6+22
as a negative $g$ gives the alues of $\Delta G$	<ul> <li>(3) contains replaceable H<sup>+</sup> ion</li> <li>(4) gives up a proton</li> <li>74. AlF<sub>3</sub> is soluble in HF only in presence of KF. It is due to the formation of</li> <li>(1) AlH<sub>3</sub></li> <li>(2) K[AlF<sub>2</sub>H]</li> </ul>	polymeric. 35 +
	(3) $K_3[AIF_3H_3]$ (4) $K_3[AIF_6]$	(3) It is rendered passive by nitric acid.  19 Re20  [P.T.O.
		[ 1.1.0.

www.FirstRanker.com	www.FirstRanker.com
www.FirstRanker.com	