


Mathematics

Section - D

23.

Radius of the hemisphere =  $\frac{d}{2}$   
 $= \frac{4.2 \text{ cm}}{2}$

$$= 2.1 \text{ cm}$$



Side of cube = 6 cm.

a) Total surface area of block = Total surface area of cube + Curved surface area of hemisphere - area enclosed by base of hemisphere

$$= 6a^2 + \frac{4\pi r^2}{2} - \pi r^2$$

$$= 6a^2 + \pi r^2$$

$$= 6 \times 6^2 + \frac{22 \times (2.1)^2}{7} \text{ cm}^2$$

$$= 216 + \frac{22 \times 2.1 \times 2.1}{7} \text{ cm}^2$$

$$= [216 + 13.86] \text{ cm}^2$$

$$= 229.86 \text{ cm}^2$$

b) Volume of block formed = volume of cube + volume of hemisphere  
 $= a^3 + \frac{2}{3}\pi r^3$

$$= 6^3 + \frac{2 \times 22 \times 2.1^3}{3} \text{ cm}^3$$

$$= 216 + \frac{2 \times 22 \times 21 \times 21}{3 \times 1000} \text{ cm}^3$$

$$= 216 + 19.404 \text{ cm}^3$$

$$= \underline{235.404 \text{ cm}^3}$$

24. To prove: Square of hypotenuse, in a right triangle, is equal to the sum of squares of other two sides. (Pythagoras theorem)

That is,  $AC^2 = AB^2 + BC^2$ .

Construction: Construct  $BD \perp AC$

Name  $\angle BAC = \theta$ .

Then,  $\angle BCA = 90 - \theta$ ,  $\angle ABD = 90 - \theta$ ,  $\angle DBC = \theta$




Fig.

It is clear that,

$$\triangle ABD \sim \triangle ACB \sim \triangle BCD.$$

Using  $\triangle ABD \sim \triangle ACB$ , we get:

$$\frac{AB}{AC} = \frac{AD}{AB} \Rightarrow AB^2 = AC \times AD \quad \text{--- (1)}$$

Similarly, using  $\triangle BCD \sim \triangle ACB$ , we get:

$$\frac{BC}{AC} = \frac{CD}{BC} \Rightarrow BC^2 = AC \times CD \quad \text{--- (2)}$$

Adding (1) and (2), gives:

$$AB^2 + BC^2 = AC \times AD + AC \times CD$$

$$\Rightarrow AB^2 + BC^2 = AC (AD + CD)$$

$$\Rightarrow AB^2 + BC^2 = AC \times AC$$

$\Rightarrow$

$$AB^2 + BC^2 = AC^2.$$

Hence, proved that in a right triangle, sum of squares of two sides is equal to the square of hypotenuse.

25. graph paper (near graph, on Pg. no 22).

26. given - as per figure

Shadow of tower is 40 m longer at sun's altitude at  $30^\circ$

$$\therefore BD - BC = 40 \text{ m}$$

$$\Rightarrow CD = 40 \text{ m}$$

In  $\Delta ACB$ ,


$$\tan 60^\circ = \frac{AB}{BC}$$

$$\Rightarrow \sqrt{3} \times BC = AB \Rightarrow BC = \frac{AB}{\sqrt{3}} \quad \text{--- (1)}$$

In  $\Delta ADB$ ,

$$\tan 30^\circ = \frac{AB}{BD} \Rightarrow \frac{1}{\sqrt{3}} = \frac{AB}{40 + BC}$$

$$\Rightarrow 40 + BC = \sqrt{3} \times AB$$



$$\Rightarrow 40 + \frac{AB}{\sqrt{3}} = \sqrt{3} \times AB$$

[Put  $BC = \frac{AB}{\sqrt{3}}$  from ①]

$$\Rightarrow AB \left( \sqrt{3} - \frac{1}{\sqrt{3}} \right) = 40$$

$$\Rightarrow AB \left( \frac{3-1}{\sqrt{3}} \right) = 40$$

$$\Rightarrow AB \times \frac{2}{\sqrt{3}} = 40 \Rightarrow AB = \frac{40 \times \sqrt{3}}{2} \text{ m}$$

$$\Rightarrow AB = 20\sqrt{3} \text{ m}$$

Given, use  $\sqrt{3} = 1.732$

$$\therefore AB = 20 \times 1.732 \text{ m} = 34.64 \text{ m}$$

$\therefore$  Height of tower = 34.64 m.

27. Let the first term of given A.P. be 'a' and the common difference be 'd'.  
and  $a_p$  denotes  $p^{\text{th}}$  term.

$$\text{Given: } m(a_m) = n(a_n) \quad [m \neq n]$$

$$\text{To show: } a(m+n) = 0$$

$$m(a_m) = n(a_n)$$

$$\Rightarrow m[a + (m-1)d] = n[a + (n-1)d]$$

$$\Rightarrow am + md(m-1) = an + nd(n-1)$$

$$\Rightarrow am - an = nd(n-1) - md(m-1)$$

$$\Rightarrow a(m-n) = d[n(n-1) - m(m-1)]$$

$$\Rightarrow a(m-n) = d[n^2 - n - m^2 + m]$$

$$\Rightarrow a(m-n) = d[n^2 - m^2 + m - n]$$

$$\Rightarrow a(m-n) = d[(m+n)(n-m) + (m-n)]$$

$$\Rightarrow a(m-n) = d(m-n)[-(m+n) + 1]$$

$$\Rightarrow a - d[-(m+n) + 1] = 0$$

$$\Rightarrow [a + (m+n-1)d] = 0$$

$$\Rightarrow a + m + n = 0$$

$$\therefore a_{m+n} = 0$$

Hence, proved!

28.

Let the no. of books bought by the shopkeeper be 'n'.

Total money spent = Rs 80

$$\therefore \text{Cost of each book} = \frac{\text{Rs } 80}{n}$$

Now, given: He buys 4 more books, no. of books bought =  $n+4$   
(for same amount)

$$\text{New cost of each book} = \frac{\text{Rs } 80}{n+4}$$

Given, new cost of each book is Rs. 1 less than earlier.

$$\therefore \frac{80}{n} - \frac{80}{n+4} = 1$$

$$\Rightarrow 80 \left( \frac{1}{n} - \frac{1}{n+4} \right) = 1$$

$$\Rightarrow \frac{n+4-n}{n(n+4)} = \frac{1}{80} \Rightarrow \frac{4 \times 80}{n(n+4)} = 1$$

$$\Rightarrow n^2 + 4n - 320 = 0$$

Using quadratic formula;

$$\Rightarrow n = \frac{-4 \pm \sqrt{16 + 4 \times 320}}{2}$$

$$\Rightarrow x = \frac{-4 \pm \sqrt{64}}{2}$$

$$= \frac{-4 \pm 8}{2} \Rightarrow x = \frac{-4+8}{2} \text{ or } \frac{-4-8}{2}$$

$$\Rightarrow x = \frac{-4 + \sqrt{1296}}{2}$$

$$\Rightarrow x = \frac{-4 \pm 36}{2}$$

$$\Rightarrow x = \frac{-40 \text{ or } 32}{2} \Rightarrow x = -20 \text{ or } 16$$

Since, no. of books is a whole no., it cannot be negative

$x = -20$  can be ignored.

$$\therefore \boxed{x = 16}$$

No. of books bought by the shopkeeper = 16.


29.

To construct: a pair of tangents to a circle of radius  $= 4\text{ cm}$ ,  
from a point at a distance  $6\text{ cm}$  from centre.

Steps of construction:

- 1) Draw a circle of radius  $4\text{ cm}$  with  $O$  as the centre.
- 2) Take a point  $P$  at  $PO = 6\text{ cm}$ .

- 3) Join  $PO$ . Construct a perpendicular bisector of  $PO$  at  $M$  ( $PM = MO$ ,  $AB \perp PO$ )



- 4) With  $M$  as centre and  $PM (= MO)$  as radius, draw a circle touching the circle with centre  $O$  at  $T$  and  $Q$ .
- 5) Join  $PT$  and  $PQ$ .  
 $\therefore PT$  and  $PQ$  are required tangents.

So. To prove:  $\frac{1}{1+\sin^2\theta} + \frac{1}{1+\cos^2\theta} + \frac{1}{1+\sec^2\theta} + \frac{1}{1+\csc^2\theta} = 2$ .

Taking from LHS,

$$= \frac{1}{1+\sin^2\theta} + \frac{1}{1+\cos^2\theta} + \frac{1}{1+\csc^2\theta} + \frac{1}{1+\sec^2\theta}$$

$$= \frac{1}{1+\sin^2\theta} + \frac{1}{1+\cos^2\theta} + \frac{1}{1+\cos^2\theta} + \frac{1}{1+\sec^2\theta} \quad [\text{Re-arranging}]$$

$$= \frac{1}{1+\sin^2\theta} + \frac{1}{1+\left(\frac{1}{\sin^2\theta}\right)} + \frac{1}{1+\cos^2\theta} + \frac{1}{1+\left(\frac{1}{\cos^2\theta}\right)} \quad \left[ \begin{array}{l} \csc^2\theta = \frac{1}{\sin^2\theta} \\ \sec^2\theta = \frac{1}{\cos^2\theta} \end{array} \right]$$

$$= \frac{1}{1+\sin^2\theta} + \frac{1 \times \sin^2\theta}{1+\sin^2\theta} + \frac{1}{1+\cos^2\theta} + \frac{1 \times \cos^2\theta}{1+\cos^2\theta}$$

$$= \frac{1+\sin^2\theta}{1+\sin^2\theta} + \frac{1+\cos^2\theta}{1+\cos^2\theta}$$

$$= 1 + 1$$

$$= 2$$

$$= \text{RHS}$$

LHS = RHS  
Hence, proved!

Section - C

18. Given  $\tan(A+B) = 1$  and  $\tan(A-B) = \frac{1}{\sqrt{3}}$

$$\tan(A+B) = 1.$$

$$\Rightarrow \tan(A+B) = \tan 45^\circ$$

$$\Rightarrow A+B = 45^\circ \quad \text{--- ①}$$

Now taking,

$$\tan(A-B) = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \tan(A-B) = \tan 30^\circ$$

$$\Rightarrow A-B = 30^\circ \quad \text{--- ②}$$

Adding ① and ②;


$$A+B+A-B = 45^\circ + 30^\circ$$

$$\Rightarrow 2A = 75^\circ \Rightarrow A = \frac{75^\circ}{2} \Rightarrow A = 37.5^\circ$$

$$B = 45^\circ - A \Rightarrow B = 45^\circ - 37.5^\circ \Rightarrow B = 7.5^\circ$$

$$\therefore \boxed{A = 37.5^\circ, B = 7.5^\circ}$$

14. To prove: opposite sides of a quadrilateral circumscribing a circle subtend equal angles at the centre.



Construction: Constructed a quadrilateral ABCD, circumscribing a circle (centre O). Circle touches AB, BC, CD, DA at P, Q, R, S respectively.

To prove:  $\angle AOB + \angle COD = 180^\circ$   
Or  $\angle AOD + \angle BOC = 180^\circ$

We know, that tangents from same exterior point subtend equal angle at the centre of circle with radius.

$$\therefore \angle AOP = \angle AOS = \angle 1 \text{ (say)}$$

$$\text{Similarly, } \angle BOP = \angle BOQ = \angle 2$$

$$\angle COR = \angle COQ = \angle 3$$

$$\angle DOR = \angle DOS = \angle 4$$

$$\therefore \angle AOP + \angle BOP + \angle BOQ + \angle COQ + \angle COR + \angle DOR + \angle DOS + \angle AOS = 360^\circ$$

[Complete angle around a point]

$$\Rightarrow 2\angle 1 + 2\angle 2 + 2\angle 3 + 2\angle 4 = 360^\circ$$

$$\Rightarrow \angle 1 + \angle 2 + \angle 3 + \angle 4 = 180^\circ$$

$$\Rightarrow (\angle 1 + \angle 2) + (\angle 3 + \angle 4) = 180^\circ$$

$$\Rightarrow \angle AOB + \angle COD = 180^\circ$$

$$\text{Or } (\angle 1 + \angle 4) + (\angle 2 + \angle 3) = 180^\circ$$

$$\Rightarrow \angle AOD + \angle BOC = 180^\circ$$

Hence, proved!

P.T.O.

Calculating mean using step deviation method.

| No. of days<br>(class interval) | No. of students<br>( $f_i$ ) | $x_i$<br>( $\frac{\text{Upper} + \text{Lower limit}}{2}$ ) | $u_i = \frac{x_i - A}{h}$ | $f_i \times u_i$       |
|---------------------------------|------------------------------|------------------------------------------------------------|---------------------------|------------------------|
| 0-6                             | 10                           | 3                                                          | $\frac{3-21}{6} = -3$     | $10 \times -3 = -30$   |
| 6-12                            | 11                           | 9                                                          | $\frac{9-21}{6} = -2$     | $11 \times -2 = -22$   |
| 12-18                           | 7                            | 15                                                         | $\frac{15-21}{6} = -1$    | $7 \times -1 = -7$     |
| 18-24                           | 4                            | 21                                                         | $\frac{21-21}{6} = 0$     | $4 \times 0 = 0$       |
| 24-30                           | 4                            | 27                                                         | $\frac{27-21}{6} = 1$     | $4 \times 1 = 4$       |
| 30-36                           | 3                            | 33                                                         | $\frac{33-21}{6} = 2$     | $3 \times 2 = 6$       |
| 36-42                           | 1                            | 39                                                         | $\frac{39-21}{6} = 3$     | $1 \times 3 = 3$       |
| Total :                         | $\Sigma f_i = 40$            |                                                            |                           | $\Sigma f_i u_i = -46$ |

Class size ( $h$ ) =  $6 - 0 = 6$

Assumed mean ( $A$ ) = 21

Mean =  $A + \frac{\Sigma f_i u_i \times h}{\Sigma f_i}$

$$\Rightarrow \text{Mean} = 21 + \left( \frac{-46}{40} \right) \times 6$$

$$= 21 - \frac{23 \times 6}{10}$$

$$= 21 - 6.9$$

$$= 14.1 \text{ days}$$

Mean of no. of days student remains absent = 14.1 days

16. Given, each ripper blade has length ( $r$ ) = 21 cm.  
and sweeps angle =  $120^\circ$

Area swept by one blade =  $\frac{\theta}{360} \times \pi r^2$  unit square

$$= \frac{120}{360} \times \frac{22}{7} \times 21 \times 21 \text{ cm}^2$$

$$= 462 \text{ cm}^2$$

Blades don't overlap.

$\therefore$  Area swept by 2 blades =  $462 \times 2 \text{ cm}^2 = 924 \text{ cm}^2$


17.

Given,

$$\triangle ABC \sim \triangle PQR$$

AD and PM

are medians of  $\triangle ABC$  and  $\triangle PQR$  respectively.



Since,  $\triangle ABC \sim \triangle PQR$

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR} \quad \text{--- (1)}$$

D is the midpoint of BC (AD is median)  
M is the midpoint of QR (PM is median)

$$\therefore BC = 2BD$$

$$QR = 2QM$$

$$\therefore \frac{AB}{PQ} = \frac{BC}{QR}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{2BD}{2QM}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{BD}{QM}$$

$\Rightarrow$

$$\therefore \triangle ABD \sim \triangle PQM \quad \text{--- (2)}$$

That is,  $\frac{AB}{PQ} = \frac{BD}{QM} = \frac{AD}{PM}$

Similarly,  $\frac{BC}{QR} = \frac{AC}{PR}$  [from ①]

$$\Rightarrow \frac{2BD}{2QM} = \frac{AC}{PR}$$

$$\Rightarrow \frac{AC}{PR} = \frac{BD}{QM}$$

$$\therefore \triangle ACD \sim \triangle PRM$$

From both ③ and ⑤ we get that.

$$\frac{AB}{PQ} = \frac{AD}{PM}$$

hence proved!

18.

Given:  $P(x) = x^5 - 4x^3 + x^2 + 3x + 1$ .

$$Q(x) = x^3 - 3x + 1$$

To check: if  $Q(x)$  is a factor of  $P(x)$  or not.

Method: simply divide.

P.T.O.

$$\begin{array}{r} x^3 - 3x + 1 \quad x^5 - 4x^3 + 3x + 1 \quad (x^2 - 1) \\ \underline{x^5 - 3x^3 + x^2} \phantom{+ 1} \\ -x^3 + 3x + 1 \end{array}$$

$$\begin{array}{r} -x^3 + 3x + 1 \\ \underline{-x^3 + 3x - 1} \\ + \phantom{-} + \phantom{-} + \phantom{-} \\ 2 \end{array}$$

We get remainder = 2

Therefore,  $p(x)$  is not completely divisible by  $q(x)$

$q(x)$  is not a factor of  $p(x)$

19. Let us assume, if possible, that  $\sqrt{3}$  is rational  
Then,  $\sqrt{3}$  can be expressed as  $\frac{p}{q}$  where  $q \neq 0$  and  $p, q$  are coprimes [ $\text{HCF}(p, q) = 1$ ]

$$\therefore \sqrt{3} = \frac{p}{q} \quad [p, q \in \mathbb{Z}; \text{HCF}(p, q) = 1]$$

On squaring both sides,

$$3 = \frac{p^2}{q^2}$$

$$\Rightarrow p^2 = 3q^2 \quad \text{--- (1)}$$

$\therefore$  3 divides  $p^2$   
 $\therefore$  3 divides  $p$ .

Then,  $p$  can be written as;

$$p = 3a \quad \text{for some integer 'a'}$$

On squaring,

$$p^2 = 9a^2$$

$$\text{Put } p = 3a \text{ from (1)}$$

$$\Rightarrow 3q^2 = 9a^2$$

$$\Rightarrow q^2 = 3a^2$$

$$3 \text{ divides } q^2$$

$$\therefore 3 \text{ divides } q$$

$\therefore$  3 divides both  $p$  and  $q$ , 3 is common factor of  $p$  and  $q$ .

But,  $p$  and  $q$  are co-primes.

Therefore, our assumption is wrong.  
This is irrational.

from Section D (4 marks)

25. More than series

More than series  
More than 20 - 100

More than 30 - 90

More than 40 - 82

More than 50 - 70

More than 60 - 46

More than 70 - 40

More than 80 - 15

More than 90 - 0

| Class Interval | 20-30 | 30-40 | 40-50 | 50-60 | 60-70 | 70-80 | 80-90 |
|----------------|-------|-------|-------|-------|-------|-------|-------|
| Frequency      | 10    | 8     | 12    | 24    | 6     | 25    | 15    |


$$\sum f_i = 100$$

$$n = 100 \quad n = 50$$

For more than ogive, we plot the point A (20, 100), B (30, 90), C (40, 82), D (50, 70), E (60, 46), F (70, 40), G (80, 15) and H (90, 0).

**www.FirstRanker.com**








Q6.

Given: Triangle ABC with

$$A(x_1, y_1) = A(1, -4)$$

D and E are midpoints of AB and AC.



Let coordinates of B be  $(x_2, y_2)$  and that of C be  $(x_3, y_3)$ .  
Using section formula for mid-point;

$$\frac{1+x_2}{2} = 2, \quad \frac{-4+y_2}{2} = -1$$

$$\Rightarrow x_2 = 3, \quad y_2 = 2$$

$$(x_2, y_2) = (3, 2)$$

Similarly,  $\frac{1+x_3}{2} = 0, \quad \frac{-4+y_3}{2} = -1$

$$\Rightarrow x_3 = -1, \quad y_3 = 2$$

$$(x_3, y_3) = (-1, 2)$$

$$\text{Area of triangle} = \frac{1}{2} |x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)| \text{ unit square}$$

$$= \frac{1}{2} |1(-2-2) + 3(2+4) + (-1)(-4-2)| \text{ unit square}$$

$$= \frac{1}{2} |1 \times 0 + 3 \times 6 + (-1 \times -6)| \text{ sq. units}$$

$$= \frac{1}{2} |18 + 6| \text{ sq. units} = 12 \text{ sq. units}$$

$$\therefore \text{Area of } \Delta = 12 \text{ sq. units}$$

21. Let the required two numbers be  $5x$  and  $6x$ .  
Given, if 7 is subtracted from both nos, ratio becomes 4:5.  
New nos. =  $(5x-7)$  and  $(6x-7)$ .

According to the question;

$$\frac{5x-7}{6x-7} = \frac{4}{5} \quad \text{--- (1)}$$

Solving ①;

$$5(5x-7) = 4(6x-7)$$

$$\Rightarrow 25x - 35 = 24x - 28$$

$$\Rightarrow x = 35 - 28$$

$$\Rightarrow x = 7$$

The required nos. are :  ~~$5x = 35$~~   
 ~~$6x = 42$~~

Q2.

Let the radius of cylindrical

pipe be  $r$  metres


given -

Radius ( $R$ ) of cylindrical tank = 40 cm

$$= \frac{2}{5} \text{ m}$$

Height of tank filled = 3.15 m

Time taken =  $\frac{1}{2}$  h = 30 minutes =  $30 \times 60$  s.



Rate of flow of water =  $2.52 \text{ km/h} = \frac{2.52 \times 1000}{60} \text{ m/s} = \frac{126}{10} \text{ m/s}$

$$= 0.7 \text{ m/s.}$$

Volume of

To find - internal diameter of pipe ( $2r$ ).

Solution:

Volume of water passed through pipe  
in  $\frac{1}{2}$  hour =  $\pi r^2 \times h$  unit cube

$$= \pi r^2 \times \text{rate of flow} \times \text{time}$$

$$= \pi r^2 \times 0.7 \times 30 \times 60 \text{ m}^3.$$

Volume of water in tank in  $\frac{1}{2}$  hour =  $\pi R^2 \times h$

$$= \pi \left(\frac{2}{5}\right)^2 \times 3.15 \text{ m}^3$$

But, volume of water passed through pipe = volume of water collected in tank

$$\therefore \pi r^2 \times \frac{7}{18} \times 30 \times 60 = \pi \left(\frac{2}{5}\right)^2 \times \frac{315}{100}$$

$$\Rightarrow r^2 = \frac{4}{5 \times 8} \times \frac{315}{100} \times \frac{1}{7} \times \frac{1}{3} \times \frac{1}{60} \times \frac{60 \times 205}{205}$$

$$\Rightarrow r^2 = \frac{1}{2500} \Rightarrow r = \sqrt{\frac{1}{2500}} \Rightarrow r = \pm \frac{1}{50}$$

Radius is <sup>always</sup> positive, so  $r = -\frac{1}{50}$  can be ignored.

$$\Rightarrow r = \frac{1}{50} \text{ m.}$$

$$\Rightarrow r = \frac{1}{50} \times 100 \text{ cm} \Rightarrow r = 2 \text{ cm}$$

Internal diameter of pipe =  $d = 2r = 4 \text{ cm}$   
Or  $0.04 \text{ m.}$

### Section - B.

7. Event: Dice is thrown.

Outcomes: 1, 2, 3, 4, 5, 6 (6 outcomes)

Favourable events :-

i) Composite number = 4, 6

Probability of getting a composite no. =  $\frac{\text{no. of favourable outcomes}}{\text{Total outcomes}}$

$$= \frac{2}{6} = \frac{1}{3}$$

ii)

prime no. : 2, 3, 5.

Probability =  $\frac{\text{no. of outcomes favourable to the event}}{\text{Total possible outcomes}}$

or  $\frac{\text{no. of prime nos.}}{\text{Total outcomes}}$

$$= \frac{3}{6} = \frac{1}{2}$$

$$\frac{40}{34}$$

8. Event : cards numbered from 7-40 are chosen  
Total possible outcomes : (7, 8, 9, ..., 40) = 34 cards.


For Favourable event : cards multiple of 7.

Favourable outcomes : (7, 14, 21, 28, 35) 5 cards

Probability of selecting a card multiple of 7 =  $\frac{\text{Favourable outcomes}}{\text{Total no. of outcomes}}$

$$= \frac{5}{34}$$

9. Points A, B, C, D are vertices of a parallelogram.



We know that diagonals of a parallelogram bisect each other.

$\therefore$  O is the midpoint of both AC and BD.

Using section formula for mid-point.

on BD,  $x = \frac{4+5}{2}, y = \frac{3+1}{2}$

$\Rightarrow x = \frac{9}{2}, y = 2.$

on AC

$x = \frac{3+a}{2}, y = \frac{b+1}{2}$

$\Rightarrow \frac{9}{2} = \frac{3+a}{2}, y = 2 = \frac{b+1}{2} \Rightarrow \boxed{a=6}, \boxed{b=3}$

$\Rightarrow a=6, b=3.$

34

10. given -

$$3x - 5y = 4 \quad \text{--- ①}$$

$$9x - 2y = 7 \quad \text{--- ②}$$

To find 'x' and 'y'

Multiplying ①  $\times 3$ , and ②  $\times 1$  and adding, we get =

$$(3x - 5y) \times 3 + (9x - 2y) \times 1 = 4 \times 3 + 7 \times 1$$

$$\Rightarrow 9x - 15y - 9x + 2y = 12 + 7$$

$$\Rightarrow -13y = 19 \Rightarrow y = -\frac{19}{13}$$

Then, putting  $y = -\frac{19}{13}$  in ①;

$$3x = 4 + 5y$$

$$\Rightarrow 3x = 4 + 5 \times -\frac{19}{13}$$

$$\Rightarrow 3x = \frac{52 - 95}{13} \Rightarrow x = -\frac{43}{13}$$

$$\Rightarrow x = -\frac{43}{13}$$

11. Using Euclid's Division Lemma (states that  $a = bq + r$ ,  $0 \leq r < b$ ) we can find the HCF of 65 and 117.

$$\begin{aligned} 117 &= 65 \times 1 + 52 \\ 65 &= 52 \times 1 + 13 \\ 52 &= 13 \times 4 + 0 \end{aligned}$$

$$\therefore \text{HCF of } 65, 117 = 13$$

But,

$$65n - 117 = 13$$

$$\Rightarrow 65n = 13 + 117$$

$$\Rightarrow n = \frac{130}{65}$$

$$n = 2$$

$$\Rightarrow$$

$$\boxed{n = 2}$$

12) Given, quadratic equation  $\Rightarrow kx^2 - 6x - 1 = 0$ , where  $a = k$ ,  $b = -6$ ,  $c = -1$ .

For no real roots (you imaginary roots), discriminant must be less than 0.

That is,  $D < 0$

$$\Rightarrow b^2 - 4ac < 0$$

$$\Rightarrow (-6)^2 - 4 \times k \times -1 < 0$$


$$\Rightarrow 36 + 4k < 0$$

$$\Rightarrow 4k < -36 \Rightarrow \boxed{k < -9}$$

$\therefore k$  should be less than -9. ( $k = -10, -11, \dots$ )

Section-A.

1) Given, 2 concentric circles



$$OP = OQ = a$$

$$OM = b$$

To find - PQ

$$PM = \sqrt{OP^2 - OM^2} \Rightarrow PM = \sqrt{a^2 - b^2}$$

$$PQ = 2PM \Rightarrow PQ = 2\sqrt{a^2 - b^2} \text{ units}$$

2.  $\tan \alpha = \frac{5}{12}$

36+41220  
12-9

Using identity;  $\sec^2 \alpha - \tan^2 \alpha = 1$

$$\sec^2 \alpha = 1 + \tan^2 \alpha$$

$$\Rightarrow \sec^2 \alpha = 1 + \left(\frac{5}{12}\right)^2$$

$$\Rightarrow 1 + \frac{25}{144}$$

$$= \frac{144+25}{144}$$

$$\Rightarrow \sec^2 \alpha = \frac{169}{144} \Rightarrow \sec \alpha = \sqrt{\frac{169}{144}}$$

$$\sec \alpha = \frac{13}{12}$$

$$\sec \alpha = \frac{13}{12}$$

3.  $(x+5)^2 = 2(5x-3)$

$\Rightarrow x^2 + 25 + 10x = 10x - 6$

$\Rightarrow x^2 + 31 = 0$

$a=1, b=0, c=19$

$\text{Discriminant} = b^2 - 4ac$

$= 0^2 - 4 \times 1 \times 19$

$= 0 - 76$

$= -76$

4.  $429$  can be expressed as -

$3(429)$   
 $11(143)$   
 $13(11)$

$429 = 3 \times 11 \times 13$

5. First 10 multiples of 6 form AP  $\rightarrow 6, 12, 18, \dots, 60$

Sum of 1st 10 multiples  $= \frac{n}{2} [a + e]$

$= \frac{10}{2} [6 + 60]$

$= 330$

6.

Given,  $A = 5, -3$   
 $B = 13, m$

$AB = 16$  units

Using distance formula;

$$\sqrt{(13-5)^2 + (m+3)^2} = 10$$

on squaring;

$$8^2 + (m+3)^2 = 100$$

$$\Rightarrow (m+3)^2 = 100 - 64$$

$$\Rightarrow \sqrt{(m+3)^2} = \sqrt{36}$$

$$\Rightarrow (m+3) = \pm 6$$

considering only positive value;

$$m = 6 - 3$$

$$\Rightarrow \boxed{m = 3}$$

Total

