

GUJARAT TECHNOLOGICAL UNIVERSITY

BE- SEMESTER– $1^{st}\,/\,2^{nd}$ (OLD) EXAMINATION – SUMMER 2018

Subject Code: 110009 Date: 17-05-2018

Subject Name: MATHEMATICS-II

Time: 02:30 pm to 05:30 pm Total Marks: 70

Instructions:

(b)

- 1. Attempt any five questions.
- 2. Make suitable assumptions wherever necessary.
- 3. Figures to the right indicate full marks.

Q.1 (a) (i) For which values of
$$k$$
, u and v orthogonal?
 $u = (2,1,3), v = (1,7,k)$

(ii) Verify Cauchy-Schwarz inequality for the vectors
$$u = (0, -2, 2, 1), v = (-1, -1, 1, 1)$$

(b)
$$\begin{bmatrix}
1 & 6 & 8 \\
2 & 5 & 3
\end{bmatrix}$$
(i) Find the rank for the matrix $\begin{bmatrix} 2 & 5 & 3 \\ 7 & 9 & 4 \end{bmatrix}$.

(ii) Solve the following linear system by using Gauss Jordan method.
$$3x - 2y + 8z = 9$$
$$-2x + 2y + z = 3$$
$$x + 2y - 3z = 8$$

Q.2 (a) (i) Solve the following linear system by using Gauss Elimination method
$$x + y + z = 6$$
$$x + 2y + 3z = 10$$

$$x + 2y + 4z = 1$$

[1 0 1]

O4

- (ii) Find the inverse of the matrix $\begin{vmatrix} -1 & 1 & 1 \\ 0 & 1 & 0 \end{vmatrix}$. (i) Prove that the matrix $A = \begin{bmatrix} -1 & 2+i & 5-3i \\ 2-i & 7 & 5i \\ 5+3i & -5i & 2 \end{bmatrix}$ is a hermitian matrix.
- [5+3i -5i 2]

 (ii) Find the eigenvalues and eigenvectors of the matrix $A = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \end{bmatrix}$
- (ii) Find the eigenvalues and eigenvectors of the matrix $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix}$ Q.3 (a) Show that the set of all 2×2 matrices of the form $\begin{bmatrix} a & 1 \\ 1 & b \end{bmatrix}$ with addition defined

by
$$\begin{bmatrix} a & 1 \\ 1 & b \end{bmatrix} + \begin{bmatrix} c & 1 \\ 1 & d \end{bmatrix} = \begin{bmatrix} a+c & 1 \\ 1 & b+d \end{bmatrix}$$
 and scalar multiplication defined by $k \begin{bmatrix} a & 1 \\ 1 & b \end{bmatrix} = \begin{bmatrix} ka & 1 \\ 1 & kb \end{bmatrix}$ is a vector space.

03



- First (b) ker's choice (i) Determine whether whow First Ranker a com 2,1) and www. First Ranker a com 03 dependent or linearly independent.
 - (ii) Determine whether the vectors (1,-1,1), (0,1,2), (3,0,-1) forms basis for \mathbb{R}^3
- **Q.4** (a) Extend the subset $A = \{(1, -2, 5, -3), (2, 3, 1, -4)\}$ of R^4 to the basis for vector space R^4 .
 - (b) (i) Find two vector in \mathbb{R}^2 with Euclidean norm whose inner product with (-3,1) is 2ero.
 - (ii) Obtain the matrix of a linear transformation $T: \mathbb{R}^3 \to \mathbb{R}^3$ defined by T(x, y, z) = (2x, x + y + z, x + 3z) with respect to the basis $B_1 = \{(1,0,0), (1,1,0), (1,1,1)\}$ and $B_2 = \{(1,0,0), (0,1,0), (0,0,1)\}$.
- **Q.5** (a) For the basis $S = \{u, v, w\}$ of R^3 , where u = (1,1,1), v = (1,1,0) and w = (1,0,0), let $T: R^3 \to R^3$ be a linear transformation such that T(u) = (2,-1,4), T(v) = (3,0,1), T(w) = (-1,5,1). Find a formula for T(x,y,z) and use it to find T(2,4,-1).
 - (b) State Rank-Nullity theorem. Let $T: R^4 \to R^3$ be a linear transformation defined by T(1,0,0,0) = (1,1,1), T(0,1,0,0) = (1,-1,1), T(0,0,1,0) = (1,0,0), T(0,0,0,1) = (1,0,1). Then verify the rank-nullity theorem.
- **Q.6** (a) Find the least square solution of the linear system AX = b given by $x_1 + x_2 = 7$ $-x_1 + x_2 = 0$ $x_1 + 2x_2 = -7$
 - (b) Let R^3 have the Euclidean inner product. Use the Gram Schmidt process to transform the basis $\{u_1, u_2, u_3\}$ into an orthonormal basis, where $u_1 = (1, 0, 0), u_2 = (3, 7, -2), u_3 = (0, 4, 1)$
- Q.7 (a)
 Find a matrix that diagonalizes and determine $P^{-1}AP$, where $A = \begin{bmatrix} 2 & 0 & -2 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$.
 - (b)
 (i) Find the algebraic and geometric multiplicity of $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -3 & 3 \end{bmatrix}$.
 - (ii) Verify Caley-Hamilton theorem for the matrix, $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$.
