

Subject Code: 2110015
Date: 17-05-2018
Subject Name: Vector Calculus and Linear Algebra
Time: 02:30 pm to 05:30 pm
Total Marks: 70
Instructions:

1. Question No. 1 is compulsory. Attempt any four out of remaining Six questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

Q.1 Objective Question (MCQ) Mark
(a) Choose the appropriate answer for the following questions. 07

1. A square matrix whose determinant is non zero is called
(A) Singular (B) non-singular (C) invertible (D) both B and C
2. If A and B are non singular matrices then $(AB)^{-1} = \text{_____}$
(A) $A^{-1}B^{-1}$ (B) AB (C) $B^{-1}A^{-1}$ (D) none of these
3. If $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$ then A is in
(A) Row echelon form (B) Reduced Row echelon form (C) both A and B (D) none of these
4. For what values of k does the system $x + y = 2, 3x + 3y = k$ has infinitely many solutions
(A) $k=5$ (B) $k=4$ (C) $k=6$ (D) $k=1$
5. If in a set of vectors atleast one member can be expressed as a linear combination of the remaining vectors then the set is
(A) Linearly independent (B) Linearly dependent (C) basis (D) none of these
6. If V is any vector space and S be a subset of V then S is called basis for V if
(A) S is Linearly independent (B) S spans V (C) both A and B
(D) S is Linearly dependent
7. For what value of k the vectors u and v are orthogonal where $u=(2,1,3)$, $v=(1,7, k)$
(A) $k=-3$ (B) $k=1$ (C) $k=5$ (D) $k=2$

(b) Choose the appropriate answer for the following questions. 07

1. The eigen values of a matrix $A = \begin{bmatrix} 1 & 0 \\ 2 & 4 \end{bmatrix}$ are
(A) 1,4 (B) -1,-4 (C) 1,3 (D) -1,3
2. If A is a nxn size invertible matrix then rank of A is
(A) $n-1$ (B) n (C) $2n$ (D) $n+1$

3. If F is solenoidal then www.FirstRanker.com www.FirstRanker.com
 (A) $\nabla F = 0$ (B) $\nabla \times F = 0$ (C) $\nabla \cdot F = 0$ (D) none of these
4. The mapping $T : R^3 \rightarrow R^3$ defined by $T(x, y, z) = (x, y, -z)$ is called as
 (A) Contraction (B) Projection (C) Reflection (D) Rotation
5. The linear transformation $T : V \rightarrow W$ is one to one if and only if the nullspace of T consists of only
 (A) Identity vector (B) zero vector (C) any non zero vector (D) none of these
6. If $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ then the rank of the matrix A is
 (A) 1 (B) 2 (C) 0 (D) 4
7. Let A be a skew-symmetric matrix then
 (A) $a_{ij} = a_{ji}$ (B) $a_{ij} = -a_{ji}$ (C) $a_{ii} = 0$ (D) both B and C

Q.2 (a) Find the unit vector normal to the surface $xy^3z^2 = 4$ at $(-1, -1, 2)$ **03**

(b) **04**
 Express the matrix $A = \begin{bmatrix} 3 & -2 & 6 \\ 2 & 7 & -1 \\ 5 & 4 & 0 \end{bmatrix}$ as the sum of a symmetric and skew-symmetric matrix.

(c) Investigate for what values of λ and μ the equations **07**
 $2x + 3y + 5z = 9, 7x + 3y - 2z = 8, 2x + 3y + \lambda z = \mu$ have
(1) No solution (2) a unique solution (3) infinite number of solutions

Q.3 (a) **03**
 Find the rank of the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$

(b) **04**
 Find the inverse of the matrix $\begin{bmatrix} 2 & 3 & 4 \\ 4 & 3 & 1 \\ 1 & 2 & 4 \end{bmatrix}$ by Gauss Jordan Method

(c) For the basis $S = \{v_1, v_2, v_3\}$ of R^3 where **07**
 $v_1 = (1, 1, 1), v_2 = (1, 1, 0), v_3 = (1, 0, 0)$ Let $T : R^3 \rightarrow R^2$ be the linear transformation such that
 $T(v_1) = (1, 0), T(v_2) = (2, -1), T(v_3) = (4, 3)$ find a formula for $T(x_1, x_2, x_3)$ and then use the formula to find $T(4, 3, -2)$

- Q.4 (a)** Determine whether the vector $v = (-5, 1, -7)$ is a linear combination of the vectors $v_1 = (1, -2, 2), v_2 = (0, 5, 5), v_3 = (2, 0, 8)$ **03**
- (b)** Solve the linear system $x + y + z = 4, -x - y + z = -2, 2x - y + 2z = 2$ by gauss elimination method. **04**
- (c)** Let R^3 have the Euclidean inner product. Use the gram schmidt process to transform the basis (u_1, u_2, u_3) in to Orthonormal basis where $u_1 = (1, 0, 0), u_2 = (3, 7, -2), u_3 = (0, 4, 1)$ **07**
- Q.5 (a)** Find the eigen values and corresponding eigen vectors of **03**
- $$A = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix}$$
- (b)** Let $A = \begin{bmatrix} -2 & 3 \\ 1 & -2 \\ 1 & -1 \end{bmatrix}$ and $b = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$ then find the least squares solutions to $AX=b$ **04**
- (c)** Let $T : R^3 \rightarrow R^3$ be a linear operator and $B = (v_1, v_2, v_3)$ a basis for R^3 . Suppose that $T(v_1) = (1, 1, 0), T(v_2) = (1, 0, -1), T(v_3) = (2, 1, -1)$ then **07**
- (1) Is $(1, 2, 1)$ in $R(T)$? (2) Find a basis for $R(T)$.**
- Q.6 (a)** Find the work done by the force $\vec{F} = (3x^2 - 3x)i + 3zj + k$ along the straight line $ti + tj + tk, 0 \leq t \leq 1$. **03**
- (b)** Check whether the vectors $(2, -3, 1), (4, 1, 1), (0, -7, 1)$ is a basis for R^3 **04**
- (c) Verify Green's Theorem for** $\vec{F} = (x - y)i + xj$, and C is $x^2 + y^2 = 1$ **07**
- Q.7 (a)** Find the directional derivative of $4xz^2 + x^2yz$ at $(1, -2, -1)$ in the direction of $2i - j - 2k$ **03**
- (b)** Show that $\vec{F} = (e^x \cos y + yz)i + (xz - e^x \sin y)j + (xy + z)k$ is conservative and find the potential function. **04**
- (c)** Let $V = \{(a, b) / a, b \in R\}$ and let $v = (v_1, v_2), w = (w_1, w_2)$ then define $(v_1, v_2) + (w_1, w_2) = (v_1 + w_1 + 1, v_2 + w_2 + 1)$ and $c(v_1, v_2) = (cv_1 + c - 1, cv_2 + c - 1)$ then verify that V is a vector space. **07**
