

www.FirstRanker.com Enwww.मtirstRanker.com GUJARAT TECHNOLOGICAL UNIVERSITY

BE - SEMESTER-V (OLD) - EXAMINATION - SUMMER 2018

Subject Code:150503 Date:30/04/2018 Subject Name: Chemical Engineering Thermodynamics-II Time:02:30 PM to 05:00 PM **Total Marks: 70 Instructions:** 1. Attempt all questions. 2. Make suitable assumptions wherever necessary. 3. Figures to the right indicate full marks. **Q.1** (a) Define partial molar properties. Discuss various methods for evaluation of partial **07** molar properties. List out various methods for evaluation of fugacity coefficient of pure component. **07 (b)** Discuss any two in detail. Explain Retrograde condensation in detail with figure. **07 Q.2** (a) Show that for a binary system, Henry's law is valid for component '1' then Lewis **07** Randall rule is valid for component '2'. **(b)** The enthalpy at 300 K and 1 bar of a binary liquid mixture is: **07** $\mathbf{H} = 400\mathbf{X}_1 + 600\mathbf{X}_2 + \mathbf{X}_1\mathbf{X}_2(40\mathbf{X}_1 + 20\mathbf{X}_2)$ where H is in J/mol for the stated temperature and pressure, determine: (i) Expression for partial molar enthalpies of both components 1 and 2, (ii) Numerical value for pure component enthalpies H_1 and H_2 , (iii) Numerical values for the partial molar enthalpies for both the components at infinite dilution. Discuss various methods for checking the consistency of experimental VLE data. **07** Q.3 (a) **(b)** Explain T-x,y diagram for partially miscible system. **07** At 300 K and 1 bar, the volumetric data for a liquid mixture of benzene and **07** Q.3 (a) cyclohexane are represented by: $V=109.4 \times 10^{-6} - 16.8 \times 10^{-6} \text{ x} - 2.64 \times 10^{-6} \text{ x}^2$ where x is the mole fraction of benzene and V has the units of m³/mol. Find expressions for the partial molar volumes of benzene and cyclohexane. **(b)** Construct P-x-y diagram for the cyclohexane (1) / benzene (2) system at 313 K. 07 Use the following expressions for the liquid-phase activity coefficients: $\ln \gamma_1 = 0.458 \text{ x}_2^2, \ln \gamma_2 = 0.458 \text{ x}_1^2$ At 313 K, $P_1^S = 24.62$ KPa and $P_2^S = 24.41$ KPa. 0.4 (a) Discuss the criteria of chemical reaction equilibrium. **07** Discuss about the effect of temperature on equilibrium constant. 07 Write short note on Van Laar equation and Margules equation. **07 Q.4** (a) Discuss minimum and maximum boiling azeotropes giving examples for each **07 (b)** with neat diagrams. In the synthesis of ammonia, stoichiometric amounts of nitrogen and hydrogen are 07 **Q.5** (a) sent to a reactor where the following reaction occurs. $N_2 + 3H_2 \rightarrow 2NH_3$ The equilibrium constant for the reaction at 675 K is 2 x 10⁻⁴. Determine the percent conversion of nitrogen to ammonia at 675 K & 20 bar. Prove that $\Delta G^0 = -RT \ln K$ **07 (b)**



$\begin{array}{c} \text{www.FirstRanker.com} \\ \text{OR} \end{array}$

www.FirstRanker.com

Q.5 (a) Write a brief note on Excess Properties.

07

(b) Discuss in detail: Ideal solutions and Non-ideal solutions.

07

MWW.FilestRanker.com