

R13

Code No: 113AA

JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD
B.Tech II Year I Semester Examinations, November - 2015
MATHEMATICS – II
(Common to CE, CHEM, MMT, AE, PTE, CEE)
Time: 3 Hours
Max. Marks: 75
Note: This question paper contains two parts A and B.

Part A is compulsory which carries 25 marks. Answer all questions in Part A.

Part B consists of 5 Units. Answer any one full question from each unit.

Each question carries 10 marks and may have a, b, c as sub questions.

PART- A
(25 Marks)

- 1.a) If ϕ satisfies Laplacian equation, show that $\Delta\phi$ is both solenoidal. [2M]
- b) Find curl $x^2i + yj + zk$. [3M]
- c) Write the Dirichlet's conditions for the existence of Fourier series of a function $f(x)$ in the interval $(\alpha, \alpha + 2\pi)$. [2M]
- d) State and prove linearity property of Fourier transforms. [3M]
- e) State interpolation. [2M]
- f) If the interval of difference is unity, Find $\Delta(x^2 + 2x)$. [3M]
- g) Under what conditions Gauss-Seidal method is applicable. [2M]
- h) Derive an iteration formula to find square root of a number n by Newton Raphson method. [3M]
- i) State merits and demerits of Runge-Kutta method. [2M]
- j)

| | | | | |
|---|---|-----|--------|------|
| x | 0 | 1 | 2 | 3 |
| y | 1 | 0.5 | 0.3333 | 0.25 |

 Evaluate $\int_0^3 \frac{dx}{1+x}$ by Trapezoidal rule. [3M]

PART-B
(50 Marks)

2. State and verify Stokes theorem for the function $\vec{f} = x^2i + xyj$ integrated round the square in the plane $z = 0$ whose sides are along the lines $x = 0 = y$, $x = a = y$. [10]

OR

3. Find the work done in moving in a particle in the force field $\vec{f} = 3x^2i + (2zx - y)j + zk$, along the curve defined by $x^2 = 4y, 3x^3 = 8z$ from $x = 0$ to $x = 2$. [10]

- 4.a) Expand the Fourier series expansion of the function

$$f(x) = 0, -\pi \leq x \leq 0$$

$$= \sin x, 0 \leq x \leq \pi$$

$$\text{and deduce } \frac{1}{1.3} - \frac{1}{3.5} + \frac{1}{5.7} - \dots = \frac{\pi - 2}{4}$$

- b) Obtain the half range cosine series for the function $f(x) = x^2$ when $0 < x < \pi$ and

$$\text{find the sum of the series } \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots \quad [5+5]$$

OR

- 5.a) State and prove Change of scale property of Fourier transforms.
- b) Find Fourier sine transform of $f(x) = \frac{1}{x(x^2 + a^2)}$ and hence deduce cosine transform of $\frac{1}{x^2 + a^2}$. [5+5]

- 6.a) Find the missing term in the following data:

| | | | | | |
|---|---|---|---|---|----|
| x | 0 | 1 | 2 | 3 | 4 |
| y | 1 | 3 | 9 | - | 81 |

Why this value is not equal to 3^3 . Explain.

- b) Fit a straight line to the following data by the method of least square. [5+5]

| | | | | | |
|---|---|-----|-----|-----|-----|
| x | 0 | 1 | 2 | 3 | 4 |
| y | 1 | 1.8 | 3.3 | 4.5 | 6.5 |

OR

- 7.a) Given $\sin 45^\circ = 0.7071$, $\sin 50^\circ = 0.7660$, $\sin 55^\circ = 0.8192$ and $\sin 60^\circ = 0.8660$. Find $\sin 52^\circ$ using Newton's interpolation formula.
- b) A curve passes through the points (0, 18), (1, 10), (3, -18) and (6, 90). Find the slope of the curve at $x = 2$. [5+5]

8. Find the inverse of the matrix $\begin{bmatrix} 2 & 3 & 1 \\ 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix}$ using LU decomposition method. [10]

OR

- 9.a) Find the root of $2x - \log x = 7$, correct to three places of decimal using iteration method.
- b) Find the root of the equation $x \log_{10}(x) = 1.2$ using False position method. [5+5]
10. Find the successive approximate solution of the differential equation $y' = y$, $y(0) = 1$ by Picard's method and compare it with exact solution. [10]

OR

11. Determine the largest Eigen value and the corresponding eigenvector of the matrix $\begin{bmatrix} 4 & 1 & 0 \\ 1 & 20 & 1 \\ 0 & 1 & 4 \end{bmatrix}$ to 3 correct decimal places using the power method. [10]

---ooOoo---