

### www.FirstRanker.com

www.FirstRanker.com

Code No: R161110

**R16** 

**SET - 1** 

## I B. Tech I Semester Supplementary Examinations, May/June - 2019 MATHEMATICS-II (NM&CV)

(Com to ECE, EIE, ECom E)

Max. Marks: 70 Time: 3 hours

Note: 1. Question paper consists of two parts (Part-A and Part-B)

2. Answering the question in **Part-A** is Compulsory

3. Answer any **FOUR** Questions from **Part-B** 

# PART -A

1. a) Show that  $f(z) = |z|^2$  is not analytic at any point. (2M)

b) Evaluate  $\int_0^{1+i} (x^2 + iy) dz$  along the paths (i) y = x(2M)

Classify the Singularity of f(z) = Cosec z at z = 0(2M)

Find the Residue of  $f(z) = \frac{z}{(z+1)(z-2)}$  at z = -1(2M)

Write the merits of modified Euler's method. (2M)

Write relation between E and  $\delta$ . (2M)

What is quadratic convergence? (2M)

Find analytic function whose Real part  $u(x, y) = x^3 - 3xy^2 + 3x^2 - 3y^2 + 2x + 1$ (7M)

b) Show that a analytic function f(z) = u + iv form an Orthogonal system. (7M)

3. a) find the Laurent's series of  $f(z) = \frac{z^2 - 1}{(z+2)(z+3)}$  for (7M)

(i) |z| > 3 (ii) 2 < |z| < 3b) Evaluate  $\int_c \frac{e^{2z}}{(z-1)(z-2)} dz$  where C: |z| = 3 using Cauchy's integral formula. (7M)

4. a) Evaluate  $\oint_C \frac{1+z}{z(2-z)} dz$  Where c :|z| = 1 using residue theorem. (7M)

b) Evaluate  $\int_{0}^{\infty} \frac{\sin mx}{x(x^2 + a^2)} dx$  using residue theorem. (7M)

By Taylor's method find y(0.4) given that  $\frac{dy}{dx} = 3x + y^2$ , y(0) = 1 (7M)

b) Apply RK method of fourth order to find y(1.2) given that  $y^1 = x^2 + y^2$ , y(1) = 1.5(7M)

1 of 2



### www.FirstRanker.com

### www.FirstRanker.com

Code No: R161110



SET - 1

6. a) Interpolate by means of Newton backward formula, the population of a town for (7M) the year 1985, given that.

year	1939	1949	1959	1969	1979	1989
population	12	15	20	27	39	52

b) Evaluate y(7) from the following table.

(7M)

X	1	3	5	6	8
Y	2	1.5	2.4	4	5.6

7. a) Find the Real root of  $x + \log_{10} x - 2 = 0$  using Newton Raphson method. (7M)

b) Find the Real root of tanx = x using False position method. (7M)

www.FirstRanker.com