

Code No: 07A30802

R07**Set No. 2**

II B.Tech I Semester Examinations, MAY 2011

CHEMICAL PROCESS CALCULATIONS

Chemical Engineering

Time: 3 hours

Max Marks: 80

Answer any FIVE Questions

All Questions carry equal marks

- What is solution pressure? Explain.
 - A solution of sodium nitrate in water contains 1 kg of NaNO_3 per 10 kg of water. Calculate the amount of ice formed in cooling 1000 kg of this solution to a temperature of -15°C . Concentration of saturated solution at -15°C is given by 6.2 kmol of NaNO_3 per 1000 kg of water. [4+12]
- A large chamber contains dry N_2 at 27°C and 101.3 kPa. water is injected in to the chamber. After saturation of the N_2 with water vapor, the temperature in the chamber is 27°C :

 - What is the pressure inside the chamber after saturation?
 - How many moles of H_2O per mole of N_2 are present in the saturated mixture? [8+8]
- Ammonia is made by the reaction between hydrogen and N_2 according to the reaction

$$\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$$

If the reaction is carried out at 50 bar and 600 K, What volumes of nitrogen and H_2 at these conditions are theoretically required for producing 1000 kg of ammonia and what will be the volume of NH_3 produced at the reactor conditions. [16]
- Dry air at 295K and 100 kPa is bubbled through benzene. If the saturated air leaves at 300K and 100 kPa; how many kilograms of benzene are evaporated per 100m^3 of entering air. The vapour pressure of benzene (kPa.) is given by: [16]

$$\ln P^* = 13.885 - \frac{2788.51}{T - 52.36}$$
- Calculate the amount of heat needed to raise the temperature of 1 mol of a gaseous mixture containing 79% N_2 and 21% O_2 by volume from 200 to 1200°C .

$$C_p \text{ for } \text{N}_2 = 6.457 + 1.389 \times 10^{-3} T - 0.069 \times 10^{-6} T^2$$

$$C_p \text{ for } \text{O}_2 = 6.117 + 3.167 \times 10^{-3} T - 1.005 \times 10^{-6} T^2$$

Here C_p is in Cal/(mol.K) and T is in K. [16]
- A solution has a gravity of 100° Twaddell. Calculate its specific gravity and its gravity in degrees Baume.
 - An oil has a specific gravity at $60^\circ / 60^\circ\text{F}$ of 0.790. Calculate its gravity in degrees API and degrees Baume. [8+8]

Code No: 07A30802

R07**Set No. 2**

7. Molten sulfur at 140°C is burnt with 30% excess air at 120°C in a sulfur burner producing sulfur dioxide. Calculate the adiabatic reaction temperature using the following data:

Standard heat of reaction = -70.96 kcal/mol

Mean heat capacities in cal/mol.K :

$\text{S(l)} = 7.4$

$\text{SO}_2(\text{g}) = 12.15$

$\text{O}_2(\text{g}) = 7.14$

$\text{N}_2(\text{g}) = 7.51$

Air = 7.00

[16]

8. (a) To prepare a solution of 50% sulfuric acid, a dilute waste acid containing 25% H_2SO_4 is combined with a concentrated acid containing 90% H_2SO_4 . How many kg of the concentrated acid must be purchased for each 100 kg of dilute waste acid?
- (b) A mixed acid containing 65% H_2SO_4 , 20% HNO_3 and 15% H_2O is to be made by blending the following:
- A spent acid containing 10% HNO_3 , 60% H_2SO_4 and 30% H_2O .
 - A concentrated nitric acid containing 92% HNO_3 and 8% H_2O .
 - A concentrated sulfuric acid containing 95% H_2SO_4 and 5% H_2O .
- All percentages are on weight basis. How many kg of each acid must be used to obtain 1700 kg of the mixed acid?

[8+8]

Code No: 07A30802

R07**Set No. 4****II B.Tech I Semester Examinations, MAY 2011****CHEMICAL PROCESS CALCULATIONS****Chemical Engineering****Time: 3 hours****Max Marks: 80****Answer any FIVE Questions****All Questions carry equal marks**

- (a) To prepare a solution of 50% sulfuric acid, a dilute waste acid containing 25% H_2SO_4 is combined with a concentrated acid containing 90% H_2SO_4 . How many kg of the concentrated acid must be purchased for each 100 kg of dilute waste acid?

(b) A mixed acid containing 65% H_2SO_4 , 20% HNO_3 and 15% H_2O is to be made by blending the following:

 - A spent acid containing 10% HNO_3 , 60% H_2SO_4 and 30% H_2O .
 - A concentrated nitric acid containing 92% HNO_3 and 8% H_2O .
 - A concentrated sulfuric acid containing 95% H_2SO_4 and 5% H_2O .

All percentages are on weight basis. How many kg of each acid must be used to obtain 1700 kg of the mixed acid? [8+8]
- Molten sulfur at 140°C is burnt with 30% excess air at 120°C in a sulfur burner producing sulfur dioxide. Calculate the adiabatic reaction temperature using the following data:

Standard heat of reaction = -70.96 kcal/mol
Mean heat capacities in cal/mol.K :

$\text{S(l)} = 7.4$
 $\text{SO}_2(\text{g}) = 12.15$
 $\text{O}_2(\text{g}) = 7.14$
 $\text{N}_2(\text{g}) = 7.51$
Air = 7.00 [16]
- Dry air at 295K and 100 kPa is bubbled through benzene. If the saturated air leaves at 300K and 100 kPa; how many kilograms of benzene are evaporated per 100m^3 of entering air. The vapour pressure of benzene (kPa.) is given by: [16]

$$\ln P^* = 13.885 - \frac{2788.51}{T - 52.36}$$
- (a) A solution has a gravity of 100° Twaddell. Calculate its specific gravity and its gravity in degrees Baume.

(b) An oil has a specific gravity at $60^\circ / 60^\circ\text{F}$ of 0.790. Calculate its gravity in degrees API and degrees Baume. [8+8]
- Calculate the amount of heat needed to raise the temperature of 1 mol of a gaseous mixture containing 79% N_2 and 21% O_2 by volume from 200 to 1200°C .

C_p for $\text{N}_2 = 6.457 + 1.389 \times 10^{-3} T - 0.069 \times 10^{-6} T^2$
 C_p for $\text{O}_2 = 6.117 + 3.167 \times 10^{-3} T - 1.005 \times 10^{-6} T^2$
Here C_p is in $\text{Cal}/(\text{mol.K})$ and T is in K. [16]

Code No: 07A30802

R07**Set No. 4**

6. (a) What is solution pressure? Explain.
 (b) A solution of sodium nitrate in water contains 1 kg of NaNO_3 per 10 kg of water. Calculate the amount of ice formed in cooling 1000 kg of this solution to a temperature of -15°C . Concentration of saturated solution at -15°C is given by 6.2 kmol of NaNO_3 per 1000 kg of water. [4+12]
7. A large chamber contains dry N_2 at 27°C and 101.3 kPa. water is injected in to the chamber. After saturation of the N_2 with water vapor, the temperature in the chamber is 27°C :
 (a) What is the pressure inside the chamber after saturation?
 (b) How many moles of H_2O per mole of N_2 are present in the saturated mixture? [8+8]
8. Ammonia is made by the reaction between hydrogen and N_2 according to the reaction

$$\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$$

 If the reaction is carried out at 50 bar and 600 K, What volumes of nitrogen and H_2 at these conditions are theoretically required for producing 1000 kg of ammonia and what will be the volume of NH_3 produced at the reactor conditions. [16]

Code No: 07A30802

R07**Set No. 1****II B.Tech I Semester Examinations, MAY 2011****CHEMICAL PROCESS CALCULATIONS****Chemical Engineering****Time: 3 hours****Max Marks: 80****Answer any FIVE Questions****All Questions carry equal marks**

1. A large chamber contains dry N_2 at $27^\circ C$ and 101.3 kPa. water is injected in to the chamber. After saturation of the N_2 with water vapor, the temperature in the chamber is $27^\circ C$:
 - (a) What is the pressure inside the chamber after saturation?
 - (b) How many moles of H_2O per mole of N_2 are present in the saturated mixture? [8+8]
2. Ammonia is made by the reaction between hydrogen and N_2 according to the reaction

$$N_2 + 3H_2 \rightarrow 2NH_3$$
 If the reaction is carried out at 50 bar and 600 K, What volumes of nitrogen and H_2 at these conditions are theoretically required for producing 1000 kg of ammonia and what will be the volume of NH_3 produced at the reactor conditions. [16]
3. Molten sulfur at $140^\circ C$ is burnt with 30% excess air at $120^\circ C$ in a sulfur burner producing sulfur dioxide. Calculate the adiabatic reaction temperature using the following data:
 Standard heat of reaction = -70.96 kcal/mol
 Mean heat capacities in cal/mol.K :
 $S(l) = 7.4$
 $SO_2(g) = 12.15$
 $O_2(g) = 7.14$
 $N_2(g) = 7.51$
 Air = 7.00 [16]
4. Calculate the amount of heat needed to raise the temperature of 1 mol of a gaseous mixture containing 79% N_2 and 21% O_2 by volume from 200 to $1200^\circ C$.
 C_p for $N_2 = 6.457 + 1.389 \times 10^{-3} T - 0.069 \times 10^{-6} T^2$
 C_p for $O_2 = 6.117 + 3.167 \times 10^{-3} T - 1.005 \times 10^{-6} T^2$
 Here C_p is in $\text{Cal}/(\text{mol.K})$ and T is in K. [16]
5. (a) To prepare a solution of 50% sulfuric acid, a dilute waste acid containing 25% H_2SO_4 is combined with a concentrated acid containing 90% H_2SO_4 . How many kg of the concentrated acid must be purchased for each 100 kg of dilute waste acid?
 (b) A mixed acid containing 65% H_2SO_4 , 20% HNO_3 and 15% H_2O is to be made by blending the following:

Code No: 07A30802

R07**Set No. 1**

- i. A spent acid containing 10% HNO_3 , 60% H_2SO_4 and 30% H_2O .
- ii. A concentrated nitric acid containing 92% HNO_3 and 8% H_2O .
- iii. A concentrated sulfuric acid containing 95% H_2SO_4 and 5% H_2O .

All percentages are on weight basis. How many kg of each acid must be used to obtain 1700 kg of the mixed acid? [8+8]

6. (a) What is solution pressure? Explain.
- (b) A solution of sodium nitrate in water contains 1 kg of NaNO_3 per 10 kg of water. Calculate the amount of ice formed in cooling 1000 kg of this solution to a temperature of -15°C . Concentration of saturated solution at -15°C is given by 6.2 kmol of NaNO_3 per 1000 kg of water. [4+12]
7. Dry air at 295K and 100 kPa is bubbled through benzene. If the saturated air leaves at 300K and 100 kPa; how many kilograms of benzene are evaporated per 100m^3 of entering air. The vapour pressure of benzene (kPa.) is given by: [16]
- $$\ln P^* = 13.885 - \frac{2788.51}{T - 52.36}$$
8. (a) A solution has a gravity of 100° Twaddell. Calculate its specific gravity and its gravity in degrees Baume.
- (b) An oil has a specific gravity at $60^\circ / 60^\circ\text{F}$ of 0.790. Calculate its gravity in degrees API and degrees Baume. [8+8]


Code No: 07A30802

R07**Set No. 3**

II B.Tech I Semester Examinations, MAY 2011

CHEMICAL PROCESS CALCULATIONS**Chemical Engineering****Time: 3 hours****Max Marks: 80****Answer any FIVE Questions****All Questions carry equal marks**

1. Ammonia is made by the reaction between hydrogen and N_2 according to the reaction



If the reaction is carried out at 50 bar and 600 K, What volumes of nitrogen and H_2 at these conditions are theoretically required for producing 1000 kg of ammonia and what will be the volume of NH_3 produced at the reactor conditions. [16]

2. (a) What is solution pressure? Explain.

(b) A solution of sodium nitrate in water contains 1 kg of $NaNO_3$ per 10 kg of water. Calculate the amount of ice formed in cooling 1000 kg of this solution to a temperature of $-15^\circ C$. Concentration of saturated solution at $-15^\circ C$ is given by 6.2 kmol of $NaNO_3$ per 1000 kg of water. [4+12]

3. Molten sulfur at $140^\circ C$ is burnt with 30% excess air at $120^\circ C$ in a sulfur burner producing sulfur dioxide. Calculate the adiabatic reaction temperature using the following data:

Standard heat of reaction = -70.96 kcal/mol

Mean heat capacities in cal/mol.K :

$$S(l) = 7.4$$

$$SO_2(g) = 12.15$$

$$O_2(g) = 7.14$$

$$N_2(g) = 7.51$$

$$\text{Air} = 7.00$$

[16]

4. Calculate the amount of heat needed to raise the temperature of 1 mol of a gaseous mixture containing 79% N_2 and 21% O_2 by volume from 200 to $1200^\circ C$.

$$C_p \text{ for } N_2 = 6.457 + 1.389 \times 10^{-3} T - 0.069 \times 10^{-6} T^2$$

$$C_p \text{ for } O_2 = 6.117 + 3.167 \times 10^{-3} T - 1.005 \times 10^{-6} T^2$$

Here C_p is in Cal/(mol.K) and T is in K.

[16]

5. (a) A solution has a gravity of 100° Twaddell. Calculate its specific gravity and its gravity in degrees Baume.

(b) An oil has a specific gravity at $60^\circ / 60^\circ F$ of 0.790. Calculate its gravity in degrees API and degrees Baume. [8+8]

6. Dry air at 295K and 100 kPa is bubbled through benzene. If the saturated air leaves at 300K and 100 kPa; how many kilograms of benzene are evaporated per $100m^3$ of entering air. The vapour pressure of benzene (kPa.) is given by: [16]

Code No: 07A30802

R07**Set No. 3**

$$\ln P^* = 13.885 - \frac{2788.51}{T - 52.36}$$

7. (a) To prepare a solution of 50% sulfuric acid, a dilute waste acid containing 25% H_2SO_4 is combined with a concentrated acid containing 90% H_2SO_4 . How many kg of the concentrated acid must be purchased for each 100 kg of dilute waste acid?
- (b) A mixed acid containing 65% H_2SO_4 , 20% HNO_3 and 15% H_2O is to be made by blending the following:
- A spent acid containing 10% HNO_3 , 60% H_2SO_4 and 30% H_2O .
 - A concentrated nitric acid containing 92% HNO_3 and 8% H_2O .
 - A concentrated sulfuric acid containing 95% H_2SO_4 and 5% H_2O .
- All percentages are on weight basis. How many kg of each acid must be used to obtain 1700 kg of the mixed acid? [8+8]
8. A large chamber contains dry N_2 at 27 °C and 101.3 kPa. water is injected in to the chamber. After saturation of the N_2 with water vapor, the temperature in the chamber is 27 °C:
- What is the pressure inside the chamber after saturation?
 - How many moles of H_2O per mole of N_2 are present in the saturated mixture? [8+8]
